


Cornell University Library

The original of this book is in
the Cornell University Library.

There are no known copyright restrictions in
the United States on the use of the text.


ALBERT R. MANN LIBRARY
CORNELL UNIVERSITY



GIFT FROM THE LIBRARY OF
DOC AND KATY ABRAHAM,
THE GREEN THUMB,
NAPLES, N.Y.

G. M. Patchin.

Weyland
217



HARVEST TIME.

THE
FARMERS' AND MECHANICS'
MANUAL.

WITH MANY VALUABLE TABLES FOR MACHINISTS, MANUFACTURERS, MERCHANTS, BUILDERS, ENGINEERS, MASONS, PAINTERS, PLUMBERS, GARDENERS, ACCOUNTANTS, ETC.,

BY W. S. COURTNEY.

REVISED AND ENLARGED

BY GEORGE E. WARING, JR.,

AUTHOR OF "ELEMENTS OF AGRICULTURE," "DRAINING FOR PROFIT AND FOR HEALTH,"
"EARTH CLOSETS: HOW TO MAKE AND HOW TO USE THEM," AND FORMERLY
AGRICULTURAL ENGINEER OF THE CENTRAL PARK, NEW YORK.

TWO HUNDRED ILLUSTRATIONS.

SOLD ONLY BY SUBSCRIPTION.

NEW YORK:

E. B. TREAT & CO., 654 BROADWAY;

TREAT & LILLEY, CHICAGO; E. HANNAFORD & CO., CINCINNATI; A. H. HUBBARD, PHILA.; A. L. TALCOTT & CO., PITTSBURG; G. P. HAWKES & CO., BOSTON; H. H. BANCROFT & CO., SAN FRANCISCO; J. H. HUMMEL, NEW ORLEANS; J. C. DERBY, AUGUSTA, GA.

• MANN
SPEC.
COLL.

S
501
C86
1868

Entered according to Act of Congress, in the year 1868, by
E. B. TREAT & Co.,
In the Clerk's Office of the District Court of the United States for the Southern
District of New York.

PREFACE.

THERE are few persons, no matter what their calling or their education, who do not occasionally find themselves at a loss for information of the commonest kind, on any of the subjects pertaining to the practical arts of daily life—knowledge which was, perhaps, familiar to them in their school-boy days, but which has been forgotten or become obscured through the lapse of years. For example, how few persons can tell, without consulting books, the cubic inches contained in a bushel, the square yards in an acre, or how to measure the contents of a corn crib, or gauge a cistern. Nor is the inability to do so any reflection upon either their native capacity or their education. It is simply impossible to carry all these things in the memory so as to apply them when occasion requires. Hence the necessity for “Hand-Books,” “Mechanics’ Assistants,” “Pocket Companions,” &c.

Besides the labor involved in the almost daily necessity of calculating arithmetical, mensural, and other results, and the constant liability to error to which even the competent scholar is subject, the *time* required in the process, in this age, when time has emphatically acquired a money value, is no inconsiderable desideratum. Hence the necessity for “Ready Reckoners,” “Pocket Accountants,” “Calculators’ Assistants,” &c.

In presenting this volume, a chief aim of the author was so to combine the Manual with the Reckoner, as to furnish the inquirer, in brief, with all the necessary rules and *data* and the elementary facts and axioms relating to almost every branch of industrial science, and particularly that of agriculture, and, at the same time, whenever it was possible, to compute and tabulate the results for him in the same connection. Hence he will find in the ensuing pages the axiomatical or elementary propositions, the *data*, the standards, the units, &c., of almost every useful and practical art with which the farmer may have to deal, clearly stated, together with their simplest rules, illustrated by examples and solutions, and, wherever it was practicable, the arithmetical results calculated and tabularized.

Those who consult this book must remember that it is not a work of *recipes*, *prescriptions*, or of *directions* and *advice* as to the best mode of conducting any or all the various operations pertaining to agriculture, &c. But they will bear in mind that the subjects of which this book treats are, for the most part, *facts and figures*—assured analyses and demonstrations—about which there can be no dispute. The design was to produce a work of substantial and enduring value, and of universal application and use—something in the sphere of agriculture corresponding to Haswell in Engineering, or Fairbairn in Mechanics. How far the author's labors have tended to that end remains to be tested by experience. He is sanguine of their ultimate fruition.

So vast is the domain of agriculture, that there are few of the mechanic arts of which the farmer does not require some information, and which he is often compelled to seek through many books and journals. He is, in a certain sense, encyclopediac in his science and use. Hence many subjects

upon which he may require elementary knowledge and the assistance of computations may have escaped the vigilance of the author.

When a friend first suggested to the author the design of such a work, the latter had no adequate conception of the labor involved in such an undertaking. Although many of the tables were supplied or compiled from other authors, yet the labor involved in those he himself calculated and arranged was prodigious. Besides, the composition or type-setting of the matter was of the most tedious, difficult, and expensive kind, so that the volume of matter included within the covers would seem to bear no just proportion to the price the publisher is obliged to charge for it. Books much larger, and of many more pages of the ordinary composition, can be afforded at a much less cost. Withal, however, the author commends it to the favorable regard of those to whom it is addressed.

TO THE PRACTICAL READER.

HAVING been long engaged in the various occupations into which a life of combined farming and engineering is quite sure to lead any man of a practical turn of mind, I look back with regret on the days wasted in making long calculations to decide some simple question of size, or form, or quantity. Many a long day have I hunted through alcoves full of practical hand-books at the Astor Library,—scouring now the field of Agriculture, now of Mechanics, and now of Hydraulics,—often disappointed in my search, and compelled to go home and work far into the night, pursuing, through the long lanes of square and cube roots, the phantom of some every-day question of the discharge of water through pipes, the strength of material, or the resistance in ploughing.

I have always found less assistance than I had a right to expect from works written with the professed object of telling me what I wanted to know. After hunting them through, I have generally come to the conclusion that they contain almost everything *except* what I am looking for.

Certainly all that I have hitherto seen have been sadly incomplete.

Finally, I quite accidentally became acquainted with Mr. Courtney's Manual, and I found it much more nearly what it professes to be than any book that I had hitherto seen, for, although he very modestly complains of its incompleteness, it is undoubtedly much more thorough and accurate than are most works of its class.

The idea occurred to me, that by bringing my experience in the use of such books to bear upon the completion and amendment of Mr. Courtney's work, I might render a good service to the thousands who have almost daily occasion to consult a book of this character;—and in some degree make up for the loss that the community sustained in his death, although I cannot hope to bring to the task either the patience or the experience that constituted his great merit as a compiler.

It would be presumption to claim that, even in its enlarged and corrected condition, this book is complete, and all that could be desired, for there are more subjects of quite general interest to farmers and mechanics than could be properly *catalogued* in a book of this size. All that is claimed is, that so far as it goes it is correct; and that it goes as far, and in as many directions, as is compatible with its size and purpose.

The importance of having such a book as this always at one's elbow is very much greater than would at first sight be supposed by one who has not known the convenience of it.

How often, in farming, do we wish that we could know, on the spot, how to estimate the weight of hay in various conditions in the mow; the weight of cattle by measurement; the capacity of a grain bin; the weight of a piece of timber, or of a load of manure; the distance apart to which to set trees or plants in order to get a certain number within a certain space; the size of an irregular field. How often in mechanics do we need to know the strength and measurement of masonry; the contents of cisterns and small vessels; the area of circles; the quality of cements; the *power* value of fuel; the weight of bar iron, or of lead pipe; the fusing heat of metals; the strength of materials; or the board measure of scantling.

And, worst of all, how sadly we accustom ourselves to get along without knowing these things! How much we lose by *guessing* instead of *knowing*!

The object of this book is to put it within the power of every practical man to *know* these details;—to leave less to guessing, and to enable him to guide his daily operations by the light of positive knowledge. If it accomplishes this purpose, neither Mr. Courtney nor I will have worked in vain.

In addition to the many tables and statements of valuable facts with which the book abounds, I have thought it advisable to review very carefully all of its “agricultural” matter, and to add what I could, in the space allowed to me, that might be of interest to those farmers who care to look a little beyond the mere question of dollars and cents

in farming, and of value to those who believe (as, happily, a yearly increasing number do believe) that the road to surer and greater profit lies through the door that Science and Common Sense—the guardian angels of Agriculture—hold open to them.

It has not been possible to do much in this direction, for the subject is a very extended one, but I think that many a young farmer, if he will consider well the principles that are laid down under the headings of “Plants,” “Soils,” and “Manures,” will at least feel a desire to learn more of the simple truths which lie at the foundation of his practice.

I am sure, also, that it is not too much to say, that a careful study of the directions and the reasons for Tile-Draining will richly repay any occupier of cold, wet land for the purchase of the book.

This is a subject which, in this country at least, is still in the very early infancy of its progress. Not one acre in ten thousand of the land that it would pay well to drain in the best manner, has yet felt the benefit of the operation; and not one farmer in a thousand has the faintest conception of the fact,—a fact that ample experience, here and in Europe, has fully demonstrated,—that he can no more *afford* to farm an undrained heavy soil, than a carpenter can afford to work with a dull tool.

I have introduced another novelty into the work, under the head of “The Dry Earth System.” This is a bantling that has raised its head within a very few years, and is only now coming to be recognized at its full value; but it is

ushered before our attention with all the force that consideration of decency, health, and economy can lend; and the most thoughtful attention is asked for its claims. It is really the coming Reform, and promises more for civilization, and for national prosperity, than any improvement that has yet been brought to the notice of the public.

To sum up, then: this book is offered as containing more that has been proven by long use to be of value; more that it is most necessary for every farmer and mechanic to know; and more of promising novelty, than any other that has ever been presented to the farmers and mechanics of America.

It is complete in every particular in which it is possible for such a book to be complete, and, in addition to this, it is sufficiently suggestive in many other respects to induce its readers to read more, to think more, to experiment more, and to become more intelligent and more successful in the management of their business, as well as really happier and wiser men.

If it should be thought that I claim too much for a single Hand-Book, which is mainly filled with dry details concerning the measurement of boards, and the spacing of trees in an orchard, I trust that I shall at least not be condemned as an enthusiast until the reader has taken the trouble to examine carefully what I have to say, and to consider well to what better things the helping hand of Nature may lead him if he has the wisdom to heed its beckonings.

GEO. E. WARING, JR.

OGDEN FARM, NEWPORT, R. I., September, 1868.

TESTIMONIALS FOR THE FARMERS' AND MECHANICS' MANUAL.

From the New York Tribune.

A new edition of "THE FARMERS' AND MECHANICS' MANUAL," by W. S. Courtney, deceased, revised and enlarged by George E. Waring, Jr., introduces several improvements on the original work, forming a valuable book of general reference on practical affairs. It comprises a variety of tables and rules, and a thousand other points which perpetually occur in the experience of industrial life, and which are often decided by guess rather than by knowledge. The agricultural portions of the volume have been thoroughly revised by Mr. Waring, who has also enriched it with a variety of original matter, especially in relation to his favorite topics of "Tile-Draining," "The Dry Earth System," and others. In its present form, the work challenges the attention of every tiller of the soil and every lover of improvement. It is a sound, honest, instructive publication, doing all which it professes to do, and more, full of information suited not only to put money into the purse of the farmers and mechanics who consult its pages, but to increase their stock of valuable intelligence, and add to their resources for a happy and useful life.

From the New York Evening Post.

FARMERS' AND MECHANICS' MANUAL.—The work, as its title implies, is designed not less for the wants of the mechanic than the husbandman, but for both it would not be easy to exaggerate its usefulness. The entire matter of measurement, in its connection with weight, bulk, liquid contents, distance and superficial area, is exhausted in the simple tables and diagrams which constitute a large part of the text, and by means of these the mechanic or farmer may in a moment resolve a problem which might otherwise occupy a day.

Besides the tabular information which the book contains, there are hints upon the subjects of drainage, manures, stock-raising, rotation of crops, gardening for market, and steam cultivation, which the agriculturist will thankfully receive. In one respect the book is entitled to very high commendation—the accuracy of its typography. Crowded as it is with figures, but seven errata have been discovered in its five hundred pages. Messrs. Courtney and Waring have performed a most laborious and meritorious public service in its preparation. The moderate price at which it is offered to the public, three dollars a copy, must insure it an extended circulation.

From the Rural New Yorker.

This is a valuable and will be found a useful book to almost all classes of business men. Facts and figures of practical utility relating to all sorts of industry, the results of patient, elaborate calculation, are here crystallized into a condensed form ready for use, and so far as we have had opportunity to examine, the rules and tables are correct. We recommend it cordially to our readers.

From the New York Day-Book.

It would not answer for us to give a table of the contents of this very excellent and practical book, as an alphabetical arrangement of the same shows no less than eight hundred different items of information of value to the mechanic, manufacturer, merchant, builder, engineer, mason, painter, plumber, farmer, gardener, accountant, etc. The book is fairly crammed with solid and useful knowledge suited to these trades and professions; and it would seem as if the compilers themselves must have had years of service in each of these branches to arrive at so complete an understanding of what each branch ought to know. It is one of the most complete books of the day, and every member of the above professions should own a copy.

From the Hartford Courant, Conn.

A VALUABLE BOOK FOR ALL.—One of the most useful and valuable books for farmers, mechanics and working men which we have seen, is "THE FARMERS' AND MECHANICS' MANUAL." It is the useful and comprehensive "MANUAL" of W. S. Courtney, now deceased, and revised and enlarged by George E. Waring, Jr. It contains a great amount of information and statistics, arranged conveniently for reference, concerning matters that every farmer, mechanic and business man must inform himself about almost every day. It is an invaluable hand-book and book of reference.

From San Francisco Bulletin, Cal.

We have been favored with a look at a work entitled "THE FARMERS' AND MECHANICS' MANUAL," which we are sure is destined to achieve a marked success; and do not hesitate to add our unqualified commendation of the aim and execution of the work.

From W. S. CLARK, Esq., President of the Massachusetts Agricultural College.

Please accept thanks for a copy of a very useful book, styled "THE FARMERS' AND MECHANICS' MANUAL." It is full of valuable information, and Mr. Waring's name is a sufficient guarantee of its correctness. I shall advise my students to act as agents for its sale.

LIST OF ILLUSTRATIONS.

ENGRAVINGS.	PAGE
1. HARVEST TIME.....	Frontispiece
2. ILLUSTRATING SEASONS, LONGITUDE, &c.....	19-22
2. " CIRCULAR MEASURE.....	23, 24
2. " MEASURE OF TIME.....	25-30
1. " PENDULUMS.....	31
1. " WEATHER.....	34
2. " WINDMILLS.....	35, 36
14. " MEASUREMENT OF LAND.....	43-46
1. " GOVERNMENT LAND MEASURE.....	50
3. " MEASUREMENT OF HAY.....	51-56
2. " " " CORN IN THE CRIB.....	57-59
1. " " " GRAIN IN GRANARIES....	60
1. " " " TIMBER.....	61
1. " " " WOOD.....	62
1. " " " ROUND TIMBER.....	64
2. " GAUGING OF CASKS.....	79, 80
3. " CAPACITY OF WAGON BEDS.....	82, 83
2. " FALSE BALANCES.....	84, 85
3. " CISTERNS.....	86-92
1. " HYDRAULICS.....	97
3. " HYDRAULIC RAM.....	102-109
1. " " PRESS.....	110
2. " FUEL.....	115-124
1. " FENCES.....	125
1. " HEDGES.....	133
1. " HORSE POWER.....	137
1. " PLOUGHING.....	141
2. " FREIGHTING VESSELS.....	142-144
25. " UNITED STATES MONEY.....	145-148
16. " ENGLISH MONEY.....	149-151

LIST OF ILLUSTRATIONS.

ENGRAVINGS.		PAGE
12.	ILLUSTRATING AVOIRDUPOIS WEIGHT.....	152-154
5.	" TROY "	156, 157
6.	" APOTHECARIES' "	158, 159
5.	" LIQUID MEASURE	160, 161
6.	" DRY "	162-164
1.	" SQUARE "	165
1.	" LONG "	167
4.	" CUBIC "	170, 171
2.	" METRIC SYSTEM OF WEIGHTS AND MEASURES	177
1.	" SPECIFIC GRAVITY.....	183
1.	" CORN AND PORK.....	196
1.	" LIFE AND INCREASE OF ANIMALS.....	197
4.	" THE AGE OF ANIMALS.....	201-205
2.	" " COMPUTED WEIGHT OF CATTLE.....	209-211
2.	" " FOOD OF ANIMALS.....	212-215
1.	" LIGHTNING RODS.....	250
1.	" WEIGHT OF SQUARE AND ROLLED IRON....	273
1.	" MASONRY.....	276
1.	" MECHANICAL POWERS—INCLINED PLANE....	282
1.	" " " WEDGE.....	286
1.	" " " SCREW.....	288
1.	" " " PULLEY.....	290
24.	" MATHEMATICAL DEFINITIONS.....	292-295
1.	" MANURES.....	327
21.	" TILE DRAINING.....	363-372
1.	" BUTTER AND CHEESE.....	400
3.	" STEAMING FOOD FOR STOCK.....	414-423
1.	" GARDENING FOR MARKET.....	428
1.	" STEAM PLOUGHING.....	448

COMMERCIAL ABBREVIATIONS.

@.,	At.	Fr't.,	Freight.
q/c.,	Account.	Inst.,	This month.
¢.,	Cents.	Int.,	Interest.
#,	Number.	Mdse.,	Merchandise.
Am't.,	Amount.	Mo.,	Month.
Ass'd.,	Assorted.	Net,	Without discount.
Bal.,	Balance.	No.,	Number.
Bbl.,	Barrel.	Pay't.,	Payment.
Blk.,	Black.	Pk'gs.,	Packages.
Cons't.,	Consignment.	Per or pr.,	By.
Dft.,	Draft.	Prem.,	Premium.
Disc't.,	Discount.	Prox.,	Next month.
E. E.,	Errors excepted.	Ps.,	Pieces.
Exps.,	Expenses.	Sunds.,	Sundries.
Fol.,	Folio.	Ult.,	Last month.
Fwd.,	Forwarded.		

EXPLANATION OF ARITHMETICAL CHARACTERS USED IN THIS BOOK.

$=$ *Equal* ; as 12 inches $=$ 1 foot, or $4 \times 5 = 20$.

$+$ *Plus or more* ; signifies addition, as $3 + 5 + 7 = 15$.

$-$ *Minus or less* ; signifies subtraction, as $12 - 4 = 8$.

\times *Multiplied by* ; signifies multiplication, as $8 \times 7 = 56$.

\div *Divided by* ; signifies division, as $56 \div 8 = 7$.

$:$ $::$ $:$ *Proportion* ; as $2 : 4 :: 8 : 16$; that is, as 2 is to 4 so is 8 to 16.

$\sqrt{}$ Prefixed to a number denotes that the *square root* of that number is required, as, $\sqrt{36} = 6$.

$\sqrt[3]{}$ Prefixed to a number denotes that the *cube root* of that number is required, as, $\sqrt[3]{27} = 3$.

2 Added to a number signifies that the number is to be *squared*, as 4^2 means that 4 is to be multiplied by 4.

3 Added to a number signifies that the number is to be *cubed*, as 4^3 means $4 \times 4 \times 4 = 64$.

$.$ *Decimal point*, when prefixed to a number signified that that number has an unit (1) for its denominator, as .1 is $\frac{1}{10}$, .2 is $\frac{2}{10}$, .12 is $\frac{12}{100}$, .125 is $\frac{125}{1000}$, &c.

$^{\circ}$ Signifies degrees : ' minutes, and " seconds.

SEASONS, LONGITUDE, &c.



To reduce longitude to time.

The English count their degrees of longitude east and west from Greenwich, which, owing to our early dependence upon the mother country for books and science, became extensively adopted in this country, and still prevails to a considerable extent, especially in our nautical charts, and

works on navigation. But by an act of Congress, passed some thirty years ago, the meridian of Washington was established as the point of departure, and accordingly our maps, charts, &c., have since been adapted to that meridian.

The sun passes over a degree of longitude in 4 minutes—the 360° in 24 hours. Thus, when we travel west, or on a line with the sun, our watch is four minutes *fast* for every 60 geographical miles we travel. If we travel east, or on a line with the sun, it is four minutes *slow* for every degree we travel. Hence, when it is noon at Greenwich, that is, when the sun is on the meridian there, if we multiply 74° , the longitude of New York west from Greenwich, by 4, and subtract the result from 12 o'clock M., it will give the corresponding time at New York. Thus, $74^\circ \times 4 = 296$ minutes, which, divided by 60, gives 4 hours and 56 minutes for the sun to travel from Greenwich to New York. Subtracting this from 12 o'clock (the Greenwich time) gives 7 o'clock and 4 minutes A.M. as the corresponding time at New York. So also by reverse, when it is noon at New York, it is 4 hours and 56 minutes past noon at Greenwich. Hence results the following

RULE.—Multiply the number of degrees, minutes, and seconds west or east of the given meridian by 4, reduce the product to hours, &c., and for west longitude subtract

from 12 hours, and for east longitude add to 12 hours (*i. e.*, so many hours past 12), and the result will be the corresponding time.

EXAMPLE.—Required the time at longitude $50^{\circ} 31'$ west, corresponding to noon at Greenwich?

SOLUTION.— $50^{\circ} 31' \times 4 = 3$ hours 22 min. 4 sec.—12 = 8 h. 37 min. 56 sec. A.M. *Ans.*


NOTE.—Time is both *apparent* and *mean*. The sun is on the meridian at 12 o'clock on four days only in the year. It is sometimes as much as $16\frac{1}{4}$ minutes before or after 12 when its shadow strikes the noon mark on the sundial. This is occasioned by the irregular motion of the earth on its axis and the inclination of its poles. This is called *apparent time*. *Mean time* is determined by the *equation* of these irregularities for every day in the year, and is noted in all good almanacs. The latter is the true or correct time. The foregoing rule is applicable to either.

When you buy an almanac, buy one that expresses on each calendar page the *mean time* when the sun reaches the meridian, or the shadow the noon-mark on the dial, and set your time-piece fast or slow as indicated in the almanac.

To ascertain the length of the day and night.

At any time in the year, add 12 hours to the time of the sun's setting and from the sum subtract the time of rising

for the *length of the day*. Subtract the time of setting from 12 hours, and to the remainder add the time of rising the next morning for the *length of the night*. This rule is true of either apparent or mean time.



CIRCULAR OR ANGULAR MEASURE.


This Measure is used to measure angles or the arcs of circles. It is used in astronomy, geography, navigation, and surveying, and for calculating differences of time.

TABLE.

60 seconds (") make	1 minute,	marked '
60 minutes "	1 degree,	" °
30 degrees "	1 sign,	" sig.
90 degrees "	{ 1 quadrant,	" quad.
	{ 1 right angle,	" r. a.
4 quadrants or	{ 1 { circumference	
12 signs "		" cir.

NOTES.—1. The greatest distance across a circle is called its *diameter*. The distance around it is called its *circumference*. Any part of the circumference is called an *arc*.

2. If any circumference, whether large or small, be divided into 360 equal arcs, each arc is called a degree. The



degree is divided into 60 minutes, and the minute into 60 seconds. The length of a degree, minute, or second, depends on the size of the circle. If the size of the circle is increased or decreased, the length of the degree, minute, or second is also increased or decreased.

3. The greatest circumference of the earth's surface is about 24,930 miles ; 1° of that circumference is one 360th of 24,930 miles, which is $69\frac{1}{4}$ miles.

4. A geographical or nautical mile is equal to $1'$ of the earth's greatest circumference, which is found to be a little more than one statute mile and 49 rods.

5. Latitude is measured north or south from the equator on any meridian, and is expressed in degrees, minutes, and seconds ; thus, $43^\circ 17' 31''$ north lat. denotes a position $43^\circ 17' 31''$ north from the equator.

6. The linear extent of a degree of longitude depends upon the latitude, and diminishes as the latitude increases ; thus, at latitude 10° its extent is 359640 feet ; at lat. 40° it is 280106 feet ; and at lat. 80° it is only 63612 feet.



MEASURE OF TIME.

Time is the measure of duration.




TABLE.

60 seconds (sec.)	make 1 minute,	marked min.
60 minutes	“ 1 hour,	“ h.
24 hours	“ 1 day,	“ da.
7 days	“ 1 week,	“ wk.
4 weeks 2 days, or }	“ 1 month,	“ mo.
30 days		

365 days, or	}	make 1 year,	marked yr.
52 weeks 1 day			
12 calendar months			
100 years		" 1 century,	" C.

The calendar year is divided as follows :

Season.	Months.	No. of days.	Abbreviations.
Winter	1. January	31	Jan.
	2. February	28 or 29	Feb.
Spring	3. March	31	Mar.
	4. April	30	Apr.
	5. May	31	—
Summer	6. June	30	Jun.
	7. July	31	—
	8. August	31	Aug.
Autumn	9. September	30	Sept.
	10. October	31	Oct.
	11. November	30	Nov.
Winter	12. December	31	Dec.

365 or 366.

NOTES.—1. The exact length of the solar year is 365 days 5 h. 48 min. 49 sec. ; but, for convenience, it is reckoned 11 min. 11 sec. more than this, or 365 da. 6 h. = 365½ days. This ½ day in four years makes 1 day, which every fourth year (called Bissextile or leap year) is added to the shortest month, giving it 29 days. The numbers de-

noting leap years are exactly divisible by 4 ; as, 1856, 1860, 1864 ; except years whose number can be divided without a remainder by 100, but not by 400.

2. Owing to an error in the Julian calendar, it was decreed by the British Government that the day following the second day of September, 1752, should be called the fourteenth day of September, or that 11 days should be stricken from the calendar.

3. Time, previous to this decree, is called *Old Style* (O. S.), and since, *New Style* (N. S.). Russia still reckons time by the Old Style, hence their dates are 12 days behind ours.

4. In most business transactions 30 days are called a month, and 52 weeks a year.

5. The centuries are numbered from the commencement of the Christian era ; the months from the commencement of the year ; the days from the commencement of the month ; and the hours from the commencement of the day (12 o'clock, midnight), and from mid-day or noon. A.M. denotes time before noon, M., at noon, and P.M., after noon. Thus, 9 o'clock A.M., May 23, 1860, is the end of the ninth hour of the 23d day of the fifth month of the 60th year of the 19th century.

6. A decade is a period of 10 years.

7. The *Lunar Cycle*, or *Golden Number*, is a period of 19 years, after which the changes of the moon return on the same days of the month.

8. The *Solar Cycle* is a period of 28 years, when the days of the week again return to the same days of the month.

To find the golden number or lunar cycle.

RULE.—Add 1 to the given year; divide the sum by 19, and the remainder is the golden number.

EXAMPLE.—What is the golden number for 1857?

SOLUTION.— $1857 + 1 \div 19 = 97$, rem. 15. *Ans.*

NOTE.—If 0 remain, it will be 19. Hence, in 1861, the changes of the moon occur on the same days of the month they did in 1842, 1823, 1804, &c.

TABLE showing the number of days from any day in one month to the same day in any other.

FROM	Jan.	Feb.	Mar.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
January.....	365	31	59	90	120	151	181	212	243	273	304	334
February.....	334	365	28	59	89	120	150	181	212	242	273	303
March.....	306	337	365	31	61	92	122	153	184	214	245	275
April.....	275	306	334	365	30	61	91	122	153	183	214	244
May.....	245	276	304	335	365	31	61	92	123	153	184	214
June.....	214	245	273	304	334	365	30	61	92	122	153	183
July.....	184	215	243	274	304	335	365	31	62	92	123	153
August.....	153	184	212	243	273	304	334	365	31	61	92	122
September...	122	153	181	212	242	273	304	334	365	30	61	91
October.....	92	123	151	282	211	243	273	304	335	365	31	61
November...	61	92	120	151	181	212	242	273	304	334	365	30
December...	31	62	90	121	151	182	212	243	274	304	335	365

EXPLANATION.—Find, in the left-hand column, the month from any day of which you wish to compute the number of days to the same day in any other month, and follow the line along until under the latter, and you have the

required number of days. Thus, from the 12th of April to the 12th of October, is 183 days; from the 7th of March to the 7th of June, 92 days.

TABLE for finding the number of days between two dates—
new method.


Jan.	Feb.	Mar.	April	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.
1	32	60	91	121	152	182	213	244	274	305	335
2	33	61	92	122	153	183	214	245	275	306	336
3	34	62	93	123	154	184	215	246	276	307	337
4	35	63	94	124	155	185	216	247	277	308	338
5	36	64	95	125	156	186	217	248	278	309	339
6	37	65	96	126	157	187	218	249	279	310	340
7	38	66	97	127	158	188	219	250	280	311	341
8	39	67	98	128	159	189	220	251	281	312	342
9	40	68	99	129	160	190	221	252	282	313	343
10	41	69	100	130	161	191	222	253	283	314	344
11	42	70	101	131	162	192	223	254	284	315	345
12	43	71	102	132	163	193	224	255	285	316	346
13	44	72	103	133	164	194	225	256	286	317	347
14	45	73	104	134	165	195	226	257	287	318	348
15	46	74	105	135	166	196	227	258	288	319	349
16	47	75	106	136	167	197	228	259	289	320	350
17	48	76	107	137	168	198	229	260	290	321	351
18	49	77	108	138	169	199	230	261	291	322	352
19	50	78	109	139	170	200	231	262	292	323	353
20	51	79	110	140	171	201	232	263	293	324	354
21	52	80	111	141	172	202	233	264	294	325	355
22	53	81	112	142	173	203	234	265	295	326	356
23	54	82	113	143	174	204	235	266	296	327	357
24	55	83	114	144	175	205	236	267	297	328	358
25	56	84	115	145	176	206	237	268	298	329	359
26	57	85	116	146	177	207	238	269	299	330	360
27	58	86	117	147	178	208	239	270	300	331	361
28	59	87	118	148	179	209	240	271	301	332	362
29		88	119	149	180	210	241	272	302	333	363
30		89	120	150	181	211	242	273	303	334	364
31		90		151		212	243		304		365

NOTE.—To find from the above table the number of days between two dates, we give the following—


RULE.—I. When the dates are in the same year, subtract the number of days of the earlier date from the number of days of the later date; the result will be the number of days required.

II. When the dates are in consecutive years, subtract the number of days of the earlier date from 365, and add to the remainder the number of days of the later date; the result will be the number of days required.

When the year is a leap year, add one day to the result.



PENDULUMS.



The vibrations of pendulums are as the square roots of their lengths. The length of one that will vibrate seconds in New York, at the level of the sea, is 39.1013 inches.

To find the length of a pendulum for any given number of vibrations per minute.

RULE.—As the number of vibrations given is to the square root of 39.1013 inches, so is 60 to the square root of the length of the pendulum required.

EXAMPLE.—What is the length of a pendulum that will make 50 vibrations per minute?

SOLUTION.—50 : 6.25 (the sq. root of 39.1013) :: 60 : 7.5, then $7.5^2 = 56.25$ inches. *Ans.*

To find the number of vibrations per minute, the length of the pendulum being given.

RULE.—As the square root of the length of the pendulum is to 60, so is the square root of 39.1013 to the number of vibrations required.

EXAMPLE.—How many vibrations will a pendulum 64 inches long make in a minute?

SOLUTION.—8 (square root of 64) : 60 :: 6.25 (sq. root of 39.1013) : 46.875 vibrations. *Ans.*

TABLE showing the planets, comparative size, &c., in the solar system.

NAMES.	Mean Diameter.	Mean distance from the Sun.	Revolution ar'd the Sun.	Revolution on axis.	Velocity pr. minute in orbit.	Size—the Earth being 1.	Density: Earth being 1.	Light: Earth being one.
	Miles.	Miles.	yrs. days	d. h. m.	Miles.			
THE SUN.....	888,246	25 9 59	1,412,921.100	0.252	infin.
Mercury.....	3,224	36,814,000	...	88 1 0 5	1,827	0.053	1.120	6.680
Venus.....	7,687	68,787,000	...	224 .. 23 21	1,338	0.909	0.923	1.911
The Earth.....	7,912	95,103,000	1 ..	23 56	1,138	1.000	1.000	1.000
The Moon.....	2,180	95,103,000	1 ..	27 7 43	38	0.020	0.615	1.000
Mars.....	4,189	144,908,000	1 321	1 0 37	921	0.125	0.948	0.431
Jupiter.....	89,170	494,797,000	11 215	.. 9 56	496	1,456.000	0.238	0.037
Saturn.....	79,042	907,162,000	29 167	.. 10 29	368	771.000	0.138	0.011
Uranus.....	35,112	1,824,290,000	84 6	1 13 33	259	80.000	0.242	0.003
Neptune.....	41,500	2,854,000,000	164 226	208	143.000	0.140	0.001

THE WEATHER.

The following table, and the accompanying remarks, originally formed by Dr. Herschel, and approved with some alterations by the experienced Dr. Adam Clarke, are the result of many years' close observation; the whole being on a due consideration of the attraction of the sun and moon, in their several positions respecting the earth, and will, by inspection, show the observer what kind of weather will *most probably* follow the entrance of the moon into any of its quarters—so probably, indeed, that it has seldom been found to fail.

TABLE, for telling the weather through all the lunations of each year forever.

MOON	TIME OF CHANGE.	IN SUMMER.	IN WINTER.
If the new moon, the 1st quarter, full moon, or last quar., happens	Between midnight and two in the morning,	Fair.	Hard frost, unless the wind be S. or W.
	— 2 and 4, morning,	Cold, with frequent showers.	Snow and stormy.
	— 4 and 6 “	Rain.	Rain.
	— 6 and 8 “	Wind and rain.	Stormy.
	— 8 and 10 “	Changeable.	Cold rain, if wind be W.; snow if E.
	— 10 and 12 “	Frequent showers.	Cold and high wind.
	At 12 o'clock at noon, and 2, P. M.	Very rainy.	Snow or rain.
	Between 2 and 4, P. M.	Changeable.	Fair and mild.
	— 4 and 6, “	Fair.	Fair.
	— 6 and 8, “	Fair, if wind N. W. } Rainy, if S. or S. W. }	Fair and frosty, if wind N. or N. E.
	— 8 and 10, “	Do.	Rain or snow, if S. or S. W.
	— 10 and midnight,	Fair.	Do. Fair and frosty.

OBSERVATIONS.—1. The nearer the time of the moon's

change, first quarter, full, or last quarter are to *midnight*, the fairer will the weather be during the seven days following.


2. The space for this calculation occupies from ten at night till two next morning.

3. The nearer to *mid-day* or *noon*, the phases of the moon happen, the more foul or wet weather may be expected during the next seven days.

4. The space of this calculation occupies from ten in the forenoon to two in the afternoon. These observations refer principally to the summer, though they affect spring and autumn nearly in the same ratio.



WIND.



The force of the wind increases directly as the square of the velocity. Thus, a wind blowing 10 miles an hour exerts a pressure four times as great as at 5 miles an hour, and 25 times as great as at 2 miles an hour.

To find the force of wind acting directly against a surface.

RULE.—Multiply the surface in square feet by the lbs. pressure per square foot as given in the following table.


EXAMPLE.—What is the pressure of a wind of a velocity of 20 miles per hour against a barn door 10 feet by 6?

SOLUTION.— $10 \times 6 = 60$ sq. ft., surface, $\times 2$ lbs., pressure per square foot, $= 120$ lbs. *Ans.*

TABLE, *showing the force and velocity of wind.*

Miles per hour.	Feet per minute.	Lbs pressure on 1 sq. foot.	Description.
1	88	.005	Barely observable.
2	176	.020	Just perceptible.
3	264	.045	
4	352	.080	
5	440	.125	Gentle, pleasant wind.
6	528	.180	
8	704	.320	
10	880	.500	
15	1320	1.125	Brisk blow.
20	1760	2.000	Very brisk.
25	2200	3.125	
30	2640	4.500	High wind.
35	3080	6.125	
40	3520	8.000	Very high.
45	3960	10.125	
50	4400	12.500	Storm.
60	5280	18.000	Great storm.
80	7040	32.000	Hurricane. [ing off buildings, &c.
100	8800	50.000	Tornado, uprooting trees, sweep-

The mechanical force of wind is well illustrated in the old-fashioned windmills, which were used for the purpose of



Windmill.

raising water and grinding grain, where facilities for steam or water-power were wanting.

AVERAGE TEMPERATURE AND FALL OF RAIN.

TABLE, *showing the average temperature of the four Seasons at points on the Pacific and Atlantic coasts, and the interior of this continent.*

	TEMPERATURE.					
	Latitude.	Spring.	Summer.	Autumn.	Winter.	Year.
PACIFIC COAST.						
Monterey,.....	36°36'	53°·99	58°·64	57°·29	51°·22	55°·29
San Francisco.....	37°48'	54°·41	57°·33	56°·83	50°·86	54°·88
Astoria.....	46°11'	51°·16	61°·58	53°·76	42°·43	52°·23
INTERIOR.						
St. Louis Arsenal,.....	38°40'	54°·15	76°·19	55°·44	32°·27	54°·51
Chicago.....	41°52'	44°·90	67°·33	48°·85	25°·90	46°·75
Fort Ripley.....	46°19'	39°·33	64°·94	42°·91	10°·01	39°·30
ATLANTIC COAST.						
Fort Monroe, near Norfolk,....	37°	56°·87	76°·57	61°·68	40°·45	58°·89
Fort Columbus, N. Y. Harbor,	40°42'	48°·74	72°·10	54°·55	31°·38	51°·69
Fort Sullivan, Eastport,.....	44°15'	40°·15	60°·50	47°·52	23°·90	43°·02

From this table it will be perceived that Astoria, on the Pacific coast, and Fort Ripley in the interior, are in about the same latitude. Astoria, though 650 miles north of Monterey, is only 3 degrees colder. Fort Ripley is *fifteen* degrees colder than St. Louis, although it is only about 500 miles further north.

San Francisco, St. Louis, and Fort Monroe, are in about the same latitude. The difference between the mean summer and winter temperature of San Francisco is less than *seven* degrees; of St. Louis, nearly *forty-four* degrees; and of Fort Monroe, *thirty-six* degrees. Eastport is two degrees south of Astoria, but is *nine* degrees colder.

The United States may be divided with reference to the fall of rain into three regions, namely: the region of periodical rains, the region of frequent rains, and the region of scanty rains.

The region of periodical rains comprises the western division of the Pacific slope.

In that portion of this division south of the 40th parallel of latitude, scarcely any rain falls in summer, and very little in autumn. The quantity in winter somewhat exceeds that which falls during the spring.

A much greater quantity of rain falls upon that part of the division north of lat. 40° than south of it; but, as in the southern division, the largest amount belongs to the winter and spring.

The region of frequent rains extends from the Atlantic coast westward to about the 100th meridian of longitude. This region, considered as a whole, is exceedingly well watered, the rain being quite equally distributed through the different seasons.

From an examination of the table, it will appear that along the Atlantic slope, as far south as Washington, very nearly the same annual quantity of rain falls; and that it is very equally distributed throughout the year. In the Gulf States, and along the Atlantic slope south of Washington, the annual amount of rain is much greater than in the other sections, and the summer rains are much more abundant than those of the winter. In the interior the

annual quantity is less, and generally much less rain falls in winter than in the other seasons.

The region of scanty rains embraces the country between about the 100th meridian of longitude and the Cascade and Sierra Nevada Mountains. It includes the northern and southern divisions of the Pacific slope, the inland basin of Utah, the table-lands of the Texas slope, and the sterile region east of the Rocky Mountains.

Among the mountains of this region a considerable quantity of rain falls, and violent showers are experienced in all seasons of the year. Some of the mountain valleys are also well watered. Thus the annual fall of rain at Santa Fe, situated on a plateau enclosed by mountains, is 19.83 inches; and the fall at Fort Massachusetts, which is situated in a valley 100 miles further north, is 20.54 inches.

The annual fall of rain in the desert region, through which the great Colorado flows, is estimated at three inches; that of the inland basin of Utah, at five inches; of the Great Plain south of the Columbia River, ten inches; of the Llano Estacado, ten inches; and of the sterile region east of the Rocky Mountains, from fifteen to twenty inches. In all these sections scarcely any rain falls in summer.

The greatest amount of rain reported in the "Army Meteorological Register," for any given year, was the fall, in 1846, at Baton Rouge, of 116.6 inches; the least, a fall, in 1853, at Fort Yuma, California, of 1.78 inches.

[This valuable Table is compiled from the "Army Meteorological Register," and presents the results of all the records, in the Army Medical Bureau, for 33 years, from 1822 to the close of 1854.]

TABLE, showing the latitude and longitude, the elevation above the level of the sea, the mean annual temperature, and the average annual fall of rain at various places in the United States.

NAME OF PLACE OF OBSERVATION	Latitude.	Longitude West from Greenwich.	Elevation above the level of the sea in feet.	Mean Annual temperature.	Average annual fall of rain in inches.
Fort Kent, Maine.....	47° 15'	68° 35'	575	37° 04	36.46
Fort Fairfield, Maine.....	46 46	67 49	415	38.11	
Hancock Barracks, Maine.....	46 07	67 49	620	40.51	36.97
Fort Sullivan, Eastport, Maine....	44 54	66 58	70	43.02	39.39
Fort Preble, Portland, Maine.....	43 39	70 20	20	45.22	45.25
Fort Constitution, Portsmouth, N.H.	43 04	70 49	40	45.81	35.57
Fort Independence, Bost. Har., Mass	42 20	71	50	48.92	35.30
Watertown Arsenal, Mass.....	41 21	71 09		47.34	42.07
Fort Adams, Rhode Island.....	41 29	71 20	40	49.70	52.46
Fort Wolcott, Newport Harbor, R. I.	41 30	71 20	20	50.72	
Fort Trumbull, New London, Conn.	41 21	72 06	23	49.62	45.69
Fort Columbus, N. Y. Harbor.....	40 42	74 01	23	51.69	42.23
Fort Hamilton, N. Y. Harbor.....	40 37	74 02	25	51.54	43.65
West Point, New York.....	41 23	74	167	50.73	54.15
Watervliet Arsenal, New York....	42 43	73 43	50?	48.07	34.55
Plattsburg Barracks, New York....	44 41	73 25	186	44.	33.39
Sackett's Harbor, New York.....	43 57	76 15	262	46.38	39.78
Fort Ontario, New York.....	43 20	76 40	250	46.44	30.88
Fort Niagara, New York.....	43 18	79 08	250	47.91	31.77
Buffalo Barracks, New York.....	42 53	78 58	660	46.25	38.80
Alleghany Arsenal, Pittsburg, Pa..	40 32	80 02	704	50.86	34.96
Carlisle Barracks, Carlisle, Pa.....	40 12	77 14	500	51.10	34.01
Fort Mifflin, Pa.....	39 53	75 13	20	53.85	45.27
Fort Delaware, Del.....	39 35	75 34	10	56.06	
Fort McHenry, Md.....	39 17	71 35	36	54.36	42.
Fort Severn, Md.....	38 58	76 27	20	55.42	48.61
Washington City, D. C.....	38 53	77 02	50-90	56.14	41.20
Fort Washington, Md.....	38 43	77 06	60	57.87	45.02
Bellona Arsenal, Richmond, Va....	37 20	77 25	120	59.27	
Fort Monroe, Va.....	37	76 18	8	58.89	50.89
Fort Macon, N. C.....	34 41	76 40	20	62.23	
Fort Johnston, N. C.....	34	78 05	20	65.68	46.01
Augusta Arsenal, Ga.....	33 28	81 53	600?	64.01	23.
Fort Moultrie, Charleston, S. C....	32 45	79 51	25	66.58	44.92
Oglethorpe Barracks, Ga.....	32 05	81 07	40	67.44	53.33
Fort Marion, St. Augustine, Fla....	29 38	81 35	25	69.61	31.80
Fort Shannon, Palatka, East Fla....	29 34	81 48	25	69.64	48.68
New Smyrna, East Fla.....	28 54	81 02	20	69.17	


Table continued.

NAME OF PLACE OF OBSERVATION	Latitude.	Longitude West from Greenwich.	Elevation above the lev- el of the sea, in feet.	Mean Annual temperature.	Average An- nual fall of rain in inches.
Fort Pierce, East Fla.	27°30'	80°20'	30	73°·20	62·98
Fort Dallas, East Fla.	25 55	80 20	20	74·75	
Key West, Fla.	24 32	81 48	10	76·51	47·65
Fort Myers, South Fla.	26 38	82 02	50	75·04	62·26
Fort Brooke, Tampa Bay, Fla.	28	82 28	20	71·92	55·47
Fort Meade, Fla.	28 01	82	80	71·48	40·22
Fort Micanopy, Fla.	29 30	82 28	60?	70·09	
Fort King, Fla.	29 10	82 10	50	70·	
Cedar Keys, Fla.	29 07	83 03	35	69·60	48 50
Fort Fanning, Fla.	29 35	83	50	70·20	
Fort Barrancas, Pensacola, Fla.	30 18	87 27	20	68·74	56·98
Fort Morgan, Ala.	30 14	88	20	66·88	
Mt. Vernon Arsenal, Ala.	31 12	88 02	200?	65·84	63·56
Fort Pike, La.	30 10	89 38	10	69·86	71·92
Fort Wood, La.	30 08	89 51	20	69·25	60·63
New Orleans, La.	29 57	90	10	69·86	50 96
Baton Rouge, La.	30 26	91 78	41	68·14	62·10
Fort Jessup, La.	31 33	93 32	80?	66·34	45·85
Fort Towson, Ind. Ter.	34	95 33	300?	61·69	51·08
Fort Washita, Indian Ter.	24 14	96 38	645	62·21	41·66
Fort Smith, Arkansas.	35 23	94 29	460	60·02	42 10
Fort Gibson, Ind. Ter.	34 47	95 10	560	60·81	36·46
Fort Scott, Mo.	37 45	94 35	1000?	54·50	42·12
Jefferson Barracks, Mo.	38 28	90 15	472	55·46	37·83
St. Louis Arsenal, Mo.	38 40	90 05	450	54·51	41·95
Newport Barracks, Newport, Ky. ...	39 05	84 29	500	55·26	
Detroit, Mich.	42 20	82 58	580	47·25	30·07
Fort Gratiot, Mich.	42 55	82 23	598	46·29	32·62
Fort Mackinac, Mich.	45 51	84 32	728	40·65	23·87
Fort Dearborn, Chicago, Ill.	41 52	87 35	591	46·75	
Fort Brady, Mich.	46 30	84 43	600	40·37	31·35
Fort Wilkins, Mich.	47 30	88	620	41·	
Fort Howard, Wis.	44 30	88 05	620	44·49	34·65
Fort Winnebago, Wis.	43 31	89 28	770?	44·80	27·49
Fort Crawford, Wis.	43 05	91	642	47·63	31·40
Fort Armstrong, Ill.	41 30	90 40	528	50·31	
Fort Atkinson, Iowa.	43	92	700?	45·50	39·74
Fort Des Moines, Iowa.	41 32	93 38	780	49·74	26·56
Fort Ripley, Minnesota.	46 19	94 19	1130	39·30	29·48
Fort Snelling, Minn.	44 53	93 10	820	44·54	25·43
Fort Leavenworth, Kansas.	39 21	94 44	896	52·78	30·29
Council Bluffs, Nebraska.	41 30	95 48	1250	49·28	
Fort Kearney, Nebraska.	40 38	98 57	2360	47·67	27·98
Fort Laramie, Nebraska.	42 12	104 47	4519	50·06	19·98
Fort Arbuckle Ind Ter.	34 27	97 09	1000?	60·83	30·57
Fort Belknap, Texas.	33 08	98 48	1600?	63·99	22·

Table continued.

NAM. OF PLACE OF OBSERVATION.	Latitude.	Longitude West from Greenwich.	Elevation above the lev- el of the sea, in feet.	Mean Annual temperature.	Average An- nual fall of rain in inches.
Fort Worth, Texas.....	32°40'	97°25'	1100?	63° 54	40·86
Phantom Hill, Texas.....	32 30	99 45	2300?	63·73	17·22
Fort Chadbourne, Texas.....	31 38	100 40	2120	62·38	31·88
Fort Graham, Texas.....	31 56	97 26	900?	65·76	40·58
Fort Gates, Texas.....	31 26	97 49	1000?	66·12	
Fort Croghan, Texas.....	30 40	98 31	1000?	65·74	36·56
San Antonio, Texas.....	29 25	98 25	600	69·25	33·77
Fort Merrill, Texas.....	28 17	98	150?	71·37	
Fort Ewell, Texas.....	28 05	98 57	200	71·30	
Corpus Christi, Texas.....	27 47	97 27	20	70·95	30·82
Fort Brown, Texas.....	25 54	97 26	50	73·75	33·65
Ringgold Barracks, Texas.....	26 23	99 02	200?	74·21	20·95
Fort McIntosh, Texas.....	27 31	99 21	400	73·24	18·66
Fort Duncan, Eagle Pass, Texas...	28 42	100 30	800	70·85	22·20
Fort Inge, Texas.....	29 09	99 07	845	67·69	27·99
Fort Lincoln, Texas.....	29 22	99 33	900?	68·03	20·58
Fort Clark, Texas.....	29 17	100 25	1000?	61·04	21·80
Fort Fillmore, New Mexico.....	32 13	106 42	3937	63·98	9·28
Fort Webster, New Mexico.....	32 48	108 04	6350	54·84	8·79
Fort Conrad, New Mexico.....	33 34	107 09	4576	59·40	6·76
Albuquerque, New Mexico.....	35 06	106 38	5032	56·32	9·42
Cebolleta and Laguna, New Mexico	35 03	107 14	6000	55·12	12·05
Santa Fé, New Mexico.....	35 41	106 02	6846	50·59	19·83
Las Vegas, New Mexico.....	35 35	105 16	6418	49·14	19·24
Fort Union, New Mexico.....	35 54	104 57	6670	49·14	19·24
Fort Massachusetts, New Mexico...	37 32	105 23	8365	49·11	20·54
Fort Defiance, New Mexicc.....	35 44	109 15	7200?	46·92	16·64
Fort Yuma, California.....	32 43	114 36	120	73·62	3·24
San Diego, California.....	32 42	117 14	150	62·	10·43
Posts Del Chino and Jurupa, Cal'a.	34	117 25	1000?	63·28	13·77
Monterey, California.....	36 36	121 52	140	55·29	12·20
Fort Miller, California.....	37	119 40	402	66·	24·51
San Francisco, California.....	37 48	122 26	150	54·88	23·59
Benicia Barracks, California.....	38 03	122 08	64	58·29	16·62
Sacramento, California.....	38 33	121 20	50	59·89	21·32
Fort Reading, California.....	40 30	122 05	674	62·09	29·02
Fort Humboldt, California.....	40 46	124 09	50	52·80	16·77
Fort Jones, California.....	41 36	122 52	2570	51·40	16·77
Fort Orford, California.....	42 44	124 29	50	53·62	68·52
Fort Vancouver, Oregon.....	45 40	122 30	50	52·65	45·50
Fort Dalles, Oregon.....	45 36	120 55	350	52·79	14·32
Fort Steilacoom, Washington Ter..	47 10	123 25	300?	50·82	51·75
Astoria, Oregon.....	40 11	123 48	50	52 23	
Great Salt Lake, Utah.....	40 46	112 06	4351	53·24	

MEASUREMENT OF LAND.



Every farmer should know the contents, in acres, of each of his fields, meadows, and lots, to ascertain which he should have a rod measure, a light stiff pole, just $16\frac{1}{2}$ feet long, with division marks on it of a yard each, making $5\frac{1}{2}$ yards. Provided with this measure, and proceeding according to the following rules, he can ascertain the area in acres of each of his fields, lots, &c.


Where the field is a square, a parallelogram, a rhombus, or a rhomboid.



Square.



Parallelogram.



Rhombus.





Rhomboid.

RULE.—Multiply the length in rods by the breadth in rods, and divide the product by 160, and the quotient will be the number of acres.

EXAMPLE.—What is the area in acres of a field of 30 rods long by 28 rods wide.

SOLUTION.— $30 \times 28 = 840 \div 160 = 5$ acres and 40 rods, or $5\frac{1}{4}$ acres. *Ans.*

Where the field is triangular.




RULE.—Multiply the base or longest side, in rods, by the perpendicular height (*i.e.*, the greatest width), in rods, and divide half the product by 160, and the quotient will be the number of acres.


EXAMPLE.—What is the area in acres of a triangular field, the base of which is 60 rods long, and its perpendicular height 28 rods?

SOLUTION.— $60 \times 28 = 1680 \div 2 = 840 \div 160 = 5$ acres and 40 rods, or $5\frac{1}{4}$ acres. *Ans.*

When the field is a trapezium or a trapezoid.





Trapezium.



Trapezoid.

RULE.—Divide it diagonally by a line running from one extreme corner to the other, which will cut the field into two triangles; then proceed with each as in the foregoing rule, and add the areas of the two triangles together. The product will be the number of acres.


Where the field is an irregular polygon.



RULE.—Draw diagonals to divide the field into triangles; find the area of each separately, and the sum of the whole will be the number of acres.

NOTE.—There are very few fields or lots which cannot be measured by cutting them into triangles, and proceeding by the above rule. In fact, all straight-sided fields can be so measured.

Where the field is long, and the sides crooked and irregular.




RULE.—Take the breadth in rods in a number of places, at equal distances apart; add them, and divide the sum by the number of breadths for the mean average or breadth; then multiply that by the length in rods and divide the product by 160, and the quotient will be the number of acres.

EXAMPLE.—What is the area in acres of a long irregular-sided field, the length of which is 80 rods, and its breadths at 10 rods apart are as follows, viz.: 8, 10, 11, 9, 8, 7, 9, 10 rods?

SOLUTION. — $8 + 10 + 11 + 9 + 8 + 7 + 9 + 10 = 72 \div 8 = 9$ rods mean breadth; then $9 \times 80 = 720 \div 160 = 4$ acres and 80 rods, or $4\frac{1}{2}$ acres. *Ans.*

Where the field is long, and the sides and ends crooked and irregular.



RULE.—Find the mean breadth in rods by the foregoing rule, and proceed in like manner to find the mean length in rods; then multiply the mean length by the mean breadth, and divide the product by 160, and the quotient will be the number of acres.

EXAMPLE.—What is the area in acres of a field of irregular sides and ends, the various breadths of which are as follows, viz.: 9, 6, 7, 8, 10 and 8 rods, and the lengths as follows, viz.: 50, 40, 30 and 40 rods?

SOLUTION.— $9 + 6 + 7 + 8 + 10 + 8 = 48 \div 6 = 8$ rods mean breadth.

$50 + 40 + 30 + 40 = 160 \div 4 = 40$ rods mean length.

Then $40 \times 8 = 320 \div 160 = 2$ acres. *Ans.*

Where the field is a circle.

RULE.—Take the diameter in rods, and find the area of the circle in the table of circles on page 298, and divide it by 160, and the quotient will be the number of acres.

EXAMPLE.—What is the area in acres of a circular field 22 rods in diameter?

SOLUTION.—380, area of circle, $\div 160 = 2$ acres and 80 rods, or $2\frac{1}{2}$ acres. *Ans.*

An acre of land is contained in a plot,

3 by $53\frac{1}{8}$ rods	7 by $22\frac{2}{7}$ rods	10 by 16 rods
4 by 40 “	8 by 20 “	11 by $14\frac{6}{11}$ “
5 by 32 “	9 by $17\frac{2}{3}$ “	12 by $13\frac{1}{3}$ “
6 by $26\frac{2}{3}$ “		

12 rods 10 feet and $8\frac{1}{2}$ inches square make an acre.

It is often desirable, for experimental and other purposes, for a farmer to lay off small portions of his ground. To enable him to do so, we have compiled the following:
 TABLE, *showing the square feet and the feet square of the fractions of an acre.*

Fractions of an acre.	Square feet.	Feet square.	Fractions of an acre.	Square feet.	Feet square.
$\frac{1}{16}$	2722 $\frac{1}{2}$	52 $\frac{1}{2}$	$\frac{1}{2}$	21780	147 $\frac{1}{2}$
$\frac{1}{8}$	5445	73 $\frac{1}{2}$	1	43560	208 $\frac{1}{2}$
$\frac{3}{8}$	10890	104 $\frac{1}{2}$	2	87120	295 $\frac{1}{2}$
$\frac{1}{4}$	14520	120 $\frac{1}{2}$			

TABLE, *showing the number of hills or plants on an acre of land, for any distance apart, from 10 inches to 6 feet—the lateral and longitudinal distances being unequal.*

	10 in.	12 in.	15 in.	18 in.	20 in.	2 ft.	2 $\frac{1}{2}$ ft.	3 ft.	3 $\frac{1}{2}$ ft.	4 ft.	4 $\frac{1}{2}$ ft.	5 ft.	5 $\frac{1}{2}$ ft.	6 ft.
10 in.	62726													
12 "	52272	43560												
15 "	41817	34848	27878											
18 "	34848	29040	23232	19360										
20 "	31363	26136	20908	17424	15681									
2 feet	26136	21780	17424	14520	13068	10890								
2 $\frac{1}{2}$ "	20908	17424	13939	11616	10454	8712	6969							
3 "	17424	14520	11616	9680	8712	7260	5808	4840						
3 $\frac{1}{2}$ "	14935	12446	9953	8297	7467	6223	4976	4148	3565					
4 "	13068	10890	8712	7260	6534	5445	4356	3630	3111	2722				
4 $\frac{1}{2}$ "	11616	9680	7744	6453	5808	4840	3872	3226	2767	2420	2151			
5 "	10454	8712	6969	5808	5227	4356	3484	2904	2489	2178	1936	1742		
5 $\frac{1}{2}$ "	9504	7920	6336	5280	4752	3960	3168	2640	2263	1980	1760	1584	1440	
6 "	8712	7260	5808	4840	4356	3630	2904	2420	2074	1865	1615	1452	1320	1210

EXPLANATION.—Find the distance between your plants or hills the widest way in the left hand column, then trace the line in which it stands to the right, until it intersects the column headed by the number that expresses the distance of the narrow way, where you will find the number sought.

EXAMPLE.—The rows of corn in a corn-field are $5\frac{1}{2}$ feet apart, and the plants 20 inches apart, in drill or hill; required, the number of hills or plants in an acre?

SOLUTION.—Find $5\frac{1}{2}$ feet (the distance of the rows apart), in the left hand column, then trace the line along unto the column headed by 20 inches (the distance of the plants or hills apart), and you have 4752. *Ans.*

TABLE, *showing the number of plants, hills, or trees contained in an acre at equal distances apart, from 3 inches up to 66 feet.*

Distance apart.	No. of plants.	Distance apart.	No. of plants.
3 inches by 3 inches	696,960	6 feet by 6 feet	1,210
4 " by 4 "	392,040	6½ " by 6½ "	1,031
6 " by 6 "	174,240	7 " by 7 "	881
9 " by 9 "	77,440	8 " by 8 "	680
1 foot by 1 foot	43,560	9 " by 9 "	537
1½ feet by 1½ feet	19,360	10 " by 10 "	435
2 " by 1 foot	21,780	11 " by 11 "	360
2 " by 2 feet	10,890	12 " by 12 "	302
2½ " by 2½ "	6,960	13 " by 13 "	257
3 " by 1 foot	14,520	14 " by 14 "	222
3 " by 2 feet	7,260	15 " by 15 "	193
3 " by 3 "	4,840	16 " by 16 "	170
3½ " by 3½ "	3,555	16½ " by 16½ "	160
4 " by 1 foot	10,890	17 " by 17 "	150
4 " by 2 feet	5,445	18 " by 18 "	134
4 " by 3 "	3,630	19 " by 19 "	120
4 " by 4 "	2,722	20 " by 20 "	108
4½ " by 4½ "	2,151	25 " by 25 "	69
5 " by 1 foot	8,712	30 " by 30 "	48
5 " by 2 feet	4,356	33 " by 33 "	40
5 " by 3 "	2,904	40 " by 40 "	27
5 " by 4 "	2,178	50 " by 50 "	17
5 " by 5 "	1,742	60 " by 60 "	12
5½ " by 5½ "	1,417	66 " by 66 "	10

GOVERNMENT LAND MEASURE.

A township is 6 miles square, and contains 36 sections, or 23,040 acres.

A section is 1 mile square, and contains 640 acres.


A quarter-section is half a mile square, and contains 160 acres.

A half quarter-section is half a mile long, almost universally north and south, and one-fourth of a mile wide, and contains 80 acres.

A quarter quarter-section is one-fourth of a mile square, and contains 40 acres. It is the smallest sized tract, except fractions, sold by the government.



MEASUREMENT OF HAY.



There is no accurate mode of measuring hay but by weighing it. This, on account of its bulk and character, is very difficult, unless it is baled or otherwise compacted. This difficulty has led farmers to estimate the weight by the bulk or cubic contents, a mode which, from the nature of the commodity, is only approximately correct. Some kinds of hay are light, while others are heavy, their equal bulks varying in weight. But for all ordinary farming purposes of estimating the amount of hay in meadows, mows, and stacks, the following rules will be found sufficient.

As nearly as can be ascertained, 25 cubic yards of average meadow hay, in windrows, make a ton.

When well settled in mows or stacks, 15 or 18 cubic yards make a ton.

When taken out of mows or old stacks, and loaded on wagons, 20 or 25 cubic yards make a ton.

Twenty or twenty-five cubic yards of clover, when dry, make a ton.

To find the number of tons of meadow hay raked into windrows.

RULE.—Multiply the length of the windrow in yards by the width in yards, and that product by the height in yards, and divide by 25; the quotient will be the number of tons in the windrow.

EXAMPLE.—How many tons of hay in a windrow 40 yards long by 2 wide and 2 high?

SOLUTION.— $40 \times 2 \times 2 = 160 \div 25 = 6\frac{2}{5}$. Ans.


To find the number of tons of hay in a mow.

RULE.—Multiply the length in yards by the height in yards, and that by the width in yards, and divide the product by 15; the quotient will be the number of tons.

EXAMPLE.—How many tons of well-settled hay in a mow 10 yards long by 6 wide and 8 high?

SOLUTION.— $10 \times 6 \times 8 = 480 \div 15 = 32$ tons. Ans.

To find the number of tons of hay in old stacks.



RULE.—Find the area in square yards of the base in the table of the areas of circles on page 298, or by the rule given on page 296; then multiply the area of the base by half the altitude of the stack in yards, and divide the product by 15; the quotient will be the number of tons.

EXAMPLE.—How many tons of hay in a circular stack, whose diameter at the base is 8 yards, and height 9 yards?

SOLUTION.—50.265, area of base in sq. yards, $\times 4\frac{1}{2}$, half the altitude, $= 226.192 \div 15 = 15.079$ tons. *Ans.*

To find the number of tons in long square stacks.

RULE.—Multiply the length in yards by the width in yards, and that by half the altitude in yards, and divide the product by 15; the quotient will be the number of tons.

EXAMPLE.—How many tons of hay in a square stack 10 yards long, 5 wide, and 9 high?

SOLUTION.— $10 \times 5 \times 4\frac{1}{2} = 225 \div 15 = 15$ tons. *Ans.*

To find the number of tons of hay when taken out of mows or old stacks.

RULE.—Multiply the length of the load in yards by the width in yards, and that by the height in yards, and divide the product by 20; the quotient will be the number of tons.

EXAMPLE.—How many tons of hay taken from an old stack, in a load 6 yards long by 3 wide and 3 high?

SOLUTION.— $6 \times 3 \times 3 = 54 \div 20 = 2\frac{7}{10}$ tons. *Ans.*

These estimates are for medium sized mows or stacks. If the hay is piled to a great height, as it often is where horse hay-forks are used, the row will be much heavier per cubic yard.

TABLE, showing the price per cwt. of hay, at given prices per ton.

Price per ton.	$\frac{1}{2}$ hundred.		1 hundred.	2 hundred	3 hundred.	4 hundred.	5 hundred.	6 hundred.	7 hundred.	8 hundred.	9 hundred.	10 hundred.	11 hundred.
\$	cts.	cts.	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts
4	10	20	40	60	80	1.00	1.20	1.40	1.60	1.80	2.00	2.20	2.40
5	12	25	50	75	1.00	1.25	1.50	1.75	2.00	2.25	2.50	2.75	3.00
6	15	30	60	90	1.20	1.50	1.80	2.10	2.40	2.70	3.00	3.30	3.60
7	17	35	70	1.05	1.40	1.75	2.10	2.45	2.80	3.15	3.50	3.85	4.20
8	20	40	80	1.20	1.60	2.00	2.40	2.80	3.20	3.60	4.00	4.40	4.80
9	22	45	90	1.35	1.80	2.25	2.70	3.15	3.60	4.05	4.50	4.95	5.40
10	25	50	1.00	1.50	2.00	2.50	3.00	3.50	4.00	4.50	5.00	5.50	6.00
11	27	55	1.10	1.65	2.20	2.75	3.30	3.85	4.40	4.95	5.50	6.00	6.60
12	30	60	1.20	1.80	2.40	3.00	3.60	4.20	4.80	5.40	6.00	6.60	7.20
13	32	65	1.30	1.95	2.60	3.25	3.90	4.55	5.20	5.85	6.50	7.15	7.80
14	35	70	1.40	2.10	2.80	3.50	4.20	4.90	5.60	6.30	7.00	7.70	8.40
15	37	75	1.50	2.25	3.00	3.75	4.50	5.25	6.00	6.75	7.50	8.25	9.00

Table continued.

Price per ton.	12 hundred.	13 hundred.	14 hundred.	15 hundred.	16 hundred.	17 hundred.	18 hundred.	19 hundred.	20 hundred.
\$	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts	\$ cts
4	2.40	2.60	2.80	3.00	3.20	3.40	3.60	3.80	4.00
5	3.00	3.25	3.50	3.75	4.00	4.25	4.50	4.75	5.00
6	3.60	3.90	4.20	4.50	4.80	5.10	5.40	5.70	6.00
7	4.20	4.55	4.90	5.25	5.60	5.95	6.30	6.65	7.00
8	4.80	5.20	5.60	6.00	6.40	6.80	7.20	7.60	8.00
9	5.40	5.85	6.30	6.75	7.20	7.65	8.10	8.55	9.00
10	6.00	6.50	7.00	7.50	8.00	8.50	9.00	9.50	10.00
11	6.50	7.15	7.70	8.25	8.80	9.35	9.90	10.45	11.00
12	7.20	7.80	8.40	9.00	9.60	10.20	10.80	11.40	12.00
13	7.80	8.45	9.10	9.75	10.40	11.05	11.70	12.35	13.00
14	8.40	9.10	9.80	10.50	11.20	11.90	12.60	13.30	14.00
15	9.00	9.75	10.50	11.25	12.00	12.75	13.50	14.25	15.00

An easy mode of ascertaining the value of a given number of lbs. of hay, at a given price per ton of 2000 lbs.

RULE.—Multiply the number of pounds of hay (coal, or anything else which is bought and sold by the ton) by one half the price per ton, pointing off three figures from the right hand; the remaining figures will be the price of the hay (or any article by the ton).

EXAMPLE.—What will be the cost of 658 lbs. of hay, at \$7.50 per ton?

SOLUTION.—\$7.50 divided by 2 equals \$3.75, by which multiply the number of pounds, thus:

$$\begin{array}{r}
 658 \\
 \times 3.75 \\
 \hline
 3290 \\
 4606 \\
 1974 \\
 \hline
 \end{array}$$

\$2.46750. *Ans.*

NOTE.—The principle in this rule is the same as in interest—dividing the price by two gives us the price of half a ton, or 1000 lbs.; and pointing off three figures to the right is dividing by 1000.


A *truss* of hay, new, is 60 lbs.; old, 56 lbs.; straw, 40 lbs.

A *load* of hay is 36 trusses.

A *bale* of hay is 300 lbs.



TO MEASURE CORN ON THE COB IN CRIBS.



When the crib is equilateral.

RULE.—Multiply the length in inches by the breadth in inches, and that again by the height in inches, and divide the product by 2748 (the number of cubic inches in a heaped bushel), and the quotient will be the number of heaped bushels of ears. Take two-thirds of the quotient for the number of bushels of shelled corn.

EXAMPLE.—Required the number of bushels of shelled corn contained in a crib of ears, 15 feet long by 5 feet wide and 10 feet high?

SOLUTION.—180 in., length, \times 60 in., width, \times 120 in.,
3*

height, $=1296000 \div 2748 = 471.6$ heaped bushels, $\frac{2}{3}$ of which is 314.6 bushels shelled. *Ans.*

NOTE.—The above rule assumes that three heaping half bushels of ears make one struck bushel of shelled corn. This proportion has been adopted upon the authority of the major part of our best agricultural journals. Nevertheless, some journals claim that two heaping bushels of ears to one of shelled corn is a more correct proportion, and it is the custom in many parts of the country to buy and sell at that rate. Of course, much will depend upon the kind of corn, the shape of the ear, the size of the cob, &c. Some samples are to be found, three heaping half bushels of which will even overrun one bushel shelled; while others again are to be found, two bushels of which will fall short of one bushel shelled. Every farmer must judge for himself, from the sample on hand, whether to allow one and a half or two bushels ears to one of shelled corn. In either case, it is only an approximate measurement, but sufficient for all ordinary purposes of estimation. The only true way of measuring all such products is by weight.

When the crib is flared at the sides.

RULE.—Multiply half the sum of the top and bottom widths in inches by the perpendicular height in inches, and that again by the length in inches, and divide the product by 2748, and the quotient will be the number of heaped bushels of ears. Take two-thirds of the quotient for the number of bushels of shelled corn.

EXAMPLE.—Required, the number of bushels of shelled corn contained in a crib of ears 4 feet wide at the bottom, 8 feet at the top, 10 feet in perpendicular height, and 15 feet long?

SOLUTION.—48 inches, bottom width, +96 inches, top width, $=144 \div 2 = 72 \times 120$ inches perpendicular height, \times 180 inches length, $=1555200 \div 2748 = 565.9$ bus. ears, $\frac{2}{3}$ of which is 377.28 bus. shelled corn. *Ans.*

NOTE.—A barrel of corn is 5 bushels shelled. By this latter measure crops are estimated, and corn bought and sold throughout most of the Southern and Western States. At New Orleans a barrel of corn is a flour-barrel full of ears. In some parts of the West, it is common to count 100 ears to the bushel.



MEASUREMENT OF GRAIN IN GRANARIES.



To find the number of bushels of grain in a granary.

RULE.—Multiply the length in inches by the breadth in inches, and that again by the depth in inches, and divide the product by 2150 (the number of cubic inches in a bushel), and for heaped bushels by 2748, and the quotient will be the answer.

EXAMPLE.—Given a granary 9 feet long by 4 wide and 6 deep. How many bushels will it contain?

SOLUTION.—108 inches length, \times 48 inches width, \times 72 in. depth, $= 373248 \div 2150 = 173.65$ bus. *Ans.*

MEASUREMENT OF TIMBER.



The unit of board measure is a superficial foot 1 inch thick.

Besides inch-boards, plank and scantling are usually bought and sold by board measure.

Round, sawed, or hewn timber is bought and sold by the cubic foot.

Pine and spruce spars, from 10 to $4\frac{1}{2}$ inches in diameter inclusive, are measured by taking the diameter, clear of bark, at one-third of their length from the large end.

Spars are usually purchased by the inch diameter; all under 4 inches are considered *poles*.


Spruce spars of 7 inches and less, should have 5 feet in length for every inch in diameter.

WOOD MEASURE.

To ascertain the contents or number of cords in a given pile of wood.

RULE.—Multiply the length by the width, and that product by the height, which will give you the number of cubic feet. Divide that product by 128, and the quotient will be the number of cords.

A pile of wood 4 feet wide, and 4 feet high, and 8 feet long, contains 1 cord; and a cord foot is 1 foot in length of such a pile, thus :



BOARD MEASURE.

To ascertain the contents (board measure) of boards, scantling, and plank.

RULE.—Multiply the breadth in inches by the thickness in inches, and that by the length in feet, and divide the product by 12, and the quotient will be the contents.

TABLE, showing the contents of inch-boards from 3 to 30 in. broad, and from 4 to 24 feet long.

Ins.	LENGTH IN FEET.																							
	4.	5.	6.	7.	8.	9.	10.	11.	12.	13.	14.	15.	16.	17.	18.	19.	20.	21.	22.	23.	24.			
3	1.00	1.03	1.06	1.09	2.00	2.03	2.06	2.09	3.00	3.03	3.06	3.09	4.00	4.03	4.06	4.09	5.00	5.03	5.06	5.09	6.00			
4	1.04	1.08	2.00	2.04	2.08	3.00	3.04	3.08	4.00	4.04	4.08	5.00	5.04	5.07	6.00	6.04	6.08	7.00	7.05	7.08	8.00			
5	1.08	2.01	2.06	2.11	3.04	3.09	4.02	4.07	5.00	5.05	5.10	6.03	6.08	7.01	7.03	7.11	8.04	8.09	9.02	9.07	10.00			
6	2.00	2.06	3.00	3.06	4.00	4.06	5.00	5.06	6.00	6.06	7.00	7.06	8.00	8.06	9.00	9.06	10.00	10.06	11.00	11.06	12.00			
7	2.04	2.11	3.06	4.01	4.08	5.03	5.10	6.05	7.00	7.07	8.02	8.09	9.04	9.11	10.06	11.01	11.08	12.03	12.10	13.05	14.00			
8	2.08	3.04	4.00	4.08	5.04	6.00	6.08	7.04	8.00	8.08	9.04	10.00	10.08	11.04	12.00	12.08	13.04	14.00	14.08	15.04	16.00			
9	3.00	3.09	4.06	5.03	6.00	6.09	7.06	8.03	9.00	9.09	10.06	11.03	12.00	12.09	13.06	14.03	15.00	15.09	16.06	17.03	18.00			
10	3.04	4.02	5.00	5.10	6.08	7.06	8.04	9.02	10.00	10.10	11.08	12.06	13.04	14.02	15.00	15.10	16.08	17.06	18.04	19.02	20.00			
11	3.08	4.07	5.06	6.05	7.04	8.03	9.02	10.01	11.00	11.11	12.10	13.09	14.08	15.07	16.06	17.05	18.04	19.03	20.00	21.01	22.00			
12	4.00	5.00	6.00	7.00	8.00	9.00	10.00	11.00	12.00	13.00	14.00	15.00	16.00	17.00	18.00	19.00	20.00	21.00	22.00	23.00	24.00			
13	4.04	5.05	6.06	7.07	8.08	9.09	10.10	11.11	12.12	13.13	14.14	15.15	16.16	17.17	18.18	19.19	20.20	21.21	22.22	23.23	24.24			
14	4.08	5.10	7.00	8.02	9.04	10.06	11.08	12.10	14.00	15.02	16.04	17.06	18.08	19.10	21.00	22.02	23.04	24.06	25.08	26.10	28.00			
15	5.00	6.03	7.06	8.09	10.00	11.03	12.06	13.09	15.00	16.03	17.06	18.09	20.00	21.03	22.06	23.09	25.00	26.03	27.06	28.09	30.00			
16	5.04	6.08	8.00	9.04	10.08	12.00	13.04	14.08	16.00	17.04	18.08	20.00	21.04	22.08	24.00	25.04	26.08	28.00	29.04	30.08	32.00			
17	5.08	7.01	8.06	9.11	11.04	12.09	14.02	15.07	17.00	18.05	19.10	21.03	22.08	24.01	25.06	26.11	28.04	29.09	31.02	32.07	34.00			
18	6.00	7.06	9.00	10.06	12.00	13.06	15.00	16.06	18.00	19.06	21.00	22.06	24.00	25.06	27.00	28.06	30.00	31.06	33.00	34.06	36.00			
19	6.04	7.11	9.06	11.01	12.08	14.03	15.10	17.05	19.00	20.07	22.02	23.09	25.04	26.11	28.06	30.01	31.08	33.03	34.10	36.05	38.00			
20	6.08	8.04	10.00	11.08	13.04	15.00	16.08	18.04	20.00	21.08	23.04	25.00	26.08	28.04	30.00	31.08	33.04	35.00	36.08	38.04	40.00			
21	7.00	8.09	10.06	12.03	14.00	15.09	17.06	19.03	21.00	22.09	24.06	26.03	28.00	29.09	31.06	32.03	35.00	36.09	38.06	40.03	42.00			
22	7.04	9.02	11.00	12.10	14.08	16.05	18.04	20.02	22.00	23.10	25.08	27.06	29.04	31.02	33.00	34.10	36.08	38.06	40.04	42.03	44.00			
23	7.08	9.07	11.06	13.05	15.04	17.03	19.00	21.01	23.00	24.11	26.10	28.09	30.08	32.07	34.06	36.05	38.04	40.03	42.02	44.01	46.00			
24	8.00	10.00	12.00	14.00	16.00	18.00	20.00	22.00	24.00	26.00	28.00	30.00	32.00	34.00	36.00	38.00	40.00	42.00	44.00	46.00	48.00			
25	8.04	10.05	12.06	14.07	16.08	18.09	20.10	22.11	25.00	27.01	29.02	31.03	33.04	35.05	37.06	39.07	41.08	43.09	45.10	47.11	50.00			
26	8.08	10.10	13.00	15.02	17.04	19.06	21.08	23.10	26.00	28.02	30.04	32.06	34.08	36.10	39.00	41.02	43.04	45.06	47.08	49.10	52.00			
27	9.00	11.03	13.06	15.09	18.00	20.03	22.06	24.09	27.00	29.03	31.06	33.09	36.00	38.03	40.06	42.09	44.08	46.08	49.00	51.04	54.00			
28	9.04	11.08	14.00	16.04	18.08	21.00	23.04	25.08	28.00	30.04	32.08	35.00	37.04	39.08	42.00	44.04	46.08	49.00	51.06	53.00	56.00			
29	9.08	12.01	14.06	16.11	19.04	21.09	24.02	26.07	29.00	31.05	33.10	36.03	38.08	41.01	43.06	45.11	48.04	50.09	53.02	55.07	58.00			
30	10.00	12.06	15.00	17.06	20.00	22.06	25.00	27.06	30.00	32.06	35.00	37.06	40.00	42.06	45.00	47.06	50.00	52.06	55.00	57.06	60.00			

BOARD MEASURE.

EXPLANATION.—First find the width in inches in the left hand column, and the length in feet at the heads of the other columns; then trace the two until they meet, and the figures so found will express the contents in feet and inches.

ROUND TIMBER.



Round timber when squared is estimated to lose *one-fifth*; hence a ton of round timber is said to contain only 40 cubic feet.

Sawed lumber, as joists, plank, and scantlings, are now generally bought and sold by *board measure*. The dimensions of a *foot* of board measure is 1 foot long, 1 ft. wide, and 1 inch thick.

To measure round timber.

RULE.—Take the girth in inches at both the large and small ends, add them, and divide their sum by two for the mean girth; then multiply the length in feet by the square of one-fourth of the mean girth in inches, divide the product by 144, and the quotient will be the contents in cubic feet.

EXAMPLE.—What are the cubic contents of a round log 12 feet long, 54 inches girth at the large end, and 34 at the small end?

SOLUTION.— $54 + 34 = 88 \div 2 = 44$ inches, mean girth.

Then $12 \text{ length} \times 121 \text{ inches (the square of } \frac{1}{4} \text{ mean girth)} = 1452 \div 144 = 10\frac{1}{2}$ cubic feet. *Ans.*

SQUARE TIMBER.

To measure square timber.

RULE.—Multiply the breadth in inches by the depth in inches, and that by the length in feet, and divide the product by 144, and the quotient will be the contents in cubic feet.

EXAMPLE.—What is the cubic contents of a square log 12 feet long by 20 inches broad and 18 deep?

SOLUTION.— $20 \times 18 = 360 \times 12 = 4320 \div 144 = 30$ cubic feet. *Ans.*

PLANK MEASURE.

TABLE, showing the contents (board measure) of planks of various dimensions.

Feet long	1 by 10	1 by 11	1 by 12	1 by 13	1 by 14	1 by 15	1 by 16	1 by 17	1 by 18	1 by 19
24	35	39	42	45	49	52	56	59	63	66
25	36	40	44	47	51	55	59	62	66	69
25	38	42	45	49	53	57	61	64	68	72
27	39	43	47	51	55	59	63	67	71	74
28	41	45	49	53	57	61	65	69	73	77
29	42	47	51	55	59	63	68	72	76	80
30	43	48	52	57	61	65	70	74	79	83
31	45	50	54	59	63	68	72	77	81	85
32	47	51	56	61	65	70	75	79	84	88
33	48	53	58	63	67	72	77	82	87	91
34	49	55	59	64	69	74	79	84	89	93
35	51	56	61	66	71	76	82	87	92	95
36	52	58	63	68	73	79	84	89	94	98
37	54	59	65	70	75	81	86	92	97	101
38	55	61	66	72	77	83	89	94	100	104
39	57	63	68	74	80	85	91	97	102	107
40	58	64	70	76	82	87	93	99	105	109
41	60	67	72	78	84	90	96	102	108	112
42	61	69	73	80	86	92	98	104	110	115
43	62	71	75	82	88	94	100	108	113	118
44	64	73	77	83	90	96	103	109	115	121
45	65	74	79	85	92	98	105	112	118	123
46	67	76	80	87	94	101	107	114	121	126
47	68	78	82	89	96	103	110	117	123	129
48	70	79	84	91	98	105	112	119	125	131
49	71	80	86	93	100	107	114	121	128	134
50	73	82	88	94	102	109	117	124	131	136
51	74	83	89	97	104	112	119	126	133	140
52	76	84	91	99	106	114	121	129	136	141

Table continued.

24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	255	260	265	270	275	280	285	290	295	300	305	310	315	320	325	330	335	340	345	350	355	360	365	370	375	380	385	390	395	400	405	410	415	420	425	430	435	440	445	450	455	460	465	470	475	480	485	490	495	500	505	510	515	520	525	530	535	540	545	550	555	560	565	570	575	580	585	590	595	600	605	610	615	620	625	630	635	640	645	650	655	660	665	670	675	680	685	690	695	700	705	710	715	720	725	730	735	740	745	750	755	760	765	770	775	780	785	790	795	800	805	810	815	820	825	830	835	840	845	850	855	860	865	870	875	880	885	890	895	900	905	910	915	920	925	930	935	940	945	950	955	960	965	970	975	980	985	990	995	1000
24	65	70	75	80	85	90	95	100	105	110	115	120	125	130	135	140	145	150	155	160	165	170	175	180	185	190	195	200	205	210	215	220	225	230	235	240	245	250	25																																																																																																																																																					

Table continued.

Feet. long.	3 by 24	3 by 25	3 by 26	3 by 27	3 by 28	3 by 29	3 by 30	3 by 31	3 by 32	3 by 33	3 by 34	3 by 35	3 by 36	3 by 37	3 by 38	3 by 39	3 by 40	3 by 41	3 by 42	3 by 43	3 by 44	3 by 45	3 by 46	3 by 47	3 by 48	3 by 49	3 by 50	3 by 51	3 by 52	
21	144	150	156	162	168	174	180	186	192	198	204	210	216	222	228	234	240	246	252	258	264	270	276	282	288	294	300	306	312	318
22	145	151	157	163	169	175	181	187	193	199	205	211	217	223	229	235	241	247	253	259	265	271	277	283	289	295	301	307	313	319
23	146	152	158	164	170	176	182	188	194	200	206	212	218	224	230	236	242	248	254	260	266	272	278	284	290	296	302	308	314	320
24	147	153	159	165	171	177	183	189	195	201	207	213	219	225	231	237	243	249	255	261	267	273	279	285	291	297	303	309	315	
25	148	154	160	166	172	178	184	190	196	202	208	214	220	226	232	238	244	250	256	262	268	274	280	286	292	298	304	310	316	
26	149	155	161	167	173	179	185	191	197	203	209	215	221	227	233	239	245	251	257	263	269	275	281	287	293	299	305	311	317	
27	150	156	162	168	174	180	186	192	198	204	210	216	222	228	234	240	246	252	258	264	270	276	282	288	294	300	306	312	318	
28	151	157	163	169	175	181	187	193	199	205	211	217	223	229	235	241	247	253	259	265	271	277	283	289	295	301	307	313	319	
29	152	158	164	170	176	182	188	194	200	206	212	218	224	230	236	242	248	254	260	266	272	278	284	290	296	302	308	314	320	
30	153	159	165	171	177	183	189	195	201	207	213	219	225	231	237	243	249	255	261	267	273	279	285	291	297	303	309	315	321	
31	154	160	166	172	178	184	190	196	202	208	214	220	226	232	238	244	250	256	262	268	274	280	286	292	298	304	310	316	322	
32	155	161	167	173	179	185	191	197	203	209	215	221	227	233	239	245	251	257	263	269	275	281	287	293	299	305	311	317	323	
33	156	162	168	174	180	186	192	198	204	210	216	222	228	234	240	246	252	258	264	270	276	282	288	294	300	306	312	318	324	
34	157	163	169	175	181	187	193	199	205	211	217	223	229	235	241	247	253	259	265	271	277	283	289	295	301	307	313	319	325	
35	158	164	170	176	182	188	194	200	206	212	218	224	230	236	242	248	254	260	266	272	278	284	290	296	302	308	314	320	326	
36	159	165	171	177	183	189	195	201	207	213	219	225	231	237	243	249	255	261	267	273	279	285	291	297	303	309	315	321	327	
37	160	166	172	178	184	190	196	202	208	214	220	226	232	238	244	250	256	262	268	274	280	286	292	298	304	310	316	322	328	
38	161	167	173	179	185	191	197	203	209	215	221	227	233	239	245	251	257	263	269	275	281	287	293	299	305	311	317	323	329	
39	162	168	174	180	186	192	198	204	210	216	222	228	234	240	246	252	258	264	270	276	282	288	294	300	306	312	318	324	330	
40	163	169	175	181	187	193	199	205	211	217	223	229	235	241	247	253	259	265	271	277	283	289	295	301	307	313	319	325	331	
41	164	170	176	182	188	194	200	206	212	218	224	230	236	242	248	254	260	266	272	278	284	290	296	302	308	314	320	326	332	
42	165	171	177	183	189	195	201	207	213	219	225	231	237	243	249	255	261	267	273	279	285	291	297	303	309	315	321	327	333	
43	166	172	178	184	190	196	202	208	214	220	226	232	238	244	250	256	262	268	274	280	286	292	298	304	310	316	322	328	334	
44	167	173	179	185	191	197	203	209	215	221	227	233	239	245	251	257	263	269	275	281	287	293	299	305	311	317	323	329	335	
45	168	174	180	186	192	198	204	210	216	222	228	234	240	246	252	258	264	270	276	282	288	294	300	306	312	318	324	330	336	
46	169	175	181	187	193	199	205	211	217	223	229	235	241	247	253	259	265	271	277	283	289	295	301	307	313	319	325	331	337	
47	170	176	182	188	194	200	206	212	218	224	230	236	242	248	254	260	266	272	278	284	290	296	302	308	314	320	326	332	338	
48	171	177	183	189	195	201	207	213	219	225	231	237	243	249	255	261	267	273	279	285	291	297	303	309	315	321	327	333	339	
49	172	178	184	190	196	202	208	214	220	226	232	238	244	250	256	262	268	274	280	286	292	298	304	310	316	322	328	334	340	
50	173	179	185	191	197	203	209	215	221	227	233	239	245	251	257	263	269	275	281	287	293	299	305	311	317	323	329	335	341	
51	174	180	186	192	198	204	210	216	222	228	234	240	246	252	258	264	270	276	282	288	294	300	306	312	318	324	330	336	342	
52	175	181	187	193	199	205	211	217	223	229	235	241	247	253	259	265	271	277	283	289	295	301	307	313	319	325	331	337	343	

Table continued.

L. Feet.	Diam. 12	Diam. 13	Diam. 14	Diam. 15	Diam. 16	Diam. 17	Diam. 18	Diam. 19	Diam. 20	Diam. 21	Diam. 22	Diam. 23	Diam. 24	Diam. 25	Diam. 26	Diam. 27	Diam. 28	Diam. 29
10	40	61	72	89	99	116	133	150	175	191	209	235	252	287	313	345	363	381
11	54	67	79	98	109	127	147	165	192	209	230	259	278	315	344	377	400	419
12	59	73	86	107	119	139	160	180	210	228	251	283	303	344	375	411	436	457
13	64	79	93	116	129	151	173	195	227	247	272	306	328	373	408	446	473	495
14	69	85	100	125	139	162	187	210	245	266	292	336	355	401	439	479	509	533
15	74	91	107	134	149	173	200	225	262	285	313	353	379	430	469	514	545	571
16	79	97	114	142	159	185	213	241	280	304	334	377	404	459	500	548	582	609
17	81	103	122	151	168	196	227	255	297	322	356	400	429	478	531	582	618	647
18	88	109	129	160	178	208	240	271	315	342	376	424	454	510	562	610	654	685
19	93	116	136	169	188	219	253	285	332	361	397	447	480	545	594	650	692	723
20	9	122	143	177	198	232	267	300	350	380	418	470	505	573	625	684	728	761
21	103	128	150	187	208	243	280	315	368	399	439	495	530	602	656	719	764	800
22	108	134	157	196	218	255	293	330	385	418	460	518	555	631	688	753	800	838
23	113	140	164	205	228	266	307	345	403	437	480	542	571	659	719	787	837	876
24	118	146	172	214	238	278	320	360	420	455	501	566	606	688	756	821	873	914
25	123	152	179	223	248	289	333	375	438	475	522	589	631	717	781	856	910	952

L. Feet.	Diam. 30	Diam. 31	Diam. 32	Diam. 33	Diam. 34	Diam. 35	Diam. 36	Diam. 37	Diam. 38	Diam. 39	Diam. 40	Diam. 41	Diam. 42	Diam. 43
10	411	444	460	490	500	547	577	644	669	70	752	795	840	872
11	451	448	506	539	559	603	634	708	734	770	828	874	924	959
12	493	532	552	588	600	657	692	772	801	840	903	954	1007	1046
13	534	576	598	637	650	712	750	836	868	910	978	1033	1091	1135
14	575	622	644	686	706	766	807	901	934	980	1053	1113	1175	1222
15	616	666	690	735	750	821	865	965	1001	1050	1129	1192	1259	1309
16	657	710	733	784	800	876	923	1029	1068	1120	1204	1272	1343	1396
17	698	755	782	833	850	931	980	1094	1134	1190	1279	1351	1427	1485
18	739	799	828	882	900	985	1038	1158	1201	1260	1354	1431	1511	1571
19	78	843	871	931	950	1040	1096	1222	1268	1330	1430	1510	1595	1658
20	821	888	920	980	1000	1095	1152	1287	1335	1400	1505	1590	1679	1745
21	863	932	966	1029	1050	1150	1210							
22	904	976	1012	1078	1100	1204	1268							
23	945	1021	1058	1127	1150	1259	1323							
24	986	1065	1104	1176	1200	1314	1380							
25	1027	1109	1150	1225	1250	1369	1438							

EXPLANATION.—Find the length of the log in feet in the left hand column, and its mean diameter in inches (found by adding the two end diameters, and dividing their sum by two) at the heads of the other columns, and trace them

until they meet, and the figures so found will express the number of feet board measure of inch-boards the log will furnish.

SCANTLING MEASURE.

TABLE, showing the contents (board measure) of scantling of various dimensions.

Feet Long.	2 by 2	2 by 3	2 by 4	2 by 5	2 by 6	2 by 7	2 by 8	2 by 9	2 by 10
1	0.4	0.6	0.8	0.10	1.	1.2	1.4	1.6	1.8
2	0.8	1.	1.4	1.8	2.	2.4	2.8	3.	3.4
3	1.4	1.6	2.2	2.6	3.	3.6	4.	4.6	5.
4	1.8	2.	2.8	3.4	4.	4.8	5.4	6.	6.8
5	2.4	3.	3.4	4.2	5.	5.10	6.8	7.6	8.4
6	2.8	3.6	4.4	5.	6.	7.	8.	9.	10.
7	3.2	4.	4.8	5.10	7.	8.2	9.4	10.6	11.8
8	3.6	4.6	5.4	6.8	8.	9.4	10.8	12.	13.4
9	4.	5.	6.	7.6	9.	10.6	12.	13.6	15.
10	4.4	5.6	6.8	8.4	10.	11.8	13.4	15.	16.8
11	4.8	6.	7.4	9.2	11.	12.10	14.8	16.6	18.4
12	5.	6.6	8.	10.	12.	14.	16.	18.	20.
13	5.4	7.	8.8	10.10	13.	15.2	17.4	19.6	21.8
14	5.8	7.6	9.4	11.8	14.	16.4	18.8	21.	23.4
15	6.	8.	10.8	12.6	15.	17.6	20.	22.6	25.
16	6.4	8.6	11.4	13.4	16.	18.8	21.4	24.	26.8
17	6.8	9.	12.	14.2	17.	19.10	22.8	25.6	28.4
18	7.	9.6	12.8	15.10	18.	21.	24.	27.	30.
19	7.4	10.	13.4	16.8	19.	22.2	25.4	28.6	31.8
20	7.8	10.6	14.8	17.6	20.	23.4	26.8	30.	33.4
21	8.	11.	15.4	18.4	21.	24.6	28.	31.6	34.
22	8.4	11.6	16.	19.2	22.	25.8	29.4	33.	36.8
23	8.8	12.	16.8	20.10	23.	26.10	30.8	34.6	38.4
24	9.	12.6	17.4	21.	24.	27.	32.	36.	40.
25	9.4	13.	18.	22.6	25.	28.2	33.4	37.6	41.8
26	9.8	13.6	18.8	23.4	26.	30.4	34.8	39.	43.4
27	10.	14.6	19.4	24.2	27.	31.6	36.	40.6	45.
28	10.4	15.	20.	25.	28.	32.8	37.4	42.	46.8
29	10.8	15.6	20.8	26.	29.	33.10	38.8	43.6	48.4
30	11.	16.	21.4	27.	30.	34.	40.	45.	50.

Table continued.

Feet Long.	2 by 11	2½ by 5	2½ by 6	2½ by 7	2½ by 8	2½ by 9	2½ by 10	2½ by 11	2½ by 12	3 by 3	3 by 4
1	1.10	1.1	1.3	1.6	1.8	1.11	2.1	2.4	2.6	0.9	1.
2	3.8	2.1	2.6	2.11	3.4	3.9	4.2	4.7	5.	1.6	2.
3	5.6	3.2	3.9	4.5	5.	5.8	6.3	6.11	7.6	2.3	3.
4	7.4	4.2	5.	5.10	6.8	7.6	8.4	9.2	10.	3.	4.
5	9.2	5.3	6.3	7.4	8.4	9.5	10.5	11.6	12.6	3.9	5.
6	11.	6.3	7.6	8.9	10.	11.3	12.6	13.9	15.	4.6	6.
7	12.10	7.4	8.9	10.3	11.8	13.2	14.7	16.1	17.6	5.3	7.
8	14.8	8.4	10.	11.8	13.4	15.	16.8	18.4	20.	6.	8.
9	16.6	9.5	11.3	13.2	15.	16.11	18.9	20.8	22.6	6.9	9.
10	18.4	10.5	12.6	14.7	16.8	18.9	20.10	22.11	25.	7.6	10.
11	20.2	11.6	13.9	16.1	18.4	20.8	22.11	25.3	27.6	8.3	11.
12	22.	12.6	15.	17.6	20.	22.6	25.	27.6	30.	9.	12.
13	23.10	13.7	16.3	19.	21.8	24.5	27.1	29.10	32.6	9.9	13.
14	25.8	14.7	17.6	20.5	23.4	26.3	29.2	32.1	35.	10.6	14.
15	27.6	15.8	18.9	21.11	25.	28.2	31.3	34.4	37.6	11.3	15.
16	29.4	16.8	20.	23.4	26.8	30.	33.4	36.8	40.	12.	16.
17	31.2	17.9	21.3	24.10	28.4	31.11	35.5	39.	42.6	12.9	17.
18	33.	18.9	22.6	26.3	30.	33.9	37.6	41.3	45.	13.6	18.
19	31.10	19.10	23.9	27.9	31.8	35.8	39.7	43.7	47.6	14.3	19.
20	36.8	20.10	25.	29.2	33.4	37.6	41.8	45.10	50.	15.	20.
21	38.6	21.11	26.3	30.8	35.	39.5	43.9	48.2	52.6	15.9	21.
22	40.4	22.11	27.6	32.1	36.8	41.3	45.10	50.5	55.	16.6	22.
23	42.2	24.	28.9	33.7	38.4	43.2	47.11	52.9	57.6	17.3	23.
24	44.	25.	30.	35.	40.	45.	50.	55.	60.	18.	24.
25	45.10	26.1	31.3	36.6	41.8	46.11	52.1	57.4	62.6	18.9	25.
26	47.8	27.1	32.6	37.11	43.4	48.9	54.2	59.7	65.	19.6	26.
27	49.6	28.2	33.9	39.5	45.	50.8	56.3	61.11	67.6	20.3	27.
28	51.4	29.2	35.	40.10	46.8	52.6	58.4	64.2	70.	21.	28.
29	53.2	30.3	36.3	42.4	48.4	54.5	60.5	66.6	72.6	21.9	29.
30	55.	31.3	37.6	43.9	50.	56.3	62.6	68.9	75.	22.6	30.

Table continued.

Feet Long.	3 by 5	3 by 6	3 by 7	3 by 8	3 by 9	3 by 10	3 by 11	3 by 12	4 by 4	4 by 5	4 by 6
1	1.3	1.6	1.9	2.	2.3	2.6	2.9	3	1.4	1.8	2.
2	2.6	3.	3.6	4.	4.6	5.	5.6	6	2.8	3.4	4.
3	3.9	4.6	5.3	6.	6.9	7.6	8.3	9.	4.	5.	6.
4	5.	6.	7.	8.	9.	10.	11.	12.	5.4	6.8	8.
5	6.3	7.6	8.9	10.	11.3	12.6	13.9	15.	6.8	8.4	10.
6	7.6	9.	10.6	12.	13.6	15.	16.6	18.	8.	10.	12.
7	8.9	10.6	12.3	14.	15.9	17.6	19.3	21.	9.4	11.8	14.
8	10.	12.	14.	16.	18.	20.	22.	24.	10.8	13.4	16.
9	11.3	13.6	15.9	18.	20.3	22.6	24.9	27.	12.	15.	18.
10	12.6	16.	19.3	20.	22.6	25.	27.6	30.	13.4	16.8	20.
11	13.9	16.6	21.	22.	24.9	27.6	30.3	33.	14.8	18.4	22.
12	15.	18.	21.	24.	27.	30.	33.	36.	16.	20.	24.
13	16.3	19.6	22.9	26.	29.3	32.6	35.9	39.	17.4	21.8	26.
14	17.6	21.	24.6	28.	31.6	35.	38.6	42.	18.8	23.4	28.
15	18.9	22.6	26.3	30.	34.9	37.6	41.3	45.	20.	25.	30.
16	20.	24.	28.	32.	36.	40.	44.	48.	21.4	26.8	32.
17	21.3	25.6	29.9	34.	38.3	42.6	46.9	51.	22.8	28.4	34.
18	22.6	27.	31.6	36.	40.6	45.	49.6	54.	24.	30.	36.
19	23.9	28.6	33.3	38.	42.9	47.6	52.3	57.	25.4	31.8	38.
20	25.	30.	35.	40.	45.	50.	55.	60.	26.8	33.4	40.
21	26.3	31.6	36.9	42.	47.3	52.6	57.9	63.	28.	35.	42.
22	27.6	33.	38.6	44.	49.6	55.	60.6	66.	29.4	36.8	44.
23	28.9	34.6	40.3	46.	51.9	57.6	63.3	69.	30.8	38.4	46.
24	30.	36.	42.	48.	54.	60.	66.	72.	32.	40.	48.
25	31.3	37.6	43.9	50.	56.3	62.6	68.9	75.	33.4	41.8	50.
26	32.6	39.	45.6	52.	58.6	65.	71.6	78.	34.8	43.4	52.
27	33.9	40.6	47.3	54.	60.9	67.6	74.3	81.	36.	45.	54.
28	35.	42.	49.	56.	63.	70.	77.	84.	37.4	46.8	56.
29	36.3	43.6	50.9	58.	65.3	72.6	79.9	87.	38.8	48.4	58.
30	37.6	45.	52.6	60.	67.6	75.	82.6	90.	40.	50.	60.

Table continued.

Feet Long.	4 by 7	4 by 8	4 by 9	4 by 10	4 by 11	4 by 12	5 by 5	5 by 6	5 by 7	5 by 8	5 by 9
1	2.4	2.8	3.	3.4	3.6	4.	2.1	2.6	2.11	3.4	3.9
2	4.8	5.4	6	6.8	7.4	8.	4.2	5.	5.10	6.8	7.6
3	8.	8.	9.	10.	11.	12.	6.3	7.6	8.9	10.	11.3
4	9.4	10.8	12.	13.4	14.8	16.	8.4	10.	11.8	13.4	15.
5	11.8	13.4	15.	16.8	18.4	20.	10.5	12.6	14.7	16.8	18.9
6	14.	16.	18.	20.	22.	24.	12.6	15.	17.6	20.	22.6
7	16.4	18.8	21.	23.4	25.8	28.	14.7	17.6	20.5	23.4	26.3
8	18.8	21.4	24.	26.8	29.4	32.	16.8	20.	23.4	26.8	30.
9	21.	24.	27.	30.	33.	36.	18.9	22.6	26.3	30.	33.9
10	23.4	26.8	30.	33.4	36.8	40.	20.10	25.	29.2	33.4	37.6
11	25.8	29.4	33.	36.8	40.4	44.	22.11	27.6	32.1	36.8	41.3
12	28.	32.	36.	40.	44.	48.	25.	30.	35.	40.	45.
13	30.4	34.8	39.	43.4	47.8	52.	27.1	32.6	37.11	43.4	48.9
14	32.8	37.4	42.	46.8	51.4	56.	29.2	35.	40.10	46.8	52.6
15	35.	40.	45.	50.	55.	60.	31.3	37.6	43.9	50.	56.3
16	37.4	42.8	48.	53.4	58.8	64.	33.4	40.	46.8	53.4	60.
17	39.8	45.4	51.	56.8	62.4	68.	35.5	42.6	49.7	56.8	63.9
18	42.	48.	54.	60.	66.	72.	37.6	45.	52.6	60.	67.6
19	44.4	50.8	57.	63.4	69.8	76.	39.7	47.6	55.5	63.4	71.3
20	46.8	53.4	60.	66.8	73.4	80.	41.8	50.	58.4	66.8	75.
21	49.	56.8	63.	70.	77.	84.	43.9	52.6	61.3	70.	78.9
22	51.4	58.8	66.	73.4	80.8	88.	45.10	55.	64.2	73.4	82.6
23	53.8	61.4	69.	76.8	84.4	92.	47.11	57.6	67.1	76.8	86.3
24	56.	64.	72.	80.	89.	96.	50.	60.	70.	80.	90.
25	58.4	66.8	75.	83.4	91.8	100.	52.1	62.6	72.11	83.4	93.9
26	60.8	69.4	78.	86.8	95.4	104.	54.2	65.	75.10	86.8	97.6
27	63.	72.	81.	90.	99.	108.	56.3	67.6	78.9	90.	101.3
28	65.4	74.8	84.	93.4	102.8	112.	58.4	70.	81.8	93.4	105.
29	67.8	77.4	87.	96.8	106.4	116.	60.5	72.6	84.7	96.8	108.9
30	70.	80.	91.	100.	110.0	120.	62.6	75.	87.6	100.	112.5

Table continued.

Feet Long.	5 by 10	6 by 6	6 by 7	6 by 8	7 by 7	7 by 8	7 by 9	8 by 8	8 by 9	8 by 10	9 by 9
1	4.2	3.	3.6	4.	4.1	4.8	5.3	5.4	6.	6.8	6.9
2	8.4	6.	7.	8.	8.2	9.4	10.6	10.8	12.	13.4	13.6
3	12.6	9.	10.6	12.	12.3	14.	15.9	16.	18.	20.	20.3
4	16.8	12.	14.	16.	16.4	18.8	21.	21.4	24.	26.8	27.
5	20.10	15.	17.6	20.	20.5	23.4	26.3	26.8	30.	33.4	33.9
6	25.	18.	21.	24.	24.6	28.	31.6	32.	36.	40.	40.6
7	29.2	21.	24.6	28.	28.7	32.8	36.9	37.4	42.	46.8	47.3
8	33.4	24.	28.	32.	32.8	37.4	42.	42.8	48.	53.4	54.
9	37.6	27.	31.6	36.	36.9	42.	47.3	48.	54.	60.	60.9
10	41.8	30.	35.	40.	40.10	45.8	52.6	53.4	60.	66.8	67.6
11	45.10	33.	38.6	44.	44.11	51.4	57.9	58.8	66.	73.4	74.3
12	50.	36.	42.	48.	49.	56.	63.	64.	72.	80.	81.
13	54.2	39.	45.6	52.	53.1	61.8	68.3	69.4	78.	85.8	87.9
14	58.4	42.	49.	56.	57.2	65.4	73.6	74.8	84.	93.4	94.6
15	62.6	45.	52.6	60.	61.3	70.	78.9	80.	90.	100.	101.3
16	66.8	48.	56.	64.	65.4	74.8	84.	85.4	96.	106.8	108.
17	70.10	51.	59.6	68.	69.5	79.4	89.3	90.8	102.	113.4	114.9
18	75.	54.	63.6	72.	73.6	84.	94.6	96.	108.	120.	121.6
19	79.2	57.	66.6	76.	77.7	88.8	99.9	101.4	114.	126.8	128.3
20	83.4	60.	70.	80.	81.8	93.4	105.	106.8	126.	133.4	135.
21	87.6	63.	73.6	84.	85.9	98.	110.3	112.	132.	140.	141.9
22	91.8	66.	77.	88.	89.10	102.8	115.6	117.4	138.	146.8	148.6
23	95.10	69.	80.6	92.	93.11	107.4	120.9	122.8	141.	153.4	155.3
24	100.	72.	84.	96.	98.	112.	126.	128.	144.	160.	162.
25	104.2	75.	87.6	100.	102.1	116.8	131.3	133.4	150.	168.8	169.9
26	108.4	78.	91.	104.	106.2	121.4	136.6	138.8	156.	173.4	175.6
27	112.6	81.	94.6	108.	110.3	126.	141.9	144.	162.	180.	182.3
28	116.8	84.	98.	112.	114.4	130.8	147.	149.4	168.	186.8	189.
29	120.10	87.	101.6	116.	118.5	135.4	152.3	154.8	174.	193.4	195.9
30	125.	90.	105.	120.	122.6	141.	157.6	160.	180.	200.	202.6

Table continued.

Feet Long.	9 by 10	9 by 11	10 by 10	10 by 11	10 by 12	11 by 11	11 by 12
1	7.6	8.3	8.4	9.2	10.	10.1	11.
2	15	16.6	16.8	18.4	20.	20.2	22.
3	22.6	24.9	25.	27.6	30.	30.3	33.
4	30.	33.	33.4	36.8	40.	40.4	44.
5	37.6	41.3	41.8	45.10	50.	50.5	55.
6	45.	49.6	50.	55.	60.	60.6	66.
7	52.6	57.9	58.4	64.2	70.	70.7	77.
8	60.	66.	66.8	73.4	80.	80.8	88.
9	67.6	74.3	75.	82.6	90.	90.9	99.
10	75.	82.6	83.4	91.8	100.	100.10	110.
11	82.6	90.9	91.8	100.10	110.	110.11	121.
12	90.	99.	100.	110.	120.	121.	132.
13	97.6	107.3	108.4	119.2	130.	131.1	143.
14	105.	115.6	116.8	128.4	140.	141.2	154.
15	112.6	123.9	125.	137.6	150.	151.3	165.
16	120.	132.	133.4	146.8	160.	161.4	176.
17	127.6	140.3	141.8	155.10	170.	171.5	187.
18	135.	148.6	150.	165.	180.	181.6	198.
19	142.6	156.9	158.4	174.2	190.	191.7	209.
20	150.	165.	166.8	183.4	200.	201.8	220.
21	157.6	173.3	175.	192.6	210.	211.9	231.
22	165.	181.6	183.4	201.8	220.	221.10	242.
23	172.6	189.9	191.8	210.10	230.	231.11	253.
24	180.	198.	200.	220.	240.	242.	264.
25	187.6	206.3	208.4	229.2	250.	252.1	275.
26	195.	214.6	216.8	238.4	260.	262.2	286.
27	202.6	222.9	225.	247.6	270.	272.3	297.
28	210.	231.	233.4	256.8	280.	282.4	308.
29	217.6	239.3	241.8	265.10	290.	292.5	319.
30	225.	247.6	250.	275.	300.	302.6	330.

EXPLANATION.—Find the length in feet in the left hand column, and the dimensions of the sides in inches at the head of the other column; and underneath the latter and opposite the length will be found the contents in feet and inches board measure.

CASK-GAUGING.



Casks are usually comprised under the following figures, viz. :

1. The middle frustum of a spheroid.
2. The middle frustum of a parabolic spindle.
3. The two equal frustums of a paraboloid.
4. The two equal frustums of a cone.

Their contents can be computed by the rules for ascertaining the contents of these figures.

But in almost all ordinary casks the *bilge* or *swell* from

the *bung* to the head (not from head to head) is so small, that they are, with scarcely an appreciable difference in the results, usually regarded as the two equal frustums of a cone, and are very accurately gauged by three dimensions, as follows:

To find the contents of a cask by three dimensions.

RULE.—Add the bung and head diameters in inches, and divide them by 2 for the mean diameter; find the area of the mean diameter in the table of the areas of circles on page 298 and multiply it by the length of the cask in inches; then divide the product by 231 (the cubic inches in a gallon), and the quotient will be the number of gallons the cask contains.

EXAMPLE.—What are the contents in gallons of a cask, the bung diameter of which is 22 inches, the head diameter 20 inches, and the length 32 inches?

SOLUTION.— $22 + 20 = 42 \div 2 = 21$, mean diameter: then 346.36 , area of mean diameter, $\times 32 = 11083.52 \div 231 = 47.98$ gallons. *Ans.*

When the cask is much bilged or rounded from the bung to the head, a more accurate way is to gauge by four dimensions, as follows:

To find the contents of a cask by four dimensions.

RULE.—Add the head and bung diameters in inches, and the diameter taken in inches in the middle between the bung and head, and divide their sum by 3 for the mean diameter; find the area of the mean diameter in the table

of the areas of circles on page 298 and multiply it by the length of the cask in inches and divide the product by 231 (the cubic inches in a gallon), and the quotient will be the contents of the cask in gallons.

EXAMPLE.—What are the contents in gallons of a cask, the bung diameter of which is 24 inches, the middle diameter 20 inches, the head diameter 16 inches, and its length 40 inches?

SOLUTION.— $24 + 20 + 16 = 60 \div 3 = 20$, mean diameter : then 314.16 , area of mean diameter, $\times 40$ inches, length = $12566.40 \div 231 = 54.4$ galls. *Ans.*

CAPACITY OF BOXES.

A box 24 inches by 16 inches square, and 28 inches deep, will contain a barrel (5 bushels).

A box 24 inches square and 14 inches deep, will contain half a barrel.

A box 26 inches by 15.2 inches square, and 8 inches deep, will contain one bushel.

A box 12 inches by 11.2 inches square, and 3 inches deep, will contain half a bushel.


A box 8 inches by 8.4 inches square, and 8 inches deep, will contain one peck.

A box 8 inches by 8 inches square, and 4.2 inches deep, will contain one gallon.

A box 7 inches by 4 inches square, and 4.8 inches deep, will contain half a gallon.

A box 4 inches by 4 inches square, and 4.1 inches deep, will contain one quart.

CAPACITY OF WAGON-BEDS.



Wagon-Beds.

In most of the Eastern and many of the Western cities all market-men and traders, who make use of their wagon-beds as measures, are required to have them gauged and their capacity stamped on them by an officer appointed for that purpose. The wagon-makers in the country should stamp the contents in bushels on each bed they make before it leaves the shop. Should it be neglected, the following rule will enable every farmer to measure the contents in bushels of his wagon-bed for himself:

To find the contents of wagon-beds.

RULE.—If the opposite sides are parallel, multiply the length inside in inches, by the breadth inside in inches, and that again by the depth inside in inches, and divide

the product by 2150.42 (the number of cubic inches in a bushel), and the quotient will be the capacity in bushels.

EXAMPLE.—What is the capacity of a wagon-bed 10 feet long, 4 feet wide, and 15 inches deep?

SOLUTION.—120 inches, length, \times 48 inches, width, \times 15 inches, depth, $= 86400 \div 2150.42 = 40$ bushels. *Ans.*


RULE 2.—Should the head and tail boards, or either of them, be set in bevelling, add the top and bottom lengths together and divide by 2 for the mean length, and proceed by the foregoing rule. Should the sides be sloping, add the top and bottom widths, and divide by 2 for the mean width, and proceed by the foregoing rule.

Should the contents be required in cubic feet, divide the product by 1728 (the number of cubic inches in a cubic foot), instead of 2154.42, and the quotient will be the contents in cubic feet.



2150.42

FALSE BALANCES.



To detect false balances, scales, &c.

RULE.—After weighing the article transpose the weight and the article weighed, and if the latter is too light the weight will preponderate; if too heavy the article will preponderate.

To find the true weight.

RULE.—After transposing them as above, find the additional weight that will produce an equilibrium: weigh it with the article by the same balances: multiply the two false weights thus found, together, and the square root of the product will be the true weight.


EXAMPLE.—An article weighs 7 lbs. by a false balance: transposed it is found *too light*, and requires an additional weight to produce a counterpoise: this additional weight is found by the same balances to have a false weight of $9\frac{1}{4}$ lbs. What is the true weight of the article?

SOLUTION.— $9\frac{1}{4} \times 7 = 64$, the square root of which is 8 lbs., the weight. *Ans.*

EXAMPLE 2.—An article weighs 7 lbs.: transposed it is found *too heavy*, weighing only $5\frac{1}{4}$ lbs. by the same scales. What is the true weight?

SOLUTION.— $7 \times 5\frac{1}{4} = 36$, the square root of which is 6 lbs., the true weight. *Ans.*

NOTE.—In the 1st example the additional weight is added to the article to produce the equilibrium: in the second example the deficiency is taken from the weight to produce the counterpoise.



CISTERNS.


To find the number of gallons in square or oblong square cisterns.

RULE.—Multiply the length in inches by the width in inches, and that by the depth in inches, and divide the product by 231. The quotient will be the number of gallons.

EXAMPLE.—Given, a cistern 6 feet long by 3 feet wide and 4 feet deep; how many gallons will it contain?

SOLUTION.—72 inches, length, \times 36 inches, width, \times 48 inches, depth, $= 124416 \div 231 = 538.59$ galls. *Ans.*

To find the number of gallons in triangular cisterns.



RULE.—Multiply the base $a\ b$ in inches, by the perpendicular height $c\ d$ in inches, and half that sum by the depth in inches, and divide the product by 231. The quotient will be the number of gallons.

EXAMPLE.—Given, a triangular cistern 8 feet at the base or longest side, 7 feet in perpendicular height, 4 feet deep. How many gallons will it contain?

SOLUTION.—96 inches, base, $\times 84$, perpendicular height in inches, $\div 2 = 4032 \times 48$, depth in inches, $= 112896 \div 231 = 488.72$ galls. *Ans.*

To find the number of gallons in circular cisterns.

RULE.—Find the area of the circle in square inches, in the table of the “Areas of Circles,” on page 298, or by the rule given on page 296. Then multiply the area by the depth in inches, and divide the product by 231. The quotient will be the number of gallons.

EXAMPLE.—Given, a cistern 8 feet in diameter by 5 feet deep. How many gallons will it contain?

SOLUTION.—Area, the diameter being 96 inches 7238.2
 Multiplied by 60 in., the depth, gives 43429.20
 Divided by 231, cubic in. in a gall., “ 1880. gall. *Ans.*

TABLE, showing the contents of circular cisterns from 1 foot to 25 feet in diameter, for each 10 inches in depth.

DIAMETER.	GALLONS.	DIAMETER.	GALLONS.
1	4.896	$7\frac{1}{2}$	271.072
$1\frac{1}{2}$	11.015	8	313.340
2	19.583	$8\frac{1}{2}$	353.735
$2\frac{1}{2}$	30.545	9	396.573
3	44.064	$9\frac{1}{2}$	441.861
$3\frac{1}{2}$	59.980	10	489.600
4	78.333	11	592.400
$4\frac{1}{2}$	99.116	12	705.
5	122.400	13	827.450
$5\frac{1}{2}$	148.546	14	959.613
6	176.253	15	1101.610
$6\frac{1}{2}$	206.855	20	1958.421
7	239.906	25	3059.934

To find the number of gallons in tub-shaped cisterns.

RULE.—Find the cubes of the top and bottom diameters in inches, by means of the table on page 303, divide the difference between those cubes by the difference of the diameters in inches, and multiply this quotient by .7854, and again by $\frac{1}{3}$ of the depth in inches, and divide the product by 231. The quotient will be the number of gallons.

EXAMPLE.—Given, a tub-shaped cistern of a top diameter of 10 feet, a bottom diameter of 8 feet, and 6 feet deep. How many gallons will it contain?

SOLUTION.—Cube of 120 inches, the top diameter,	1728000
“ 96 “ “ bottom “	884736
Difference between cubes of diameters,	843264
Divided by 24, difference of diameters, gives	.35136
Multiplied by .7854, gives	27595.8144
“ again by $24, \frac{1}{3}$ the depth in inches, gives	662299.5456
Divided by 231, cubic inches in a gallon, gives	2867.09
galls. <i>Ans.</i>	

RULE 2.—Add the top and bottom diameters in inches and divide by 2 for the mean diameter. Find the area in square inches of the mean diameter by means of the table on page 298 or by the rule given on page 296. Multiply the area by the depth in inches, and divide the product by 231, and the quotient will be the number of gallons.

EXAMPLE.—What are the contents in gallons of a cistern 8 feet diameter at the top, 6 feet at the bottom, and 4 feet deep?

SOLUTION.—96 inches + 72 inches = 168 ÷ 2 = 84 inches, mean diameter; then 5541.77, area of mean diameter, × 48 inches, depth, = 266004.96 ÷ 231 = 1151.53 gallons. *Ans.*

NOTE.—The quantity of water which falls upon most farm buildings is sufficient to afford an ample supply for the domestic animals of the farm, when other supplies fail, were cisterns large enough to hold it provided. The average amount of rain that falls in the latitude of the Northern States during the year, is about 3 feet per year, or 3 inches per month. Every inch in depth that falls upon a roof yields 2 barrels for each ten feet square, and 72 barrels a year are yielded by 3 feet of rain. A barn 30 by 40 feet supplies annually from its roof 864 barrels, which is more than 2 barrels per day, the year round.

The size of cisterns should vary according to their intended use. If they are to furnish a *daily* supply of water, they need not be so large as for saving supplies against summer and droughts.

The size of the cistern in *daily* use need not exceed that of a body of water on the whole roof of the building, 7 inches deep, or two months' greatest fall of rain. Cisterns intended to save the water to draw from in time of drought, should be about three times as large.

To ascertain the size of cisterns adapted to roofs, &c.

RULE.—Multiply the length of the roof in inches by the breadth in inches, and that by the depth of the fall of rain required to be saved, and divide the product by 231, and

the quotient will be the number of gallons. Divide the number of gallons by $31\frac{1}{2}$, and it will give the number of barrels.

EXAMPLE.—What must be the capacity of a cistern to contain the water running from a roof 40 feet long by 30 wide, for 2 months: estimated fall of rain 7 inches?


SOLUTION.—480 inches, length, \times 360 inches, width, \times 7 inches, depth of rain, $=1909600 \div 231 = 8266\frac{2}{3}$ galls. *Ans.*

NOTE.—To ascertain the necessary dimensions of a cistern large enough to contain $8266\frac{2}{3}$ gallons, consult the foregoing table. It will there be found that a cistern 13 feet in diameter contains 827 gallons for each 10 inches in depth. To give the cistern 10 times that depth, or 100 inches ($8\frac{1}{3}$ feet) will make it contain 8270 gallons. Hence a cistern 13 feet in diameter, and $8\frac{1}{3}$ feet deep, will be large enough.

To further aid the inquirer in ascertaining the requisite diameters of cisterns for the above purposes, we subjoin an additional

TABLE, *showing the contents of circular cisterns in barrels for each foot in depth.*

5 feet	4.66
6 "	6.74
7 "	9.13
8 "	11.93
9 "	15.10
10 "	18.65




The above cut represents the sectional view of a filtered cistern, with a brick wall partition in the middle and the box of charcoal and sand at the bottom, with alternate layers of each. The pipe at the left leads from the roof, and the one at the right connects with the pump. With this style of cistern properly constructed, no one need be in want of pure wholesome water.

To construct a filtering cistern to furnish pure water for domestic use.

RULE.—Divide the cistern into two equal compartments by a wall of brick or stone, open at the bottom to the height of about six inches, and water-tight thence to the top. Let one compartment be for receiving the water, and the other for containing it when filtered and ready for use. Put alternate layers, 6 inches deep, of gravel, sand,

and pounded charcoal at the bottom of the former, and sand and gravel at the bottom of the latter. The former will receive the water from the pipe, and it will rise filtered in the latter.

ANOTHER MODE.—Divide the cistern as above by a double open wall of stone or brick, with an interspace of about six inches between the walls. Fill the interspace with sand and pounded charcoal. Let one compartment receive the water, and it will pass through the filter into the other ready for use.



HYDRAULICS.

The science of *hydraulics* treats of the motion of non-elastic fluids; *hydrodynamics*, of the force of that motion; and *hydrostatics*, of the pressure, weight, and equilibrium.

THE FUNDAMENTAL LAWS OF HYDRAULICS, &c.

1. Descending water is governed by the same laws as falling bodies.

2. Water will fall 1 foot in $\frac{1}{4}$ of a second, 4 feet in $\frac{1}{2}$ a second, and 9 feet in $\frac{3}{4}$ of a second, and so on in the same ratio.

3. The velocity of a fluid propelled through an orifice by a head of water in a cistern or reservoir, is the same that a body would acquire by falling perpendicularly through a space equal to that between the top of the head and the centre of the opening, *less the friction which, in pipes, drains, and sluices, increases as the square of the velocity.*

4. The mean velocity of water propelled through an opening by a head of 1 foot is $5\frac{1}{2}$ feet per second.

5. Fluids press equally in all directions.

6. The pressure of a fluid on the bottom of a vessel is as

the base and perpendicular height, whatever may be the figure of the vessel.

7. The pressure of a fluid on any kind of surface, horizontal, vertical, or oblique, is equal to the weight of the column of the fluid, the base of which is equal to the area of the surface pressed, and the height of which is equal to the distance from the surface of the fluid to its centre of gravity, on the surface pressed.

8. The side of a vessel filled with water sustains a pressure equal to the area of the side multiplied by half the depth, whether the sides be vertical, oblique, or horizontal.

9. If the vessel be tub-shaped, or in the form of an inverted frustum of a cone or pyramid, the bottom sustains a pressure equal to the area of the bottom and the depth of the fluid.

10. The quantity of water that will flow out of a perpendicular slit or aperture from the surface of the head to its base, is but two-thirds of what would flow out of a slit of the same dimensions were it horizontal at the level of the base.

11. A *circular* pipe of the same area as a *square*, *triangular*, or *irregular* one, will discharge *more* water in a given time.

12. The greater the length of the discharging pipe, the less the discharge, unless the pipe be perpendicular.

13. A pipe that is inclined will discharge more water in a given time than a horizontal pipe of the same dimensions.

14. The friction of a fluid is greater in small than in large pipes, when equal quantities are discharged.

15. In perpendicular pipes, the discharge being governed by the law of gravitation, the greater the length of the pipe, the greater the discharge.

16. When a prismatic vessel empties itself through an aperture, twice the quantity would be discharged in the same time if it were kept full.

17. In a stream, sluice, or ditch, the velocity is the greatest at the surface and in the middle of the current.

18. The *time* occupied by a given quantity of water passing through pipes or sewers of equal apertures and lengths, and with equal falls, is in the following proportions, viz.: In a straight line, as 90; in a regular curve, as 100; and in passing a right angle, as 140.

To find the velocity of a stream issuing from a head of water.

RULE.—Multiply the height of the head in feet by 64.33, and the square root of the product will be the velocity in feet per second.

EXAMPLE.—What is the velocity of a stream projected through an opening by a head of 12 feet?

SOLUTION.— $12 \times 64.33 = 771.96$, the square root of which is 27.780 feet per second. *Ans.*

To find the head, the velocity being given.

RULE.—Square the velocity and divide it by 64.33, and the quotient will be the head in feet.

EXAMPLE.—What is the head of water that projects a stream 27.780 feet per second?

SOLUTION.— $27.780^2 = 771.96 \div 64.33 = 12$ feet. *Ans.*

NOTE.—In the above results no allowance is made for friction, which should be made in order to ascertain the practical results. The friction of water passing out of orifices, and not through pipes, sluices, or sewers, is, however, very small.

To find the quantity of water that will issue from an opening, the dimensions of the opening and the head being given.

RULE.—Find the velocity of the jet or stream by the foregoing rule, and multiply it by the area of the orifice in feet, and the product will be the number of cubic feet per second the orifice will discharge.

EXAMPLE.—How much water will an orifice of an area of 2 square feet discharge per second under a head of 12 feet?

SOLUTION.— $12 \times 64.33 = 771.96$, the square root of which is 27.780 feet velocity; then, 27.780×2 feet, area, $= 55\frac{7}{8}$ cubic feet per second. *Ans.*



To find the velocity of currents in drains, ditches, sluices, brooks, or rivers.

RULE.—Find the velocity of the surface of the current in the middle of the stream by taking the number of inches a floating body passes over it in one second.

This, for all ordinary practical purposes, will be sufficient. But to find the *mean* or *average* velocity, take the square root of the velocity so found, double it, and deduct it from the velocity at the top, and add one to the remainder, and the result will be the velocity at the bottom. Add the top and bottom velocities, and divide them by two for the mean velocity.

EXAMPLE.—What is the mean velocity of a current, the velocity of which at the surface, in the middle of the stream, is 36 inches per second?

SOLUTION.— $\sqrt{36} = 6 \times 2 = 12 - 36 = 24 + 1 = 25$, velocity
5

at bottom; then, $36 + 25 = 61 \div 2 = 30.5$ inches per second, mean velocity. *Ans.*

To find the volume of water discharged by drains, sluices, brooks, &c., of given dimensions, in a given time.

RULE.—Multiply the velocity of the current per second in feet, by the area of the transverse section of the drain or sluice, in feet, and the product will be the quantity discharged per second, in cubic feet.

EXAMPLE.—What volume of water will a drain 2 feet wide and 3 feet deep discharge in one hour, the mean velocity of the current being 30 inches per second?

SOLUTION.— $2 \times 3 = 6$ sq. ft., area of section $\times 2\frac{1}{2}$ ft., velocity, $= 15$ cubic feet discharged per second; then, 15×3600 seconds (one hour) $= 54,000$ cubic feet per hour. *Ans.*

NOTE.—The standard gallon contains 231 cubic inches, and a cubic foot contains 1728 cubic inches. Accordingly, a cubic foot of water contains 7.476 standard gallons. Hence, if we multiply the number of cubic feet by 7.476, it will give the number of gallons. For instance, the drain in the above example discharges 54,000 cubic feet per hour, which, multiplied by 7.476, gives 403,704 gallons discharged per hour.

To find the velocity of water running through pipes.

RULE.—Multiply the height of the head in feet by 2500; divide this product by a divisor obtained as follows: Di-

vide 13.88 by the diameter of the pipe in inches, and multiply the quotient by the length of the pipe in feet, and the result will be the divisor aforesaid. Divide the first product by this sum, and the square root of the quotient will be the velocity in feet per second of the current in the pipe.

EXAMPLE.—What is the velocity of water in a pipe 5 inches diameter and 100 feet long, and under a head of 2 feet?

SOLUTION.— $13.88 \div 5 = 2.776 \times 100 = 277.6$ and $2500 \times 2 = 5000$; then, $5000 \div 277.6 = 18$, the sq. root of which is 4.24 feet. *Ans.*

To find the quantity of water discharged through pipes.

RULE.—Multiply the velocity of the current per second in feet by the area of the transverse section of the pipe in feet, and the product will be the quantity discharged in cubic feet per second.

EXAMPLE.—What quantity of water will a pipe 6 inches diameter and 100 feet long discharge per hour under a head of 2 feet?

SOLUTION.—By the preceding rule, find the velocity of the current in the pipe, thus: 2500×2 feet, head, $= 5000$, $13.88 \div 6$ inches, the diameter of the pipe, $= 2.313 \times 100$ feet, length of the pipe, $= 231.3$, divisor; $5000 \div 231.3 = 21.62$, the square root of which is 4.65 feet, velocity per second. Then, $4.65 \times .2063$ square feet, area of pipe, $=$

.942 cubic feet discharged per second. $.942 \times 3600$ seconds (one hour) = 3391 cubic ft. discharged per hour. *Ans.*

To find the pressure of a fluid on the bottom of a vessel, cistern, or reservoir

RULE.—Multiply the area of the base in square feet by the height of the fluid in feet, and their product by the weight of a cubic foot of the fluid.

EXAMPLE.—What is the pressure on the bottom of a cistern 10 feet in diameter and 8 feet deep, filled with water?

SOLUTION.— 78.54 , area of bottom, $\times 8 = 628.32 \times 62\frac{1}{2}$ lbs., the weight of a cubic foot of water, = 39.370 lbs. *Ans.*

To find the pressure on the side of a vessel.

RULE.—Multiply the area of the side in feet by half its depth in feet, and that by the lbs. per cubic foot of the fluid.

EXAMPLE.—What is the pressure upon the sloping side of a pond 10 feet square by 8 feet deep?

SOLUTION.— $10^2 = 100 \times 4$, half the depth, = $400 \times 62\frac{1}{2}$ lbs., the weight of a cubic foot of water, = 25000 lbs. *Ans.*

NOTE.—It is proper to remark that all of these rules, while they are theoretically correct, do not pretend to embrace a variety of circumstances which affect the flow of water through apertures, and which should be taken into consideration in all cases. These circumstances cannot be


measured by rules, and the just estimate of their influence must depend on experience.

1. Water will flow more rapidly from an aperture in a vessel if a funnel-shaped tun or a rapidly widening trough be attached to it on the outside. This prevents, so to speak, the intercrossing of the currents as they flow over the sides of the aperture; instead of obstructing itself; by reason of its tendency to cross the centre of the opening, the water follows the sides of the funnel or trough, and allows the full area of the opening to discharge freely.

2. The ease with which a given quantity of water can be made to pass through a pipe depends (other things being equal) upon the proportion between the area of the opening and its circumference—the latter being a source of friction. (See Nos. 14 and 11 above.)

3. The ease of the flow depends on the perfect *uniformity* of the channel. A lump or any other inequality in the side of a pipe will disturb the current and cause the water to obstruct itself. Perfect *form* is more important than a *smooth surface*.

4. The same principle operates in the case of deflections from a straight line. If the water is turned out of its course the evenness of the flow is disturbed, and it becomes more difficult (see No. 18 above). The influence of a "regular curve" is in proportion to its radius; more water will flow through a pipe which turns in a large circle than in one which turns more abruptly.



This cut is intended to illustrate the use of the Hydraulic Ram; representing one operated by the water, from a spring near which it is located, and forcing the water through suitable leading and discharge pipe, to a considerable elevation (either perpendicular or upon an inclined plane) to a trough, which may be placed in any convenient locality for watering farm stock of every description,—affording a constant supply of fresh water the year round. It may also be used to supply a cistern or a water-tank in the house.



THE HYDRAULIC RAM.

The *hydraulic ram* is a machine for forcing a portion of a brook or stream to any required elevation and distance, when the requisite head or pressure can be obtained.

Wherever a large spring or a limited but constant stream is at hand, by which a fall of four or five feet may be produced, by building a dam or otherwise, a portion of the water of such spring or stream may be raised to a perpendicular height of more than 100 feet by its own power, through the agency of the water-ram. Thus, a stream in a deep valley, or a rivulet or brook situated some distance below a point where it is desired to have a cistern or reservoir, may be made to raise a part of its water by one of these machines. From such a cistern or reservoir the water may afterwards be conveyed to any part of the premises below it, and applied for the purpose of irrigation, watering of stock, manufactories, or domestic or ornamental use.

The power of the ram, and the height to which it will raise the water, as also the quantity raised, are in proportion to the volume of the stream and the head or fall obtained.

The ram is applicable where no more than 18 inches fall can be obtained.

The *distance* which the water has to be conveyed, and the consequent length of pipe, have also a bearing upon the quantity raised and its elevation, as the larger the pipe through which the water has to be forced, the greater the friction to be overcome, and the more the power consumed in the operation.

The ram can be applied to convey water a distance of from 100 to 200 rods, and to elevations of from 100 to 200 feet.

A fall of 10 feet from the spring or brook to the ram is sufficient to force the water to any elevation not over 150 feet above the ram, and in distance not over 150 rods from it.

Although the same fall will raise water to a much greater elevation, and force it to a greater distance, yet the quantity will *diminish* as the height and distance are increased.

When a sufficient quantity of water is raised by an adequate fall the fall should not be increased, as by so doing the strain upon the ram is unnecessarily increased, and its durability lessened.

The *proportion* which the height to which the water is raised, and the quantity raised, bear to the fall and to the volume of the spring or stream, is about five times the height of the fall, and $\frac{1}{4}$ of the volume of the stream forced a distance of 50 rods—allowing for the friction in both the supply and discharging pipes.

Thus, if the ram be placed under a fall of 5 feet, for every 7 gallons drawn from the spring, 1 gallon may be raised 25 feet, or $\frac{1}{2}$ a gallon 50 feet, and forced a distance of 50 rods. If the fall be 10 feet, it will raise one gallon 50 feet, or $\frac{1}{2}$ a gallon 100 feet, for every 7 gallons discharged by the stream. If the fall be 10 feet, and the *volume of the stream be doubled*, it will raise 1 gallon 100 feet, and so on in the same ratio.

The pipe leading from the spring or head of the fall to the ram is called the *supply pipe*.

The pipe leading from the ram to the reservoir or cistern is called the *discharging pipe*.

The shorter and straighter the supply pipe, the better. Hence, unless the supply pipe is laid to the head of a spring, it is better to dam the stream at the head of its greatest fall, and after inserting the supply pipe at the base of the dam, let it follow the depression of the bed of the stream to the ram at the lowest point.

The shorter and straighter the discharging pipe the better; there is less friction to be overcome.

Should it be necessary to curve either pipe, let the *radius* of the curve be as large as possible.

To ascertain the quantity of water and the height to which it may be elevated by a given fall and volume of water—discharging pipe not over 50 rods.

RULE.—Find, by means of a common level, the fall of your spring or stream; then find the quantity of water it

discharges per minute or hour, by means of the rule given for that purpose on page 98; then multiply the height of the fall by 5, for the elevation, and divide the number of gallons discharged by the stream by 7, for the quantity of water raised.

EXAMPLE.—Given, a spring with a fall of 8 feet, discharging 28 gallons per minute. How high and how much water will it raise per minute by means of a ram—discharging pipe not exceeding 50 rods?

SOLUTION.— $8 \times 5 = 40$ feet elevation. $28 \div 7 = 4$ gals. per minute. *Ans.*

NOTE.—In the same ratio, it will raise 2 gallons 80 feet per minute, or 1 gallon 160 feet per minute, and so on.

The following working results of water rams now in actual use, will enable the inquirer to ascertain the elevating capacity of springs, with various falls and volume of water. The rams used are "Rumsey & Co.'s Premium Hydraulic Rams," Seneca Falls, N. Y.

1.—Fall from surface of water in spring to ram.....	4 feet.
Length of supply pipe, inside diameter 1 inch.....	60 "
Volume of water discharged by spring in 10 minutes.....	25 gallons.
Length of discharging pipe, inner diameter $\frac{3}{8}$ inch, curved in three places to a semicircle.....	190 feet.
Elevation of discharging pipe from ram to cistern.....	19 "
Discharged every ten minutes.....	$3\frac{1}{2}$ gallons.
2.—Fall from surface of water in spring to ram.....	10 feet.
Length of supply pipe, inside diameter $1\frac{1}{2}$ inches.....	40 "
Volume of water discharged by spring per minute.....	20 gallons.
Length of discharging pipe, $\frac{1}{2}$ inch inside diameter.....	50 rods.
Elevation of discharging pipe from ram to cistern.....	85 feet.
Discharged per minute.....	$2\frac{1}{2}$ gallons.

- 3.—Fall from surface of water in spring to ram..... 3½ feet.
 Length of supply pipe, inside diameter 1½ inches..... 30 "
 Volume of water discharged by spring..... not given.
 Length of discharging pipe inside diameter ½ inch..... 30 rods.
 Elevation of discharging pipe from ram to cistern..... 35 feet.
 Discharged a constant stream ½ inch diameter.
- 4.—Fall from surface of water in spring..... 12 feet.
 Length of supply pipe, inside diameter 1½ inches..... 32 "
 Volume of water discharged by spring..... not given.
 Length of discharging pipe, inside diameter ½ inch..... 14 rods.
 Elevation of discharging pipe from ram to cistern at barn.... 35 feet.
 Discharged a constant stream ½ inch diameter, at barn, affording more than a supply for 52 head of cattle.
- 5.—Fall from surface of water in spring to ram..... 9 feet.
 Length of supply pipe, inside diameter one inch..... 50 "
 Volume of water discharged by spring..... not given.
 Length of discharging pipe, inside diameter ½ inch..... 100 feet.
 Elevation of discharging pipe from ram to cistern..... 35 "
 Discharges a constant stream, ½ inch diameter, into a cistern at house and after supplying water for the domestic use of a large family, passes off to the cattle yard 20 rods further, affording an abundant supply for a large herd of cattle.
- 6.—Fall from surface of water in spring to ram..... 8 feet.
 Length of supply pipe, inside diameter 1½ inches..... not given.
 Volume of water discharged by spring..... "
 Length of discharging pipe, ½ inch inside diameter..... 70 rods.
 Elevation of discharging pipe from ram to cistern..... 80 feet.
 Delivers a good supply of running water at house and barn, sufficient for all necessary purposes.
- 7.—Fall of water from surface of spring to ram..... 10 feet.
 Length of supply pipe, inside diameter 1½ inches..... not given.
 Volume of water discharged by spring..... "
 Length of discharging pipe, ½ inch inside diameter. 76 rods.
 Elevation of discharging pipe from ram to cistern..... 110 feet.
 Delivers a constant stream of ½ inch diameter.
- 8.—Fall from surface of spring to ram..... 6½ feet.
 Length of supply pipe, inside diameter 1½ inches..... 60 rods.
 Elevation of discharging pipe from ram to cistern..... 60 feet.
 Discharges sufficient water in barn yard to supply 30 head of cattle.
- 9.—Fall from surface of spring to ram..... 9 feet.
 Size of supply pipe, inside diameter 2 inches, length..... not given.
 Length of discharging pipe, inside diameter ½ inch..... 150 rods.
 Elevation of discharging pipe from ram to cistern..... 130 feet.
 Delivers an abundant supply of water for house, barn, barn-yard and hog-pen.

10.—Fall from surface of dam to ram.....	7 feet.
Length and size of supply pipe.....	not given.
Volume of water discharged by stream.....	"
Length of discharging pipe, (size not given).....	126 rods.
Elevation of discharging pipe from ram to cistern.....	75 feet.
Discharges 25 barrels of water in 24 hours.	
11.—Fall from spring to ram.....	11 feet.
Size of supply pipe, 2 inches calibre; length.....	42 "
Length of discharging pipe, $\frac{1}{2}$ inch calibre.....	75 rods.
Elevation of discharging pipe from ram to cistern.....	98 feet.
Discharges over 30 barrels of water per day.	

NOTE.—The size, strength, and weight of the supply and discharging pipes must be in proportion to the head or pressure on them. They are proportioned and adjusted to the capacity of the ram by the manufacturer, and are generally sold with the machine.

When a very large supply of water is required for manufacturing or other purposes, and a stream of sufficient volume and fall is obtained, it is better to set two or three rams of a smaller size, all playing into one discharging pipe, than to set one large ram. If one ram becomes disabled, the others supply the demand.

Should the fall and volume of one stream or spring not supply enough water, and at the required elevation, and there be other springs near by, set a ram in each, all meeting in one discharging pipe. Their combined power will increase the elevation and the quantity raised.

The pipes can be so laid, and the ram so set, as to protect them from the frost during the winter.


The fall of one spring or stream may be used to raise the water of another and better spring or stream, whose own fall is not sufficient.

Mr. H. L. Emery, of Albany, in a communication to the *Country Gentleman*, says: "The result of a water ram is calculated upon the principle that a pound of force will raise a pound of water an equal height, and a less quantity to a greater height, which height is limited only by the strength of the pipes themselves.

"To enable any one to select the size ram it is necessary to compute the elevation to be overcome, and the greatest amount of fall which can be conveniently obtained, and divide the first by the last, and the quotient will be the proportion of the water (passing through the drive-pipe) which will be raised; first, however, deducting for waste of power and friction say $\frac{1}{4}$ of the amount; thus, with ten feet fall and one hundred feet elevation, one-tenth of the water would be raised, if there were no friction or loss; but deducting, say one-quarter for loss, and $7\frac{1}{2}$ gallons for each 100 gallons would be raised, all the balance of the water being required or wasted to accomplish this result."



THE HYDRAULIC PRESS.



The Hydraulic or Hydrostatic Press is a machine by which a small force may be made to exert a great pressure. Its construction may be understood by the above cut. Two metallic cylinders, A and B, of different sizes, are joined together by a tube K. In the small cylinder there is a piston *p* which can be moved up and down by the handle

M. In the large cylinder there is also a piston P, having at its upper end a large iron plate, which moves freely up and down in a strong frame-work Q. Between the iron plate and the top of this framework the body to be pressed is placed. Now, when the small piston is raised, the cylinder A is filled with water drawn from the reservoir H, below, and when it is pushed down this water is forced into the large cylinder through the pipe K. There is a valve in this tube which prevents the water from returning, so that each stroke of the small piston pushes an additional quantity of water into the large cylinder. By this means the large piston is pushed up against the body to be pressed. To calculate the pressure exerted by the large piston we must remember that the force acting upon the piston in A, will be exerted upon every equal amount of surface in B. To illustrate this: suppose the area of the large piston to be 10 times the area of the small one; then one pound at A will produce a pressure of ten pounds at P. The handle M increases the advantage still more, according to the principle of the lever to be explained in a future chapter. By increasing the size of the large cylinder, and diminishing the size of the small one, the pressure exerted by a given power will be increased proportionately. The weight of a man's hand might thus be made to lift a ship with all its cargo. The only limit to the increase of power would be the strength of the material of which the machine is made.

WEIGHT OF LEAD PIPE.

TABLE, showing the weight of lead pipe per yard, from $\frac{1}{4}$ to $4\frac{1}{2}$ inches diameter.

Diameter.	Weight in lbs. oz.	Diameter.	Weight in lbs. oz.
$\frac{1}{4}$ inch medium.....	3	$1\frac{1}{2}$ inch extra light.....	9
" " strong.....	4	" " light.....	13
$\frac{1}{2}$ " light.....	3	" " medium.....	15 8
" " medium.....	4	" " strong.....	19
" " strong.....	5	$1\frac{3}{4}$ " medium.....	16
" " extra strong.....	6 6	" " strong.....	20
$\frac{5}{8}$ " light.....	5	2 " light.....	16 12
" " medium.....	6 6	" " medium.....	20
" " strong.....	7 8	" " strong.....	23
" " extra strong.....	8 4	$2\frac{1}{2}$ " light.....	25
$\frac{3}{4}$ " " light.....	5	" " medium.....	30
" " light.....	6 4	" " strong.....	35
" " medium.....	8	3 " light.....	30
" " strong.....	9 12	" " medium.....	35
" " extra strong.....	10 8	" " strong.....	44
1 " " light.....	6 14	$3\frac{1}{2}$ " medium.....	45
" " light.....	8 5	" " strong.....	54
" " medium.....	10 5	" " extra strong.....	70
" " strong.....	12 4	4 " waste, light.....	15 14
$1\frac{1}{2}$ " extra light.....	8 5	" " medium.....	21
" " light.....	9 12	" " strong.....	26
" " medium.....	11	$4\frac{1}{2}$ " light.....	17 4
" " strong.....	12 8	" " medium.....	24
" " extra strong.....	14 10	" " strong.....	29

Very light pipe.

$\frac{1}{4}$ inch.....	1 lb.	$\frac{3}{4}$ inch.....	3 lbs. 6 oz.
".....	$1\frac{1}{2}$ "	1 ".....	5 " 10 "
$\frac{1}{2}$ ".....	2 "	$1\frac{1}{4}$ ".....	6 " 14 "
".....	$2\frac{1}{2}$ "		

NOTE.—Should the pipe be sold by the pound, multiply the price per lb. by the weight per yard in the above table, and it will give the price per yard.

FUEL.

The following table, abridged from Browne's *Sylva Americana*, will be found valuable to housekeepers in aiding them to form an estimate of the comparative value of fire woods in a *seasoned state*, or when burnt to charcoal.

TABLE, showing the Comparative Values of Fire Woods.


WOODS.	Specific Gravity of dry Woods.	Pounds dry Wood in one cord.	Specific Gravity of dry Coals.	Pounds Charcoal in one bushel.	Pounds Charcoal from one cord of dry Wood.	Bushels Charcoal from one cord of dry Wood.	Value as compared with ash-bark Hickory.
Shellbark Hickory,	1.000	4469	.625	82.89	1172	36	100
Common Walnut,919	4241	.637	83.52	1070	32	95
White Oak,855	3821	.401	21.10	826	39	81
Thick Shellbark Hickory, ..	.829	3705	.509	26.78	848	32	81
White Ash,722	3450	.547	28.78	884	31	77
Scrub Oak,747	3339	.392	20.63	774	38	73
Witch Hazel,784	3505	.368	19.36	750	39	72
Apple Tree,697	3115	.445	22.41	779	33	70
Red Oak,728	3255	.400	21.05	630	39	60
Black Gum,703	3142	.400	21.05	696	33	67
Black Walnut,681	3044	.418	22.00	677	31	65
White Beech,724	3233	.518	27.26	635	23	65
Black Birch,697	3115	.428	22.52	604	27	63
Yellow Oak,653	2919	.295	15.52	631	41	60
Sugar Maple,644	2878	.431	22.68	617	27	60
Sassafras,618	2762	.427	22.47	624	28	59
White Elm,589	2592	.357	18.79	614	34	58
Holly,602	2691	.374	19.68	613	31	57
Wild Cherry,597	2668	.411	21.63	579	27	55
Yellow Pine,551	2463	.333	17.52	585	33	54
Sycamore, or Buttonwood, ..	.535	2391	.371	19.68	564	29	52
Chestnut,522	2333	.379	19.94	590	30	52
Spanish Oak,548	2449	.362	19.05	562	30	52
Poplar,563	2516	.383	20.15	549	27	52
Butternut,567	2534	.237	12.47	527	42	51
White Birch,530	2369	.364	19.15	450	21	48
Jersey Pine,478	2137	.385	20.26	532	26	48
Pitch Pine,426	1904	.298	15.68	510	33	43
White Pine,418	1868	.293	15.42	455	39	42
Lombardy Poplar,397	1774	.245	12.89	444	34	40

NOTE.—It will be remarked that shellbark hickory is made the *standard* in the above table, not only of the fuel but also of the specific gravity, the value and specific gravity of the other woods being determined by the proportion they severally bear to this standard. The table has a further use, namely, to determine the price that should be paid per cord for other woods, taking the price paid for shellbark hickory as the standard. For instance, should shellbark be selling for \$6.00 per cord, white oak is worth \$4.86; for, as 100, the value of shellbark, : \$6.00, its price, :: 81, the value of white oak, : \$4.86, its price; and other kinds in the same proportion.

A cord of wood is 128 cubic feet; the sticks or billets are cut 4 feet long and piled 4 feet high and 4 feet wide; 8 feet in length making a cord.

The wood-cutter has a measure of two feet marked on his axe handle with which he measures the length of each stick, making due allowance for the carf, or the bevel of the cut. All fuel should, however, be sold by weight.

When the weights of different woods are equal, that which contains the most hydrogen will, during combustion, give out the greatest amount of heat. Hence, pine is preferable to oak, and bituminous to anthracite coal. When wood is used as fuel it should be thoroughly dried, as in its green and ordinary state it contains 25 per cent. of water; the heat to evaporate which is necessarily lost. To kiln-dry it adds 12 per cent. to its value over seasoned wood.



Coal Mining in Pennsylvania.

TABLE, showing the weights per cubic foot of the different kinds of Coal.

Designation.	Weight in lbs.	Designation.	Weight in lbs.
Anthracite,	50 to 55	Western, (bitum.)	47
Bituminous,	45 to 55	English, "	50
Cumberland,	53	Charcoal, (hard wood)	18½
Virginia, (bitum.)	49	do. (soft or pine wood) ..	18

NOTE.—Soft coals are usually purchased at the rate of 28 bushels of 5 pecks each, to a ton of 43.56 cubic feet. Anthracite, 20 bushels to the ton

To prepare charcoal.

Charcoal is prepared by clearing off the top soil from a circular space of the required dimensions, and piling bil-

lets of wood in it into a pyramidal heap, with several spiracles or flues formed through the pile. Chips and brushwood are put into those below, and the whole is so constructed as to kindle through in a very short time. It must then be covered all over with clay or earth beaten close, leaving openings at all the spiracles or flues. The pile is then ignited, and carefully watched and kept from bursting into a flame, by instantly closing the flues should such happen. Whenever the white watery smoke issuing from the flues is observed to be succeeded by a thin, blue, and transparent smoke, the holes must be immediately stopped; this being the indication that all the watery vapor is gone, and the burning of the true coaly matter commencing. Thus a strong red heat is raised throughout the whole mass, and all the volatile matters are dissipated by it, and nothing now remains but the charcoal. The holes being all stopped in succession as this change of the smoke is observed, the fire goes out for want of air. The pile is now allowed to cool, which requires many days, for charcoal being a very bad conductor of heat, the pile long remains red hot in the centre, and if opened in this state would instantly burn with great fury. Even when it is opened, the heat retained by some of the larger pieces often ignites it, to guard against which water should be provided to instantly extinguish it when observed.

PROPERTIES OF CHARCOAL.

Although charcoal is so combustible, it is, in some re-

spects, a very unchangeable substance, resisting the action of a great variety of other substances upon it. Hence posts are often charred before being put into the ground. Grain has been found in the excavations at Herculaneum, which was charred at the time of the destruction of that city, eighteen hundred years ago, and yet the shape is perfectly preserved, so that you can distinguish between the different kinds of grain. While charcoal is itself so unchangeable, it preserves other substances from change. Hence meat and vegetables are packed in charcoal for long voyages, and the water is kept in casks which are charred on the inside. Tainted meat can be made sweet by being covered with it. Foul and stagnant water can be deprived of its bad taste by being filtered through it. Charcoal is a great decolorizer. Ale and porter filtered through it are deprived of their color, and sugar-refiners decolorize their brown syrups by means of charcoal, and thus make white sugar. Animal charcoal, or bone-black, is the best for such purposes, although only one-tenth of it is really charcoal, the other nine-tenths being the mineral portion of the bone.

Charcoal will absorb, of some gases, from eighty to ninety times its own bulk. As every point of its surface is a point of attraction, it is supposed to account for the enormous accumulation of gases in the spaces of the charcoal. But this accounts for it only in part. There must be some peculiar power in the charcoal to change, in some way, the condition of a gas of which it absorbs ninety times its own bulk.—*Hooker.*

NOTES.—The best quality of charcoal is made from oak, maple, beech, and chestnut.

Wood will furnish, when properly charred, about 20 per cent. of coal.

A bushel of coal from hard wood weighs 30 lbs.

A bushel of coal from pine weighs 29 lbs.

TABLE, *showing the number of parts of charcoal afforded by 100 parts of different kinds of wood.*

Woods.		Parts charcoal.	Color.
Lignum Vitæ	afforded.....	26.8	Grayish.
Mahogany	"	25.4	Brown.
Laburnum	"	24.5	Velvet black.
Chestnut	"	23.2	Glossy black.
Oak	"	22.6	Black.
Black beech	"	21.4	Fine black.
Holly	"	19.9	Dull black.
Sycamore	"	19.7	Fine black.
Walnut	"	20.6	Dull black.
Beech	"	19.9	Dull black.
Maple	"	19.9	Dull black.
Norway Pine	"	19.2	Shining black.
Elm	"	19.2	Fine black.
Sallow	"	18.4	Velvet black.
Ash	"	17.9	Shining black.
Birch	"	17.4	Velvet black.
Scottish Pine	"	16.4	Brownish.

COKE.

Sixty bushels of Newcastle lump coal, will make 92 bushels of coke.

Sixty bushels of Newcastle slack, or fine coal, will make 85 bushels of coke.

Sixty bushels Pictou or Virginia Coal, will make 75 bushels coke of an inferior quality compared with the above.

A bushel of the best coke weighs 32 lbs.

The production of coke by weight is about $\frac{2}{3}$ that of coal.

Coal furnishes 60 to 70 per cent. of coke by weight.

1 lb. of coke will evaporate in a common locomotive boiler $7\frac{1}{8}$ lbs. of water at 212° into steam.

TABLE, showing the weights, evaporative powers per weight, bulk and character of Fuel.

DESIGNATION.	Specific Gravity.	Weight per cubic foot.	Lbs. of steam from water at 212° by 1 lb. fuel.	Lbs. of steam from water at 212° by 1 cubic ft. of fuel.	Weight of clinker from 100 lbs. of coal.	No of cubic feet required to show a ton.
<i>Bituminous.</i>						
Cumberland max.,	1.313	52.92	10.7	573.3	2.13	42.3
“ min.,	1.337	54.29	9.44	532.3	4.53	41.2
Blossburgh,	1.324	53.05	9.72	522.6	3.40	42.2
Midlothian screened, . .	1.283	45.72	8.94	438.4	3.33	49.
“ average,	1.294	54.04	8.29	461.6	3.82	41.4
Newcastle,	1.257	50.82	8.66	453.9	3.14	44
Pictou,	1.318	49.25	8.41	478.7	6.13	45.
Pittsburgh,	1.252	46.81	8.20	334.1	.94	47.8
Sydney,	1.338	47.44	7.99	386.1	2.25	47.2
Liverpool,	1.262	47.88	7.84	411.2	1.86	46.7
Clover Hill,	1.285	45.49	7.67	359.3	3.86	49.2
Cannelton, Ia.,	1.273	47.65	7.34	360.	1.64	47.
Scotch,	1.519	51.09	6.95	369.1	5.63	48.8
<i>Anthracite.</i>						
Peach Mountain,	1.464	53.79	10.11	531.3	3.03	41.6
Forest Improvement, . .	1.477	53.66	6.06	577.3	.81	41.7
Beaver Meadows, No. 5, .	1.554	56.19	9.88	572.9	.60	39.8
Lackawanna,	1.421	48.89	9.79	493.	1.24	45.8
Beaver Meadows, No. 3, .	1.617	54.93	9.21	526.5	1.01	40.7
Lehigh,	1.590	55.32	8.93	515.4	1.08	40.5
<i>Coke.</i>						
Natural Virginia,	1.323	46.64	8.47	407.9	5.31	48.3
Midlothian,		32.70	8.63	282.5	10.51	68.5
Cumberland,		31.57	8.99	284.	3.55	70.9
<i>Wood.</i>						
Dry Pine wood,		20.01	4.69	98.6		106.6

Prof. W. R. Johnson.

N. B.—The above are the extreme effects; for practical use let a deduction of $\frac{1}{8}$ be made from the above.

Combustible matter of fuel.

The quantity of combustible matter of fuel, if the weight and other circumstances be equal, may be learnt from the ashes, or residuum, left after the combustion. For example, good Newcastle coal contains a greater portion of combustible matter than Nova Scotia coal, and leaves behind a smaller amount of earthy and incombustible substance. The heating power, and consequent value, of different kinds of fuel, is affected by this circumstance, though by no means dependent on it. The fitness of fuel for various purposes is furthermore affected by the facility with which it gives off a part of its combustible matter in the form of vapor or gas, which, being burnt in that state, produces *flame*. For example, the bituminous coals abound in volatile matter, which, when ignited, supports a powerful blaze. On the other hand, the Lehigh and Rhode Island coals are destitute of bitumen, and yield but little flame. It is from similar causes that dry pine wood produces a powerful blaze, while its charcoal yields comparatively little. A blaze is of great service where heat is required to be applied to an extensive surface, as in reverberating furnaces, ovens, glass-houses, &c. But when an equable, condensed, or lasting fire is wanted, the more solid fuels, which blaze less, are to be preferred.

TABLE, showing the heating power of different combustibles.

Designation.	Lbs. of water heated 1° by 1 lb. of substance.	Designation.	Lbs. of water heated 1° by 1 lb. of substance.
Alcohol.....	11,000	Coal, Newcastle.....	9,230
Olive Oil.....	14,500	“ Welsh.....	11,840
Beeswax.....	14,000	“ Anthracite.....	9,560
Tallow.....	15,000	“ Cannel.....	9,000
Oak, seasoned.....	4,600	Coke.....	9,110
“ kiln-dried.....	5,960	Peat.....	3,250
Pine, seasoned.....	5,466		

TABLE, showing the effects of heat upon certain bodies.

Designation.	Fahrenheit.	Designation.	Fahrenheit.
Gold melts.....	1983°	Tin melts.....	421°
Silver “.....	1850°	Water boils.....	212°
Copper “.....	2160°	Alcohol “.....	175°
Brass “.....	1900°	Ether.....	93°
Iron, red hot in daylight.....	1077°	Heat of human blood.....	98°
“ “ twilight.....	884°	Water freezes.....	32°
Common fire.....	790°	Strong wine freezes.....	20°
Zinc melts.....	740°	Brandy “.....	7°
Quicksilver boils.....	630°	Mercury “.....	—39°
Linseed Oil “.....	600°	Greatest cold ever produced*.....	—220°
Lead melts.....	594°	Snow and salt, equal parts.....	0°
Bismuth melts.....	476°	Acetous fermentation begins... “ “ ends....	78° 88°
Tin and Bismuth, equal parts, melts.....	283°	Phosphorus burns.....	68°

TABLE, showing the relative value of the following fuels by weight.

Designation.	Value.	Designation.	Value.
Seasoned oak.....	125	Charcoal.....	285
Oak, kiln-dried.....	140	Peat.....	115
Hickory.....	137	Welsh coal.....	312
White pine.....	137	Newcastle “.....	369
Yellow pine.....	145	Anthracite “.....	250
Good Coke.....	285		

* The lowest temperature hitherto attained, —220°, is produced by evaporating in vacuo a mixture of solid (condensed) protoxide of nitrogen, carbonic acid, and bisulphide of carbon.

TABLE, showing the number of gallons of water which may be lifted to various heights by the consumption of 112 lbs. of coal, the pumping apparatus being good, and adapted to the power of the steam engine.

Height.	Gallons.	Height.	Gallons.
1 foot.....	1,600,000	9 feet... ..	177,777
2 "	800,000	10 "	160,000
3 "	533,333	11 "	145,454
4 "	400,000	12 "	133,333
5 "	320,000	13 "	123,076
6 "	266,666	14 "	114,444
7 "	228,571	15 "	106,666
8 "	200,000	16 "	100,000

NOTES.—The evaporative power of 1 lb. of *bituminous* coal applied to a steam boiler, is from 6 to 9 lbs. fresh water in the boiler, under a pressure of 30 lbs. to the square inch, evaporated into steam. Cumberland coal being the strongest, and Scotch coal the weakest.

The evaporative power of *anthracite* coal, aided by a blast, is from $7\frac{1}{2}$ to $9\frac{1}{2}$ lbs. of fresh water evaporated into steam for 1 lb. of coal.

In practical evaporating power $2\frac{1}{2}$ to $2\frac{3}{4}$ lbs. of wood is equivalent to 1 lb. of bituminous or anthracite coal.

One cord of the ordinary seasoned fire-wood is equal in evaporating power to 12 bushels (960 lbs.) of Pittsburgh coal.

One ton of Cumberland coal is equal in evaporating power to $1\frac{1}{4}$ tons of anthracite coal, and equal to 2.12 cords of dry pine wood.

One ton of anthracite coal is equal to $1\frac{3}{4}$ cords of dry pine wood.

Each cubic foot of water evaporated in a boiler at the

pressure of the atmosphere, will heat 2,000 cubic feet of inclosed air to an average temperature of 75°.

Each square foot of surface steam-pipe will warm 200 cubic feet of space.

One pound of anthracite coal in a cupola furnace will melt 5 to 10 lbs. of cast iron.

80 bushels of bituminous coal in an air furnace will melt 10 tons of cast iron.

Small or fine coal produces about $\frac{3}{4}$ the effect of large coal of the same kind.

TABLE, *showing the price of parts of a cord of wood, at certain rates per cord.*

FEET	\$1.50	\$1.75	\$2.00	\$2.25	\$2.50	\$2.75	\$3.00	\$3.25
1	0 01	0 01	0 01	0 02	0 02	0 02	0 02	0 02
2	0 02	0 02	0 03	0 03	0 04	0 04	0 05	0 05
3	0 03	0 04	0 04	0 05	0 06	0 06	0 07	0 07
4	0 05	0 06	0 06	0 07	0 08	0 09	0 09	0 10
5	0 06	0 07	0 08	0 09	0 10	0 11	0 12	0 13
6	0 07	0 08	0 09	0 11	0 12	0 13	0 14	0 15
7	0 08	0 10	0 11	0 12	0 14	0 15	0 16	0 17
8	0 09	0 11	0 12	0 14	0 16	0 18	0 19	0 20
16	0 19	0 22	0 25	0 28	0 31	0 35	0 37	0 40
24	0 28	0 33	0 37	0 42	0 47	0 52	0 56	0 61
32	0 38	0 44	0 50	0 56	0 63	0 69	0 75	0 81
40	0 47	0 55	0 63	0 70	0 78	0 86	0 94	1 02
48	0 56	0 66	0 75	0 84	0 94	1 03	1 12	1 22
56	0 61	0 77	0 88	0 98	1 09	1 20	1 31	1 42
64	0 75	0 83	1 00	1 13	1 25	1 38	1 50	1 62
72	0 84	0 98	1 13	1 27	1 41	1 55	1 69	1 83
80	0 94	1 09	1 25	1 41	1 56	1 72	1 88	2 03
84	0 98	1 15	1 31	1 48	1 64	1 81	1 97	2 13
88	1 03	1 20	1 38	1 55	1 72	1 89	2 06	2 23
92	1 03	1 26	1 44	1 62	1 80	1 98	2 15	2 33
96	1 13	1 31	1 50	1 69	1 88	2 06	2 25	2 44
104	1 22	1 42	1 63	1 83	2 03	2 23	2 44	2 64
112	1 31	1 53	1 75	1 97	2 19	2 41	2 62	2 84
120	1 41	1 64	1 88	2 11	2 34	2 58	2 81	3 05
128	1 50	1 75	2 00	2 25	2 50	2 75	3 00	3 25


· **EXPLANATION.**—Find the number of feet in the left-hand column of the table ; then the price at the top of the page, and trace the line and column until they meet, and you will find the amount in dollars and cents.

EXAMPLE.—If a load of wood contains 98 feet, at two dollars and a half per cord—first find the amount of 96 feet, which is \$1.88 ; and then add the value of 2 feet (4 cents), making \$1.92. So of all similar examples.

Should the price per cord exceed the amount in the preceding table, the price of the parts may be found by adding or doubling, as per example, for \$3.50 double \$1.75 ; for \$3.75 add \$2.00 and \$1.75 ; for \$4.00 double \$2.00 ; for \$5.00 double \$2.50, &c.



FENCES.



In the newer portions of the country, where land is cheap and timber abundant, the old-fashioned zig-zag, or "*Virginia worm fence*," still prevails. It does not cost one-third the amount required for good post or board fence. Some are constructed altogether of rails, without any bracing or support at the corners, and are, of course, easily thrown down by cattle and the wind. They are, however, usually braced in one of the following modes:

1. By stakes and riders—either single or double riders.

2. By upright stakes, opposite each other, and placed in the *obtuse* corners, driven into the ground, and tied at the top by a wire or withe.

3. By upright stakes placed in the *acute* corners, driven into the ground, and tied at the top as above described.

4. By wedging one end of a rail into the acute corner, and letting the other end rest on the ground.

5. By placing the riders, or long poles, in a straight line on the top and at the centre of the fence, and then placing upright stakes in each inner corner, between the rider and the fence, the lower end resting on the ground and the other wedged tightly between the top and the rider.

The rails for this species of fence are cut different lengths in different sections of the country, and, indeed, in the same section. Much depends upon the nature of the timber, and much also on the kind of ground on which the fence is to be laid. Some are cut 12 feet, some 14, and some even 16½ feet or 1 rod in length. The usual lengths, however, are 12 and 14 feet.

The rails are laid at different angles; some deflecting 6 feet, some 7, and some 8 feet from a right line. The more they deflect, or in other words, the *crookeder* they are laid, the firmer the fence will be; but more rails will be required and more space occupied. The deflection for a 12 foot rail is usually 6 feet; for a 14 foot rail, 7 feet; and for a rod rail, 8 feet. A foot is generally allowed at each end for the lap.

Some fences are built 5 rails high, some 6, and some 7—the rider making an additional rail high. The height, as well as the spaces between the rails, are mostly regulated by statute in the different States. The majority of these statutes require the fence to be not less than 5 feet high, with interspaces between the rails of not more than 4 inches, to a height of 4 feet.

The number of rails, stakes, and riders required to build a certain amount of fence has hitherto been pretty much guess-work; and often the farmer, before he can finish his fence, has to quit it, and go and split more rails, or gear up and haul a few more loads. It is hoped that the following tables will obviate that necessity, by enabling him to tell within a few rails how many will be required to build a given amount of fence.

TABLE, *showing the number of rails, stakes, and riders required for each 10 rods of fence.*

Length of rail.	Deflection from right line	Length of panel	Number of panels	Number of rails for each 10 rods.			Number of stakes.	Number of riders, (single.)
				5 rails high.	6 rails high.	7 rails high.		
Feet.	Feet.	Feet.	Feet.					
12	6	8	20½	103	123	144	42	21
14	7	10	16½	83	99	116	34	17
16½	8	12	13½	69	84	95	28	14

NOTE.—Should the number of rods exceed 10, the requisite number of rails, stakes, and riders can be found by multiplying. For instance, should the length of fence be 100 rods, multiply the above number by 10; should it be 75 rods, multiply the above number by 7½; for 77 rods, multiply by 7½, and so forth.

Post and rail fence.

Post and rail is a more costly fence, but much better, and in the end more economical. There is not such a waste of either timber or land.

The rails are also cut of different lengths ; some 10, some 12, some 14, and some $16\frac{1}{2}$ feet, or 1 rod. Formerly, about 6 inches at each end were allowed for the lap, but more recently a foot has been allowed, as the longer the lap the stronger and firmer the fence. They are from 5 to 8 rails high ; posts set in the ground from 2 to 3 feet.

TABLE, *showing the number of rails and posts required for each 10 rods of post and rail fence.*

Length of rail—feet.	Length of panel—feet	Number of panels.	Number of posts.	Number of rails for each 10 rods.			
				5 rails high	6 rails high.	7 rails high	8 rails high.
10	8	$20\frac{5}{8}$	21	103	123	144	165
12	10	$16\frac{3}{4}$	17	83	99	116	133
14	12	$13\frac{3}{4}$	14	69	84	95	109
$16\frac{1}{2}$	$14\frac{1}{2}$	$11\frac{1}{2}$	12	57	69	81	93

NOTE.—Should the length exceed 10 rods, the additional number of posts and rails may be found by multiplying, as directed in the note to the preceding table.

Post and board fence.

Where timber is plenty and saw-mills abound, or where lumber is cheap, post and board fence is economical.

The boards are usually sawed 16 feet long, and the posts set 8 feet apart, 3 feet in the ground.

The fence is usually 5 boards high ; the bottom, or first board 10 inches wide ; the second 8, the third 6, and the

fourth and fifth 5 inches wide. They may be wider or narrower, as cost, taste, or use may dictate.

The first, third, and fifth boards are joined on one post, and the second and fourth joined on the next post.

To find the number of feet of boards required for each rod of post and board fence.

RULE.—Add the different widths of the boards, in inches, together, and divide the sum by 12 for the width in feet; then multiply the width by $16\frac{1}{2}$, and the product will be the number of feet, board measure, required for each rod of fence.

EXAMPLE.—Required, the number of feet, board measure, for each rod of fence, 5 boards high, the various widths of the boards being 10, 8, 7, 6 and 5 inches?

SOLUTION.— $10 + 8 + 7 + 6 + 5 = 36 \div 12 = 3 \text{ ft.} \times 16\frac{1}{2} = 49\frac{1}{2}$ feet. *Ans.*

To find the number of posts required for a given length of post and board fence.

RULE.—Reduce the number of rods to feet by multiplying by $16\frac{1}{2}$, and divide the product by the number of feet the posts are set apart; the quotient will be the number of posts required.

EXAMPLE.—Required, the number of posts for a post and board fence 160 rods long; posts set 8 feet apart?

SOLUTION.— $160 \times 16\frac{1}{2} = 2640 \div 8 = 330$. *Ans.*

HEDGE PLANTS.

The following, for the cultivation of hedges, is the condensed experience of the most successful and practical hedge-growers in the United States, and especially in the West.

Directions for Setting.—During the summer or fall thoroughly manure, plough as deep as possible a strip from five to eight feet wide, leave a dead furrow in the line where the hedge is to be set. In the following spring back furrow to the hedge-line, then harrow down smooth. Stake the ground, and by means of a line make a plain mark, then with a spade placed at right angles across the mark, push the blade in the soil to its full length at an angle of about forty-five degrees. Let an assistant place the plants under the back of the spade on the line of the mark, about one inch below the depth they stood in the nursery, and about eight inches apart. Pack the ground firmly around the plants, and mulch the ground to keep moist. Cultivate until the first of August. Before frost in the fall, back furrow and cover with coarse manure or straw, and in the spring uncover and cultivate as before. Replace all missing or feeble plants with strong ones.

Trimming.—The hedge should not be trimmed until three years old, when one-half or two-thirds should be cut

nearly off close to the ground and laid down at an angle of thirty degrees from the ground. Trim once a year in July, and do not allow the hedge to exceed twenty inches broad. The fourth year in the spring, before the buds start, take off about one-half the last year's growth. Leave the lower branches a little longer than the top, and aim to give the hedge some regular uniform shape. The hedge should be allowed to gain from eight to twelve inches annually, until it has reached the desired height.

To Preserve Plants during the Winter.—Cut a trench in a dry piece of ground at an angle of forty-five degrees, place the bundles in the trench, and cover with dirt from a new trench from six to eight inches in front, and so continue until all are trenched. Cover the plants two inches deep, firmly packing the ground around them. After the ground is frozen two inches deep, cover the whole with straw from twelve to eighteen inches; after which cover the whole bed with dirt about a foot thick. Encircle with a ditch so that no water can reach the plants. Plants can also be kept in a cellar, well covered in sand, but be careful not to expose to the sun or dry wind, in setting in the spring.

Setting Evergreens.—Cultivate and set as before, but the ground should not be manured within six months of setting the plants. Chip-dirt or rotten leaves are preferable for a mulch.

Hedge Plants.—Osage Orange.—The Osage Orange stands at the head of the list of hedge plants. It is much

planted where fencing timber is scarce, in the latitude of the Middle and Southern States. It is hardy and grows vigorously, and its thorns are absolute proof against the depredations of domestic animals, and even boys retreat from contact with them. It makes a beautiful hedge when properly pruned, but when neglected it gets beyond all control. In the Northern and Eastern States, it is liable to be killed by the frost.

Honey Locust.—This thorny, vigorous, and hardy plant has no superior as a farm hedge. It requires two annual prunings, in June and September, to keep it within control. It flourishes as far north as Canada, and for the Middle and Southern States it yields only to the Osage Orange. It is easily propagated by setting the plants about six inches apart. Some prefer sowing the seed on the line of the proposed hedge.

Buckthorn.—This plant is a native of America, and would be one of the best hedge plants did it not lack a supply of thorns.

Privet.—This thornless shrub is easily propagated from cuttings, and thickens well when set in a hedge. Its foliage is rich, and in the spring it is decorated with an abundance of beautiful small white flowers. It cannot be successfully cultivated north of the latitude of Philadelphia.

Hawthorn.—The hawthorn, so common in England, does not thrive so well in our climate.

Evergreen Hedges.—*Norway Spruce.*—A hedge of this beautiful tree should be set about four or five inches apart, and the plants not over four feet high. The side branches should be pruned, and the leaders cut out. Afterwards it should be trimmed the same as other hedges. The soil should be kept rich to insure a vigorous growth.

Arbor Vitæ.—In consequence of the cheapness of the common *Arbor Vitæ*, for an ornamental hedge, it has superseded all others. Though inferior to the Siberian species, yet it will be a long time before it will yield its place to it. Being hardy and sure to flourish under ordinary treatment, it is a valuable hedge plant.

Hemlock.—The hemlock, when properly pruned, makes a thick and beautiful hedge. With a foliage ever of the richest green, and adapted to all the northern latitudes, as a hedge plant it has no superior if an equal. Although hardy, it is somewhat difficult to transplant. Select a rainy day when the ground is wet, being careful not to expose the roots to the light or air. As soon as planted mulch with coarse manure or chip-dirt.



WIRE FENCES.

Wire fences have this advantage over hedges and other fences: they take up but little space, with no exhaustion of the soil, are not blown about by the wind; are durable, economical, and make a good protection against cattle, sheep, and other animals. For enclosing lawns and gardens, many of the designs offered in market are very desirable and ornamental. For a farm fence, such as any farmer can put up, annealed wire of the size No. 6 or 8* is preferable; for the protection of cattle five wires are sufficient; for sheep and lambs, seven should be used.

In building the fence a post six inches square or larger should be set at each end, and securely braced, from which to stretch the wire; the intervening posts should be set from eight to ten feet apart. Through these holes should be bored with a $\frac{1}{4}$ -inch brace-bit, and at appropriate distances apart, according to the protection required. Instead of putting the wires through the posts, they are often fastened by means of staples made of the same material. In putting up the wires they should be stretched as tightly as possible, care being taken in splicing that they be well secured, which can be best done by means of narrow blacksmith's tongs.

Suitable wire can be bought for 8 or 10 cents per pound, making a fence of six wires cost about 40 cents per rod; this does not include posts and labor of setting.

* The size of wire is graded from No. 1, and upwards. No. 9 is the common telegraph wire.

HUMAN STRENGTH.

The *force* of a single man, unaided by machinery, and working to the best advantage, is equivalent to the raising of 70 lbs. 1 foot per second for ten hours in a day.

The *maximum power* of a strong man, exerted for $2\frac{1}{2}$ minutes, is equivalent to 18,000 lbs. raised one foot in a minute.

A man of ordinary strength exerts a force of 30 lbs. for 10 hours in a day with a velocity of $2\frac{1}{2}$ feet in a second, which is about equal to 4500 lbs. raised 1 foot in a minute.

The average weight of men is 150 lbs. each.

A man travels, without a load, on level ground, for $8\frac{1}{2}$ hours a day, at the rate of $3\frac{7}{10}$ miles an hour, or $31\frac{1}{4}$ miles per day. He can carry 111 lbs. 11 miles in a day.

A porter going short distances, and returning unloaded, carries 135 lbs. 7 miles in a day. He can carry, in a wheelbarrow, 150 lbs. 10 miles a day.

An average strong man will, for a short period, exert a force with a—

	lbs.		lbs.
Drawing knife.....equal to	100	Pincers, compression.....equal to	60
An auger, both hands... "	100	A hand-plane	50
A screw-driver, 1 hand.. "	84	A hand-saw..... "	36
A bench-vice handle.... "	72	A thumb-vice..... "	45
A chisel, vertical pressure "	72	A brace-bit, revolving..... "	16
A windlass..... "	60		

HORSE POWER.

Before the invention and improvement of the steam-engine, the force of horses was very extensively used as a motive power; and although its application to machinery is now much less frequent, it is still resorted to, especially in places where fuel is expensive. For ordinary farm labor, it will probably never be superseded. The following are some of the more important facts relating to the horse and horse-power:—

The ordinary work of a horse is taken at 22,500 lbs. raised 1 foot in a minute, for 8 hours a day.

The strength of a horse is equivalent to that of 5 men.

A *draught-horse* can draw 1600 lbs. 23 miles a day on a level road, weight of carriage included.

In a *horse-mill*, he moves at the rate of 3 feet per second on a track 25 feet diameter, and with the machine exerts the power of $4\frac{1}{2}$ horses.


He occupies in a stall a front of $4\frac{1}{2}$ feet and a depth of 10 feet.

The average weight of horses is 1000 lbs. each.

A horse travels 400 yards, *at a walk*, in $4\frac{1}{2}$ minutes; 400 yards, *at a trot*, in 2 minutes; and 400 yards, *at a gallop*, in 1 minute.

A horse will *carry* 250 lbs. 25 miles a day of 8 hours.

HORSE POWER ILLUSTRATED.—LOADING A SHIP.





A horse will live 25 days without solid food, merely drinking water. He will live 17 days without either eating or drinking. He will live only 5 days when eating solid food, without drinking.

He attains his full growth in 5 years, and will live 25. His average life is 16 years.

Horse-power as applied to the measurement of steam-engines and waterfalls was first applied by James Watt, the inventor of the steam-engine. From a series of experiments he ascertained that the average strength of a horse was sufficient to raise 33,000 lbs. one foot per minute,* and this unit has been adopted in this country and in England as a general measure of power.

A waterfall is thus said to have a horse-power for every 33,000 lbs. of water passing a given point per minute for each foot of the fall. To compute the power of a waterfall is given the following

RULE.—Divide the continued product of the width, the depth, the velocity of the water per minute, the height of the fall, and the weight of a cubic foot of water ($62\frac{1}{2}$ lbs.) by 33,000.

EXAMPLE.—The flume of a mill is 10 feet wide, the water is 3 feet deep, the velocity is 100 feet per minute, and the fall 11 feet. What is the horse-power of the fall?

OPERATION.— $(10 \times 3 \times 100 \times 11 \times 62\frac{1}{2}) \div 33,000 = 62\frac{1}{2}$ horse-power.

* This is done by means of compound pulleys.

The power of a steam-engine is estimated by the following

RULE.—Divide the continued product of the area of the piston in inches, the mean pressure per square inch in pounds, the length of the stroke in feet, and the number of strokes per minute by 33,000.

EXAMPLE.—The area of the piston of a steam-engine is 40 inches, the pressure is 60 lbs. per square inch, the length of the stroke is 3 feet, and it makes 30 strokes per minute. What is the horse-power?

OPERATION.— $(40 \times 60 \times 3 \times 30) \div 33,000 = 6\frac{1}{2}$ horse-power (nearly).

Water-wheels lose from 10 to 50 per cent. of the power, and the actual power of the steam-engine is less than that indicated by the horse-power, owing to a loss by friction, the amount of which depends upon the arrangement of the engine and the perfection of the workmanship.

TABLE, *showing the labor one horse is able to perform at different rates of speed on canals, railroads, and turnpikes. Drawing force, $83\frac{1}{3}$ lbs.*

Speed per hour. Miles.	Duration of day's work—hours	Useful effect for 1 day in tons, drawn 1 mile.		
		On canal—tons	On a railroad—tons.	On a turnpike—tons.
$2\frac{1}{2}$	$11\frac{1}{2}$	520	115	14
3	8	243	92	12
$3\frac{1}{2}$	6	154	82	10
4	$4\frac{1}{2}$	102	72	9
5	$2\frac{3}{4}$	52	57	7.3
6	2	30	48	6
7	$1\frac{1}{2}$	19	41	5
8	$1\frac{1}{4}$	12.8	36	4.5
9	$1\frac{1}{6}$	9.	32	4.
10	$\frac{3}{4}$	6.5	28.8	3.6




TABLE, showing how much one team and plough will perform in a day, in acres and tenths.

Width of furrow in inches.	Acres and tenths.	Width of furrow in inches.	Acres and tenths.	Width of furrow in feet.	Acres and tenths.	Width of furrow in feet.	Acres and tenths.
5	1.0	12	2.4	2	4.8	5½	13.2
6	1.2	14	2.8	2½	6.0	6	14.4
7	1.4	16	3.2	3	7.2	6½	15.6
8	1.6	18	3.6	3½	8.4	7	16.8
9	1.8	20	4.0	4	9.6	7½	18.0
10	2.0	22	4.4	4½	10.8	8	19.2
11	2.2			5	12.0		

NOTE.—The above table is constructed on the presumption that the team moves at the rate of about 3 feet per second, or 2 miles per hour, for 10 hours per day. Horses and mules in good condition will do this

FREIGHTS—QUANTITY OF GOODS WHICH
COMPOSE A TON IN SHIPPING.



Wharf Scene in New York.

From By-laws of the New York Chamber of Commerce.

Resolved, That when vessels are freighted by the ton, and no special agreement is made between the owner of the vessel and freighter of the goods, respecting the proportion of tonnage which each particular article shall be computed at, the following regulation shall be the standard of computation :—

That the articles, the bulk of which shall compose a ton, to equal a ton of heavy materials, shall be in weight as follows: 1568 lbs. of coffee in casks, 1830 lbs. in bags; 1120 lbs. of cocoa in casks, 1307 lbs. in bags.

952 lbs. pimento in casks, 1110 in bags.

Eight barrels of flour, 196 lbs. each.

Six barrels of beef, pork, tallow, pickled fish, pitch, tar, and turpentine.

Twenty hundred pounds of pig and bar iron, potashes, sugar, logwood, fustic, Nicaragua wood, and all heavy dye-woods, rice, honey, copper ore, and all other heavy goods.

Sixteen hundred pounds of coffee, cocoa, and dried codfish, in bulk, and twelve hundred pounds of dried codfish in casks of any size.

Six hundred pounds of ship bread in casks, seven hundred in bags, and eight hundred in bulk.

Two hundred gallons (wine-measure), reckoning the full contents of the casks, oil, wine, brandy, or any kind of liquors.

Twenty-two bushels of grain, peas, or beans, in casks.

Thirty-six bushels of grain in bulk.

Thirty-six bushels of European salt.

Thirty-one bushels of salt from the West Indies.

Twenty-nine bushels of sea-coal.

Forty feet (cubic measure) of mahogany, square timber,


oak plank, pine, and other boards, beavers, furs, peltry, bees wax, cotton, wool, and bale goods of all kinds.

One hogshead of tobacco, and ten hundred pounds of dry hides.

Eight hundred pounds of China raw silk, ten hundred pounds of net bohea, and 800 green tea.



UNITED STATES OR FEDERAL MONEY.



Stamping Coin at the United States Mint.

Money is value, or the representative of value, used for the purposes of exchange. In different countries, at different times, various articles have been used for money, such as oxen, pieces of leather stamped, shells, wampum, iron, nails, &c. Gold and silver, at present, are used almost exclusively for money. They are called precious metals.

Paper money is a substitute for coin.

Uncoined gold and silver is called *bullion*.

Coin is a piece of metal of known weight used for money, the value of which is stamped on it.

Currency is the money of circulation.

Tokens are coins whose intrinsic value is below that assigned them by law. Such coins are said to be coins in *billion*.

United States or Federal money is a decimal currency.

TABLE.

10 mills (m.)	1 cent	ct.		
10 cents	• 1 dime	d.	100	mills.
10 dimes	1 dollar	\$	1000	" 100 cents.
10 dollars	1 eagle	E.	10000	" 1000 cents 100 dimes.

COINS.—The gold coins are the *double-eagle*, *eagle*, *half-eagle*, *quarter-eagle*, *three-dollar piece*, and *dollar*.



NOTES.—1. The fifty-dollar piece is not a legal coin. The

copper half-cent is no longer coined. The mill is not a coin.

2. Gold coins contain 9 parts of gold and 1 part of an alloy of silver and copper.

3. The silver coins are the *dollar*, *half-dollar*, *quarter-dollar*, *dime*, *half-dime*, and *three-cent piece*.



4. Silver coins contain 9 parts silver and 1 part copper, except the three-cent piece, which is 3 parts silver and 1 part copper.

5. The nickel coins are the *cent*, the new *three-cent*, and new *five-cent pieces*.

6. The nickel cent contains 88 parts copper and 12 parts nickel.

7. The copper coins are the *cent* and *two-cent pieces*.



8. The two-cent and cent pieces are made of nickel and copper.

| The term *dollar* is supposed to be derived from the German "thaler," pronounced *ta-ler*.

The term *dime* means ten, *cent* a hundred, and *mill* a thousand.

The origin of the dollar-mark is uncertain; some think it the combination of U. S., others that it is an imitation of the dollars and scroll on the "pillar-dollar."

1 eagle (gold) weighs 258 troy grains.

1 dollar (silver) " 412.5 "

1 cent (copper) " 168 "

23.2 grains of pure gold = \$1.00.

Gold coin of the United States, prior to 1834, like that of England, = 88.8 cents per dwt. By act of Congress of 1834, its value was made 94.8 cents per dwt. The old United States Eagle, coined previous to 1834, is worth \$10.66·8.

ENGLISH MONEY.

English or Sterling Money is the currency of Great Britain.

TABLE.

4 farthings (far. or qr.)	make 1 penny,	marked d.
12 pence	“ 1 shilling,	“ s.
20 shillings	“ 1 pound or sovereign, £, sov.	
21 shillings	“ 1 guinea, marked guin.	

COINS.—The gold coins are the *sovereign* (£1), and the *half-sovereign* (10s.).



The silver coins are the *crown* (5s.), the *half-crown* (2s. 6d.), the *florin* (2s.), the *shilling* (12d.), *sixpenny-piece* (6d.), and *threepenny-piece* (3d.).



The bronze coins are the *penny*, *half-penny*, and *farthing*. Farthings are generally written as fractions of a penny, thus: 1 far. = $\frac{1}{4}$ d.; 2 far. = $\frac{2}{4}$ or $\frac{1}{2}$; 3 far. = $\frac{3}{4}$.



Canadian currency is decimal, and the denominations are the same as Federal money.

The franc is the unit of the French decimal currency,

and is worth \$0.186. The denominations are *francs* and *centimes*.



NOTES.—1. The symbol £ stands for the Latin word *libra*, a pound ; *s.* for *solidus*, a shilling ; *d.* for *denarius*, a penny ; *qr.* for *quadrans*, a quarter.

2. The term *sterling* is supposed to be derived from *Easterling*, a name formerly given to the early German traders.

3. The term *farthing* is derived from "four things," denoting the divisions on the old English penny.

AVOIRDUPOIS WEIGHT.



Avoirdupois weight is used for all ordinary purposes.

TABLE.

16 drams (dr.)	1 ounce,	marked oz.
16 oz.	1 pound,	" lb.
25 lb.	1 quarter,	" qr.
4 qr.	1 hundredweight,	" cwt.
20 cwt.	1 ton,	" T.
100 lb.	1 cental,	" c.

T.	cwt.	qr.	lb.	lb.	oz.	dr.	gr.*
1	=20	=80	=2000	1	=16	=256	=7000
	1	=4	=100		1	=16	=437 $\frac{1}{2}$
		1	=25			1	=27 $\frac{1}{2}$

NOTES.—1. The gross ton of 2240 lbs. was formerly in common use, but is now seldom used except at the United States Custom House and at the Pennsylvania coal mines.

2. Butter is usually packed for market in pails or firkins, which hold from 50 to 100 pounds.

3. The term *avoirdupois* is derived from the French “avoir du poids,” meaning *goods of weight*. *Cwt.* is formed from *c.*, *centum*, *wt.*, *weight*.

4. Most of the States have adopted the following

TABLE OF MISCELLANEOUS WEIGHTS.

196 lbs.	make 1 barrel of flour.
200 “	“ 1 “ beef, pork, or fish.
280 “	“ 1 “ salt at N. Y. Salt Works.
32 “	“ 1 bushel of oats.
48 “	“ 1 “ barley.
56 “	“ 1 “ corn or rye.
60 “	“ 1 “ wheat.
60 “	“ 1 “ beans.
14 “	“ 1 “ blue-grass-seed.
46 “	“ 1 “ castor-beans.
60 “	“ 1 “ clover-seed.
56 “	“ 1 “ flax-seed.
44 “	“ 1 “ hemp-seed.

* NOTE.—The exact weight of an avoirdupois dram is $27\frac{1}{32}$ troy grains.

60 lbs.	make 1 bushel of	peas.
60 “	“ 1 “	potatoes.
45 “	“ 1 “	timothy-seed.
57 “	“ 1 “	onions.
28 “	“ 1 “	apples or peaches (dried).
50 “	“ 1 “	salt.

A sack of wool is 22 stone, that is, 14 lbs. to the stone, 308 lbs.


A pack of wool is 17 stone 2 lbs.=240 lbs.—a pack load for a horse.

A truss of hay is, new, 60 lbs.; old, 50 lbs.; straw, 40 lbs. *A load of hay* is 36 trusses. *A bale of hay* is 300 lbs.

A firkin of butter was formerly 56 lbs.

A bale of cotton is 400 lbs., but it is put up in different States varying from 280 to 720 lbs. Sea Island cotton is put up in sacks of 300 lbs.

AVOIRDUPOIS WEIGHT ILLUSTRATED.



1 firkin.



1 barrel.



1 barrel.



1 barrel.



1 bushel.



1 bushel.




1 bushel.



1 bushel.

TROY WEIGHT.




Troy weight is used in weighing gold, silver, and jewels, and in philosophical experiments.

TABLE.

24 grains (gr.)	make 1 pennyweight,	marked pwt.
20 pwt.	“ 1 ounce,	“ oz.
12 oz.	“ 1 pound,	“ lb.
$3\frac{1}{2}$ grains	“ 1 carat (diamond wt.)	“ k.

SCALE OF COMPARISON.

lb.	oz.	dwt.	gr.
1	= 12	= 240	= 5760
	1	= 20	= 480
		1	= 24
		1k.	= $3\frac{1}{2}$



NOTES.—1. A carat is a weight of about 3.2 grains, and is used by jewellers to weigh diamonds. The term *carat* is also used to denote the fineness of gold. When gold contains 18 parts pure gold and 6 parts alloy, which is usually silver and copper, it is said to be 18 carats fine. Gold 14 carats fine contains 14 parts pure gold and 10 parts alloy, &c.

2. The term *Troy* is derived from *Troyes*, where the weight was first introduced into Europe, about the 12th century.

3. The term *pennyweight* is derived from the weight of the old silver penny. The term *grain* is derived from the custom of using the grains of wheat, 24 of which were taken to determine the weight of a pennyweight.

4. The symbol *oz.* is derived from the Spanish word *onza*, an ounce; *lb.* is from the Latin *libra*, a pound.

5. The standard unit of weight is the troy pound. It equals the weight of 22.79 + cu. in. of distilled water at the temperature of 39° 83' F., the barometer being at 30 in.

APOTHECARIES' WEIGHT.




Apothecaries' weight is used in preparing prescriptions, but drugs and medicines are bought and sold by avoirdupois weight.

TABLE.

20 grains (gr.)	1 scruple,	marked sc. or ℥.
3 scruples	1 drachm,	" dr. or ʒ.
8 drachms	1 ounce,	" oz. or ʒ.
12 ounces	1 pound,	" lb. or ℔.

SCALE OF COMPARISON.

℔	℥	3	℥	gr.
1	=	12	=	96
			=	288
			=	5760
	1	=	8	=
				24
				=
				480
		1	=	3
				=
				60
			1	=
				=
				20



APOTHECARIES' FLUID MEASURE.

Apothecaries' fluid measure is used for measuring liquids in preparing medical prescriptions.

TABLE.

60 minims (℥)	1 fluid drachm,	marked f 3.
8 fluid drachms	1 fluid ounce,	" f 3.
16 fluid ounces	1 pint,	" O.
8 pints	1 gallon (wine meas.)	" Cong.

NOTE.—1. The pound, ounce, and grain are the same as in troy weight, the ounce being differently subdivided.

2. The symbols are supposed to be derived from the inscriptions upon the ancient monuments of Egypt.

3. One minim equals one drop.

LIQUID OR WINE MEASURE.



Liquid measure is, of course, used in measuring liquids.

TABLE.

4 gills (gi.)	1 pint,	marked pt.
2 pints	1 quart,	" qt.
4 quarts	1 gallon,	" gal.
$31\frac{1}{2}$ gallons	1 barrel,	" bbl.
2 barrels or 63 gallons	1 hogshead,	" hhd.

SCALE OF COMPARISON.

Wine Measure.

gal. qt. pt. gi. cu. in.

$$1=4=8=32=231$$

$$1=2=8=57\frac{3}{4}$$

$$1=4=28\frac{7}{8}$$

Dry Measure.

bu. pk. qt. pt. cu. in.

$$1=4=32=64=2150\frac{3}{4} \text{ nearly.}$$

$$1=8=16=537\frac{1}{2} \quad "$$

$$1=2=67\frac{1}{8} \quad "$$



NOTE.—1. The denominations barrel and hogshead are used in estimating the capacity of cisterns, reservoirs, vats, &c.

2. The barrel, hogshead, tierce, pipe, butt and tun, are the names of casks, which are usually gauged, having the number of gallons they hold marked on them.


3. Ale or beer measure, formerly used in measuring beer, ale, and milk, is now seldom used.

4. 1 gallon of pure water weighs nearly $8\frac{1}{8}$ lb. avoirdupois, hence a pint weighs about a pound.

5. The standard unit of wine measure is the gallon, which contains 231 cubic inches.


The Imperial, or British gallon, contains 277.274 cubic inches.

DRY MEASURE.



Dry measure is used in measuring vegetables and articles not fluid.

2 pints (pt.)	1 quart,	qt.
8 quarts	1 peck,	pk.
4 pecks	1 bushel,	bu.
36 bushels	1 chaldron,	cald.



NOTES.—The standard bushel is the Winchester, which contains 2150.42 cubic inches, or 77.627 lbs. avoirdupois of distilled water at its maximum density.

Its dimensions are $18\frac{1}{2}$ inches diameter inside— $19\frac{1}{2}$ inches

outside, and 8 inches deep, and when heaped to a cone 6 inches high, contains 2748 cubic inches.

The Imperial or British bushel contains 2218 cubic inches, so that 32 of their bushels are equal to 38 of ours.

Heaping Measure.—Potatoes, turnips and esculent roots, apples and other fruits, meal and bran, corn on the ear, and in some States, oats, are sold by the heaping bushel measure.

TABLE OF COMPARISON OF THE MEASURES OF CAPACITY.

1 gallon or 4 qt.	wine measure contains	231	cubic inches.
$\frac{1}{2}$ pk. or 4 qt.	dry measure	"	268 $\frac{1}{3}$ "
1 gallon or 4 qt.	beer measure	"	282 "
1 bushel	dry measure	"	2150 $\frac{1}{3}$ "

In England the following weights and measures are sometimes used :

WEIGHT.

3 pounds=1 stone, butchers' meat.
7 pounds=1 clove.
2 cloves=1 stone common articles.
2 stone=1 tod of wool.
6 $\frac{1}{2}$ tods=1 wey "
2 weys=1 sack "
12 sacks=1 last "
240 pounds=1 pack "

CLOTH MEASURE.

2 $\frac{1}{4}$ inches=1 nail.
4 nails=1 quarter.
4 quarters=1 yard.
3 quarters=1 Flemish ell.
5 quarters=1 English ell.
6 quarters=1 French ell.
4 $\frac{2}{3}$ quarters=1 Scotch ell.

DRY MEASURE.


2 quarts=1 pottle.
2 bushels=1 strike.
2 strikes=1 coom.
2 cooms=1 quarter.
5 quarters=1 load.
3 bushels=1 sack.
36 bushels=1 chaldron.

WINE MEASURE.


18 U. S. gal =1 runlet.
25 Eng. gal. or { =1 tierce.
42 U. S. gal. }
2 tierces=1 puncheon.
52 $\frac{1}{2}$ Eng. gal. or { =1 hogshead.
63 U. S. gal. }
2 hogsheads=1 pipe.
2 pipes=1 tun.
7 $\frac{1}{2}$ Eng. gal. =1 firkin of beer.
4 firkins=1 barrel "

TABLE OF THE COMPARISON OF WEIGHTS, &c.

1 U. S. pound Troy=5760 grs. Troy.	1 English yard=36 inches.
1 Eng. pound Troy=5760 " "	1 French metre=39.368 + inches.
1 pound Apoth. =5760 " "	1 U. S. bushel =2150.42 + cu. in.
1 U. S. pound Av. =7000 " "	1 Eng. " =2218.19+ "
1 Eng. pound Av. =7000 " "	1 U. S. gallon =231.
144 pounds Av. =175 lb. "	1 Eng. " =277.26+ "
1 French gramme =15.433 grs. Troy.	1 French litre =61.533+ "
1 U. S. yard =36 inches.	1 French are =119.664 sq. yds.



SQUARE MEASURE.



Square measure is used in calculating areas or surfaces, as of land, lumber, painting, paving, &c.

TABLE.

144 square inches (sq. in.)	make 1 square foot,	marked sq. ft.
9 square feet	" 1 square yard,	" sq. yd.
30 $\frac{1}{2}$ square yards	" 1 square rod,	" sq. rd., P.
40 square rods	" 1 rood, or qr. acre,	" R.
4 roods	" 1 acre,	" A.
640 acres	" 1 sq. mile or section,	" sq. m., sec.

SCALE OF COMPARISON.

A.	R.	P.	sq. yds.	sq. ft.	sq. in.
1	= 4	= 160	= 4840	= 43560	= 6272640.
1	=	40	= 1210	= 10890	= 1568160.
1	=	30 $\frac{1}{4}$	=	272 $\frac{1}{4}$	= 39204.
		1	=	9	= 1296.
				1	= 144.

NOTE.—Artificers usually estimate their work—1. In glazing and stone-cutting, by the square foot. 2. In painting, plastering, paper-hanging, &c., by the square yard. 3. In flooring, roofing, slating, &c., by the 100 square feet. 4. In bricklaying, by the thousand bricks, by the square yard, and 100 feet.

The painting of mouldings, cornices, &c., is estimated by measuring the entire surface.


When bricklaying is estimated by square measure, the work is understood to be 12 inches thick.

Surveyor's square measure is used in finding the area of land.

TABLE.

625 square links (sq. l.)	make 1 sq. rod,	marked sq. rd.
16 sq. rods	" 1 sq. chain,	" sq. ch.
10 sq. chains	" 1 acre,	" A
640 acres	" 1 sq. mile,	" sq. mi.
36 sq. miles (six miles square)	" 1 township,	" Tp.

LONG MEASURE.



Long measure is used for distances, &c.

TABLE.

12 lines or 3 barley-corns	1 inch,	
12 inches	1 foot,	marked ft.
3 ft.	1 yard,	" yd.
$5\frac{1}{2}$ yd.	1 rod,	" rd.
40 rd.	1 furlong,	" fur.
8 fur.	1 mile,	" mi.

SCALE OF COMPARISON.

mi.	fur.	rod.	yd.	ft.	in.
1	= 8	= 320	= 1760	= 5280	= 63360
	1	= 40	= 220	= 660	= 7920
		1	= $5\frac{1}{2}$	= $16\frac{1}{2}$	= 198
			1	= 3	= 36
				1	= 12

SURVEYORS' MEASURE.

Gunter's chain is used by land surveyors. It is 4 rods or 66 feet long, and contains 100 links.

TABLE.

25 links (li.)	1 rod,	rd.
4 rods	1 chain,	ch.
80 chains	1 mile,	mi.

TABLE OF MISCELLANEOUS LINEAR MEASURE.

3 inches	1 palm.	
4 inches	1 hand.	{ Used in measuring the height of horses at the shoulder.
9 inches	1 span.	
3 feet	1 pace or step.	
3.28 feet	1 metre.	
6 feet	1 fathom.	{ Used in measuring depths at sea.
880 fathoms	1 mile.	
3 geographical miles	1 league.	
60 " "	{ 1 degree.	{ Of latitude Of longitude on the equator.
69½ statute " "		

NOTE.—A hair's breadth is the 48th part of an inch.

A ship's cable is a chain, usually about 120 fathoms or 720 feet long, hence the term "cable length" in nautical language denotes about that distance.

NOTES.—1. A knot is a nautical or geographical mile. Thus, the phrase, "thirteen knots an hour," means thirteen geographical miles an hour.

2. 1 English mile equals 5280 feet, and 1 nautical, or geographical mile, equals 6086 feet.

3. The geographic mile equals about 1.15 English miles; the German short mile, about 3.9 English miles; the German long mile, about 5.75 English miles; the Prussian mile about 4.7 English miles; the Spanish common league, about 4.2 miles; and the Spanish judicial league about 2.6 miles.

4. Measures of length were at first derived from the different parts of the body, as the *finger*, *hand*, the *span*, or the length of the thumb and middle finger extended; *cubit*, or the length of the forearm; and the *fathom*, or the length of the two arms extended.

CLOTH MEASURE.

Cloth measure is used by merchants in the sale of cloth, ribbons, laces, &c.

TABLE.

2 sixteenths (16th)	1 eighth,	marked 8th, $\frac{1}{8}$ yd.
2 eighths	1 quarter,	“ qr., $\frac{1}{4}$ yd.
2 quarters	1 half,	“ hlf., $\frac{1}{2}$ yd.
4 quarters or 2 halves	1 yard,	“ yd.

NOTE.—The old system of measuring cloth is not now used. By it each yard is divided into 4 quarters, and each quarter into 4 nails, a nail being $2\frac{1}{4}$ inches. 3 quarters make a Flemish ell, 5 quarters an English ell, and 6 quarters a French ell.

CUBIC MEASURE.






TABLE.

1728 cubic inches (cu. in.)	1 cubic foot,	marked cu. ft.
27 cubic feet	1 cubic yard,	" cu. yd.
40 cubic ft. of round timber or }	1 ton or load,	" T.
50 cubic feet of hewn timber }	1 cord foot,	" cd. ft.
16 cubic feet	1 cord of wood,	" Cd.
8 cord feet or }		
128 cubic feet }		
24½ cubic feet	1 { perch or }	" Pch.
	stone, or }	
	masonry. }	

Cubic measure is used in estimating the contents of solids;
as wood, stone, capacity of cisterns, &c.


Cubic inch.


Cubic foot.



Cubic yard.

To find the cubic contents of any solid body.

RULE.—Multiply the length by the breadth, and that product by the thickness.

NOTES.—1. A load of earth contains a cubic yard, and weighs about 3250 lbs.

2. Railway and transportation companies estimate light freight by the number of cubic feet it occupies; but heavy freight is estimated by weight.

3. A pile of wood 4 feet wide, 4 feet high, and 8 feet long, contains 1 *cord*; and a *cord foot* is 1 foot in length of such a pile.

4. A perch of stone or masonry is $16\frac{1}{2}$ feet long, $1\frac{1}{2}$ feet wide, and 1 foot high, and contains $24\frac{3}{4}$ cubic feet.

5. A brick is usually 8 inches long, 4 inches wide, and 2 inches thick; hence 27 bricks make a cubic foot.

6. Joiners, painters, and masons make no allowance for windows, doors, &c. Masons make no allowance for the corners of the walls of houses or of cellars. The size of a

cellar is estimated by the measurement of the outside of the wall.

Ton weight and ton measure.—A ton of hay, or any other coarse bulky article usually sold by that measure, is 20 gross hundreds, that is 2240 lbs. But in many places it has become the custom to count only 2000 lbs. for a ton. In freighting ships, 42 cubic feet are allowed to a ton; in the measurement of timber, 40 solid feet if round, and 50 if square make a ton.

THE METRIC SYSTEM OF WEIGHTS AND MEASURES.*

The metric system of weights and measures had its origin in France during the Revolution in the year 1790. The following year a commission of scientific men was appointed by the government to select an appropriate unit, and as the result of their investigations the *ten-millionth* part of the earth's quadrant was chosen and called a *Metre*. To determine the unit of *weight* a cube of pure water at its greatest density, each edge of which is *one-hundredth* of a metre, was taken and called a *Gramme* (anglicized *gram*). The multiples and subdivisions were made to correspond to the decimal scale, hence its great simplicity.

This system was declared obligatory in France after Nov. 2, 1801; but no penalty was attached to non-conformity until after Jan. 1, 1841. The system has since been adopted wholly or in part by Spain, Belgium, Portugal, Holland, Great Britain, Greece, Italy, Norway, Sweden, Mexico, Guatemala, Venezuela, Ecuador, U. S. of Columbia, Brazil, Chili, San Salvador, and the Argentine Republic. In 1866

* The following article on the Metric System of Weights and Measures was prepared for this work by S. A. Felter, A.M., author of a well-known series of mathematical text-books.

Congress authorized the metric system in the United States by passing the following bill:—

AN ACT TO AUTHORIZE THE USE OF THE METRIC SYSTEM OF
WEIGHTS AND MEASURES.

Be it enacted by the Senate and House of Representatives of the United States of America in Congress assembled, That from and after the passage of this act, it shall be lawful throughout the United States of America to employ the weights and measures of the metric system; and no contract or dealing, or pleading in any court, shall be deemed invalid or liable to objection, because the weights or measures expressed or referred to therein are weights or measures of the metric system.

SEC. 2.—*And be it further enacted,* That the tables in the schedule hereto annexed, shall be recognized in the construction of contracts, and in all legal proceedings, as establishing, in terms of the weights and measures now in use in the United States, the equivalents of the weights and measures expressed therein in terms of the metric system; and said tables may be lawfully used for computing, determining, and expressing, in customary weights and measures, the weights and measures of the metric system.

The utility of the metric system commends itself, even at a glance, and hence it becomes important that all should become acquainted with it. It will doubtless soon come in to general use to the exclusion of all other systems of weight and measure. The following is a brief and condensed view

of the system, so clear and simple that a child can understand it :—

The Metric System of weights and measures is formed upon the decimal scale, and has for its base an invariable unit derived from nature, and called a *METRE*; and upon this unit all the units of weight and measure are based.

The Metre is the *ten-millionth* part of the distance from the equator to the pole; and is the principal unit of linear measure.

The Are is a square whose side is ten *metres*. It is the principal unit of superficial measure.

The Stere is a cube whose edge is a *metre*. It is the principal unit of solid or cubic measure.

The Litre is a cube whose edge is the *tenth* of a *metre*. It is the principal unit of all *measures of capacity*.

The Gram is the weight of a cube of pure water at its greatest density, whose edge is the *hundredth* part of a *metre*. A litre of water weighs 1,000 grams. It is the principal unit of weight.

The names of the derivative denominations are formed by joining a Latin or Greek prefix to the principal units. There are seven of these prefixes, derived as follows:

Latin.	{	MILLI,	from <i>Millesimus</i> , a <i>thousandth</i> .
		CENTI,	from <i>Centesimus</i> , a <i>hundredth</i> .
		DECI,	from <i>Decimus</i> , a <i>tenth</i> .

Greek.	DECA, <i>ten.</i>	
	HECTO, from <i>Hecaton</i> ,	<i>one hundred.</i>
	KILO, from <i>Chilioi</i> ,	<i>one thousand.</i>
	MYRIA, from <i>Myrioi</i> ,	<i>ten thousand.</i>

The formation of the tables can be seen at a glance by the following:—


<i>Milli</i>	}	METRE.	}	ARE.*	}	STERE.	}	LITRE.	}	GRAM.
<i>Centi</i>										
<i>Deci</i>										
<i>Deca</i>	}	METRE.	}	ARE.*	}	STERE.	}	LITRE.	}	GRAM.
<i>Hecto</i>										
<i>Kilo</i>										
<i>Myria</i>										

NAMES.	PRONUNCIATION.	ABB.	NAMES.	PRONUNCIATION.	ABB.
Millimetre	Mill'-e-mee'-ter	mm.	Hectostere	Hec'-to-steer	hs.
Centimetre	Sent'-e-mee'-ter	cm.	Kilostere	Kill'-o-steer	ks.
Decimetre	Des'-e-mee'-ter	dm.	Myriastere	Mir'-e-a-steer	mys.
<i>Metre</i>	Mee'-ter	m.	Millilitre	Mill'-e-li'-ter	ml.
Decametre	Dek'-a-mee'-ter	dkm.	Centilitre	Sent'-e-li'-ter	cl.
Hectometre	Hec'-to-mee'-ter	hm.	Decilitre	Des'-e-li'-ter	dl.
Kilometre	Kill'-o-mee'-ter	km.	<i>Litre</i>	Li'-ter	l.
Myriametre	Mir'-e-a-mee'-ter	mym.	Decalitre	Dek'-a-li'-ter	dcl.
Milliare	Mill'-e-âre	ma.	Hectolitre	Hec'-to-li'-ter	hl.
Centiare	Sent'-e-âre	ca.	Kilolitre	Kill'-o-li'-ter	kl.
Deciare	Des'-e-âre	da.	Myrialitre	Mir'-e-a-li'-ter	myl.
<i>Are</i>	Are	a.	Milligram	Mill'-e-gram	mg.
Decare	Dek'-âre	dka.	Centigram	Sent'-e-gram	cg.
Hectare	Hec'-târe	ha.	Decigram	Des'-e-gram	dg.
Kilare	Kill'-âre	ka.	<i>Gram</i>	Gram	g.
Myriare	Mir'-e-âre	mya.	Decagram	Dek'-a-gram	dkg.
Millistere	Mill'-e-steer	ms.	Hectogram	Hec'-to-gram	hg.
Centistere	Sent'-e-steer	cs.	Kilogram	Kill'-o-gram	kg.
Decistero	Des'-e-steer	ds.	Myriagram	Mir'-e-a-gram	myg.
<i>S'ere</i>	Steer	s.	Quintal	Quin'-tal	q.
Decastere	Dek'-a-steer	dks.	Tonneau	Tun'-no	T.

* The *a* in *deca* and *myria*, and the *o* in *hecto* and *kilo* are dropped when prefixed to *Are*.

LINEAR MEASURE.

Illustration.



NOTE.—By the accompanying illustration it will be seen that *one-tenth* of a metre, or ten centimetres, equals about $3\frac{1}{8}$ in., or a trifle short of 4 in.

This measure, as well as the other measures and weights, is written as whole numbers and decimals. The decimal point is placed at the right of the *unit*; thus, 4.167 *m.* may be written 416.7 *cm.* To make a metric rule, cut a piece of wood, paper, or tape, $39\frac{3}{8}$ in. long. Divide it into ten equal parts, and each part into ten *other* equal parts; each of these parts is 1 *centimetre*. Divide each centimetre into ten equal parts, and each part is a *millimetre*.

The diameter of the nickel five cent piece of 1866 is 2 *centimetres*, and its weight is 5 *grams*.

The Centimetre is the unit generally used for measure-

ments less than a metre. For its length in common measure see illustration.

The Metre is the unit commonly used by artisans. It equals 3 ft. 3 $\frac{3}{8}$ in. (nearly).

The Kilometre is the unit commonly used by surveyors in measuring distances. Its length is 198 rd. 13 ft. 10 in.

TABLE.*

<i>Full.</i>		<i>Contracted.</i>	
10 millimetres	= 1 centimetre.	10 millimetres	= 1 centimetre.
10 centimetres	= 1 decimetre.	100 centimetres	= 1 <i>metre</i> .
10 decimetres	= 1 <i>Metre</i> .	100 <i>metres</i>	= 1 kilometre.
10 metres	= 1 decametre.		
10 decametres	= 1 hectometre.		
10 hectometres	= 1 kilometre.		
10 kilometres	= 1 myriametre.		

SQUARE MEASURE.

The square Metre is the unit commonly used by artisans in specifying surfaces of small extent. It contains about 10 sq. ft. 110 sq. in.

The Are is the unit commonly used to express quantities less than the *hectare*. 100 ares make one hectare.

The Hectare is the unit commonly used by surveyors

* NOTE.—The unit of each table is divided into ten equal parts, designated by prefixing *deci* (tenth); as, *decigram*. The *tenths* are divided into ten other equal parts, designated by prefixing *centi* (hundredth); as, *centigram*. The *hundredths* are subdivided in the same manner, and are designated by prefixing *milli* (thousandth); as, *milligram*. The contracted table is the most convenient for common use.

in estimating the contents of land. It contains 2.471 acres.

TABLE.

<i>Full.</i>		<i>Contracted.</i>	
10 milliares	= 1 centiare.	100 sq. millimetres	= 1 sq. centimetre.
10 centiares	= 1 deciare.	100 sq. centimetres	= 1 sq. decimetre.
10 deciares	= 1 <i>Are</i> .	100 sq. decimetres	= 1 sq. <i>metre</i> .
10 ares	= 1 decare.	100 sq. metres	= 1 <i>are</i> .
10 decares	= 1 hectare.	100 ares	= 1 <i>hectare</i> .
10 hectares	= 1 kilare.		
10 kilares	= 1 myriare.		

CUBIC OR SOLID MEASURE.

The cubic Metre or Stere is the unit commonly used by engineers in estimating the solid contents of embankments, cellars, walls, &c. It equals 1.308 cu. yards.

TABLE.

<i>Full.</i>		<i>Contracted.</i>	
10 millisteres	= 1 centistere.	1000 cu. centimetres	= 1 litre.
10 centisteres	= 1 decistere.	1000 litres	= 1 stere.
10 decisteres	= 1 <i>Stere</i> .	1000 steres	= 1 kilostere.
10 steres	= 1 decastere.		
10 decasteres	= 1 hectostere.		
10 hectosteres	= 1 kilostere.		
10 kilosteres	= 1 myriastere.		

DRY AND LIQUID MEASURE.

The unit commonly used in the measurement of grain, roots, and liquids by the barrel is the *hectolitre*. It equals 26.417 gal. wine measure, or 2.839 bu. dry measure.

The unit commonly used by grocers is the *litre*. It equals *

1.056 qt. wine measure, or .908 qt. dry measure, or a trifle more than a wine quart.

TABLE.

<i>Full.</i>		<i>Contracted.</i>	
10 millilitres	= 1 centilitre.	100 centilitres	= 1 litre.
10 centilitres	= 1 decilitre.	100 litres	= 1 hectolitre.
10 decilitres	= 1 <i>Litre.</i>	1000 litres	= 1 kilolitre.
10 litres	= 1 decalitre.		
10 decalitres	= 1 hectolitre.		
10 hectolitres	= 1 kilolitre.		
10 kilolitres	= 1 myrialitre.		

WEIGHT.

The unit commonly used in philosophical experiments, by jewellers and druggists is the *gram*. Its weight is 15.432 gr. troy.

The unit commonly used by grocers is the *kilogram*, commonly contracted *kilo*. It is the weight of a litre of pure water, and equals 2.2046 lbs., or about $2\frac{1}{4}$ lbs. avoirdupois.

The unit commonly used in weighing heavy bodies, as coal, iron, marble, R. R. freight, &c., is the *tonneau*. It is the weight of a cubic metre of pure water, and equals 2204.6 lbs. avoirdupois.

TABLE.

<i>Full.</i>		<i>Contracted.</i>	
10 milligrams	= 1 centigram.	100 centigrams	= 1 gram.
10 centigrams	= 1 decigram.	1000 grams	= 1 kilogram.
10 decigrams	= 1 <i>Gram.</i>	1000 kilograms	= 1 tonneau.
10 grams	= 1 decagram.		
10 decagrams	= 1 hectogram.		
10 hectograms	= 1 kilogram.		
10 kilograms	= 1 myriagram.		
10 myriagrams	= 1 quintal.		
10 quintals	= 1 tonneau.		

MEASUREMENT OF ANGLES.

In the *centesimal* or French method the right angle is divided into 100 equal parts called *grades*, the grade into 100 equal parts called *minutes*, the minute into 100 equal parts called *seconds*.

TABLE.

100 seconds (")	=	1 minute (')
100 minutes	=	1 grade (gr.)
100 grades	=	1 right angle (r. a.)

NOTE.—Since the signs for both the common and centesimal methods are the same, to prevent confusion when minutes and seconds are expressed in the centesimal method, annex the abbreviation *cen.*; thus, 3' 46'' *cen.*

CURRENCY.

SCALE.

Franc
decimes
centimes
millimes
0. 0 0 0

TABLE.

10 millimes	=	1 centime.
10 centimes	=	1 decime.
10 decimes	=	1 <i>Franc</i> .

LINEAR MEASURE.

Table of equivalents.*

1 in. = 25 $\frac{1}{4}$ mm. (nearly).	1 mi. = 1609.35 m.
1 ft. = 305 mm. (nearly).	1 cm. = .3937 = $\frac{1}{2.54}$ in. (nearly).
1 yd. = 914 mm.	1 m. = 39.37 in. = 1.093 yd.
1 rd. = 5029 mm.	1 km. = .62137 mi. = 198 rd., 12 ft., 10 in.

* Authorized by Act of Congress, July 27, 1866.

SQUARE MEASURE.—*Table.*

1 sq. in. = 6.5 sq. cm.	1 sq. cm. = .155 sq. in.
1 sq. ft. = 9.3 sq. dm.	1 sq. m. = { 1550 sq. in.
1 sq. yd. = .835 sq. m.	{ 10.76 sq. ft.
1 acre = 40.47 a.	1 are. = 119.6 sq. yd.
	1 ha. = 2.471 acres.

CUBIC MEASURE.—*Table.*

1 cu. in. = 16.387 cu. centm.	1 litre = { 1.0567 qt. liq. meas.
1 cu. ft. = { 28.34 litres.	{ .908 qt. dry meas.
{ .0283 steres.	1 hecto-
1 cu. yd. = .76531 steres.	litre = { 2.837 bu. dry meas.
1 cord = 3.6281 steres.	{ 26.417 gal. liq. meas.
1 fluid oz. = .02958 litres.	1 kiloli-
1 gal. = 3.786 litres.	tre } { 35.316 cu. ft.
1 bus. = 35.24 litres.	1 cu. me- } = { 1.308 cu. yd.
	tre } { 264.17 gal. liq.
	1 stere } { meas.
	{ .2759 cord.

WEIGHT.—*Table.*

1 oz. troy = 31.1 grams.	1 ton avoird. = 907.2 kilos.
1 lb. troy } = 373.2 "	1 gram. = { 15.432 gr. troy.
1 lb. apoth. } = 373.2 "	{ .5643 dr. avoird.
1 oz. avoird. = 28.35 "	1 kilogram = 2.2046 lb. avoird.
1 lb. avoird. = 453.6 "	1 tonneau = 2204.6 lb. avoird.

ANGULAR MEASURE.—*Table.*

1 r. a. = 100 grades.	1 cir. = 400 grades.
1° = 1 $\frac{1}{4}$ grades.	1 grade = 9 deg.
1' = 1.85 minutes ('cen.).	1' cen. = 5.4'.
1" = 3.08 seconds ("cen.).	1" cen. = 3.24'.

SPECIFIC GRAVITY.

When a cubic foot of a substance is compared with the same bulk of water, and weighs a certain number of times as much, that number is called its *specific gravity*.

When any substance weighs less than water, it will float

on it, and when it weighs less than air, it will rise in it; thus, iron will float in melted lead, gas will rise in the air, and wood will float on water.


The weight of a cubic foot of water being 1000 ounces avoirdupois, it has been adopted as the standard of specific gravities. Hence the specific gravity of a body or substance is the proportion its weight bears to this standard.

To find the specific gravity of a body.

RULE.—Weigh it first in air and then in water, and take the difference of these weights; then as the difference is to the weight in air, so is 1000 to the specific gravity of the body.

EXAMPLE.—What is the specific gravity of a stone weighing 20 lbs., but in water only 15 lbs.?

SOLUTION.— $20 - 15 = 5$ difference; then $5 : 20 :: 1000 :$
4000. *Ans.*



When the body is lighter than water.

RULE.—Attach to it a piece of metal sufficient to sink it in the water; weigh the piece added and the body separately, both in and out of the water, and find how much each loses in water by subtracting its weight in water from its weight in air, and subtract the less of these differences from the greater; then as the remainder is to the weight of the light body in air, so is 1000 to the specific gravity of the body.

EXAMPLE.—Required the specific gravity of a piece of wood which weighs 20 lbs. in air; attached to it is a piece of metal, which weighs 30 lbs. in air and 25 lbs. in water, and the two pieces together weighing in water 10 lbs.?

$$\text{SOLUTION.}—20 + 30 - 10 = 40$$

$$30 - 25 = \quad 5$$

$$35 : 20 :: 1000 : 571.44. \text{ Ans.}$$

To reduce the specific gravity of a body to its weight in lbs. per cubic foot.

RULE.—Divide the specific gravity by 16, and the quotient is the weight of a cubic foot in lbs.

EXAMPLE.—Required the weight of a cubic foot of a substance the specific gravity of which is 4,800?

$$\text{SOLUTION.}—4,800 \div 16 = 300 \text{ lbs. } \text{Ans.}$$

TABLE, showing the specific gravities of various substances.

DESIGNATION.	Sp. Gravity.	DESIGNATION.	Sp. Gravity.	DESIGNATION.	Sp. Gravity
Antimony, . . .	6.712	Coral,	2.700	Cork,240
Arsenic,	5.763	Coal, bit,	1.270	Cypress,644
Bismuth,	9.823	“ anthr.,	1.556	Ebony,	1.331
Brass,	7.820	Diamond,	3.521	Elder,695
Bronze,	8.700	Earth, loose, . .	1.500	Elm,671
Copper,	8.788	Emery,	4.000	Fir, yellow,657
Copper wire, . .	8.878	Flint,	2.590	“ white,669
Gold, pure, . . .	19.258	Glass,	2.930	Lignum vitæ, . .	1.333
“ 22 carat. . .	17.486	Granite,	2.625	Live oak,	1.120
“ 20 carat . . .	15.709	Grindstone, . . .	2.143	Logwood,919
Iron, cast, . . .	7.207	Gypsum,	2.168	Mahogany,	1.063
“ bars,	7.778	Hone, white, . . .	2.876	Maple,750
Lead,	11.352	Ivory,	1.822	Mulberry,897
Mercury,	13.598	Limestone,	3.180	Orange,705
Platinum,	22.069	Lime, quick,804	Pine, yellow, . .	.660
Silver,	10.477	Manganese,	7.000	“ white,554
Steel,	7.833	Marble, par., . .	2.838	Pear,661
Tin,	7.291			Plum,785
Zinc,	6.861	DRY WOOD.		Quince,705
Alabaster,	2.730	Apple,793	Sassafras,482
Amber,	1.078	Adder,800	Walnut,671
Asbestos,	3.073	Ash,845	Willow,585
Borax,	1.714	Beech,852	Yew,798
Brick,	1.900	Box,	1.231	Hickory,838
Chalk,	2.784	Campeachy,913	Poplar,383
Charcoal,441	Cherry,715	“ white,529
Clay,	1.930	Cocoa,	1.040		

When the specific gravity of a substance is given, to find the weight of a cubic foot.

RULE.—Multiply the weight of a cubic foot of pure water (62½ lbs.) by the specific gravity of the given substance.

I wish to find the number of cubic inches in a piece of cast iron, that will displace 25 ounces of water. What will it weigh?

OPERATION.—1. 25 oz. \times 1728 = 43200.

2. $43200 \div 1000 = 43$ cu. in. (nearly). *Ans.*

3. 25 oz. $\times 450\frac{1}{2} \div 1000 = 11$ lb. (nearly).

Ans.

NOTE.—To find the number of cubic inches in any irregular body, weigh a vessel containing sufficient rain water to cover the solid, then immerse the solid in the water by means of a string or wire held in the hand, being careful not to touch the vessel. While the solid is immersed, weigh the water and vessel again; the difference will be the weight of the water displaced by the solid.

RULE.—I. Multiply the weight of the water in ounces by 1728, and divide by 1000, the result will be the contents in cubic inches.

II. To find the weight, multiply the weight of the water displaced in ounces by the weight of a cubic foot of the substance, and divide the product by 1000, and the result will be the weight in pounds.

I have a pattern of a lock that will displace 20 ounces of water; how much will 1000 copies of cast iron weigh? How much will they cost me at 9 cents per pound?

$$\text{OPERATION.}—20 \overset{\text{oz.}}{\times} 450\frac{1}{2} \div 1000 = 9.01 \text{ lb.}$$

$$9.01 \overset{\text{lb.}}{\times} 1000 \times 6.09 = \$810.90. \quad \text{Ans.}$$

I have a lead pattern of a wheel that displaces 15 ounces of water; what will 500 copies in brass cost me at 40 cents per pound?

$$\text{OPERATION.}—15 \overset{\text{oz.}}{\times} 504\frac{3}{4} \div 1000 = 7.571 \text{ lb.}$$

$$7.571 \overset{\text{lb.}}{\times} 500 \times \$.40 = \$1514.20. \quad \text{Ans.}$$

TABLE, showing the weight of a cubic foot of different substances.

				Avoir.
1 cubic foot of Brass.....	weighs	504 $\frac{1}{4}$	lb.	
“ “ Brick.....	“	125	“	
“ “ Copper.....	“	555	“	
“ “ Clay.....	“	135	“	
“ “ Coal (anthracite).....	“	54	“	
“ “ Coal (bituminous).....	“	50	“	
“ “ Granite.....	“	165	“	
“ “ Iron (wrought).....	“	486 $\frac{3}{4}$	“	
“ “ Iron (cast).....	“	450 $\frac{1}{2}$	“	
“ “ Lead.....	“	708 $\frac{1}{2}$	“	
“ “ Marble.....	“	171	“	
“ “ Soil (common).....	“	124	“	
“ “ Sand.....	“	95	“	
“ “ Tallow.....	“	59	“	
“ “ Water (pure).....	“	62 $\frac{1}{2}$	“	
“ “ Water (sea).....	“	64 $\frac{1}{3}$	“	
“ “ Wood (oak).....	“	55	“	
“ “ Wood (yellow pine).....	“	42	“	
“ “ Wood (white pine).....	“	30	“	
“ “ Charcoal (hard wood)....	“	18 $\frac{1}{2}$	“	
“ “ Charcoal (pine wood)....	“	18	“	
“ “ Cork.....	“	15	“	

VELOCITY.*

The average velocities of different objects are found in the following

* Parker's Philosophy.

188 SOLID MATTER AND WATER IN ARTICLES OF DIET.

TABLE.

	Per hour.		Per sec.
A man walks	3 miles,	or	4 feet.
A horse trots	7 "	or	10 "
A horse runs	20 "	or	29 "
Steamboat runs	18 "	or	26 "
Sailing vessel runs	10 "	or	14 "
Slow rivers flow	3 "	or	4 "
Rapid rivers flow	7 "	or	10 "
A moderate wind blows	7 "	or	10 "
A storm moves	36 "	or	52 "
A hurricane moves	80 "	or	117 "
A rifle ball "	1000 "	or	1466 "
Sound "	743 "	or	1142 "
Light "	192000 miles per sec.		
Electricity "	288000 " " "		

SOLID MATTER AND WATER IN ARTICLES OF DIET.

TABLE, *showing the proportion of solid matter and water in 100 parts each of the following articles of diet.*

Designation.	Solid matter.	Water.	Designation.	Solid matter.	Water.
Wheat	87	13	Pork	24	76
Peas	87	13	Codfish	21	79
Rice	86	14	Blood	20	80
Beans	86	14	Trout	19	81
Rye	86	14	Apples	18	82
Corn	86	14	Pears	16	84
Oatmeal	74	26	Carrots	13	87
Wheat bread	51	49	Beets	13	87
Mutton	29	71	Milk	13	87
Chicken	27	73	Oysters	13	87
Lean Beef	26	74	Cabbage	8	92
Eggs	26	74	Turnips	7	93
Veal	25	75	Water Melon	5	95
Potatoes	25	75	Cucumber	3	97

WEIGHTS OF GRAIN, SEEDS, &c.

TABLE, showing the weight of grain, seeds, &c., per bushel, as established by the Legislatures of the following States. The letter m indicates sold by measure.

ARTICLES.	New-York.	Ohio.	Pennsylvania.	Indiana.	Wisconsin.	Iowa.	Illinois.	Michigan.	Connecticut.	Massachusetts.	Rhode-Island.	Kentucky.	New-Jersey.	Vermont.	Missouri.	Canada.
Wheat, lbs.	60	60	60	60	60	60	60	60	56	60		60	60	60	60	60
Rye,	56	56	56	56	56	56	54	56	56	56		56	56	56	56	56
Corn,	58	56	56	56	56	56	56	56	56	56		56	56	56	52	56
Oats,	32	32	32	32	32	35	32	32	28	30		33	30	32	m	34
Barley,	48	48	47	48	48	48	44	48		46		48	48	46	m	48
Buckwheat,	48		48	50	42	52	40	42	45	46		52	50	46	m	48
Clover seed,	60	64		60	60	60		60				60	64		m	60
Timothy seed,	44	42		45		45		m		m		45			m	48
Flax seed,	55	56		56		56		m		m		56	55		m	56
Hemp seed,	44			44		44										
Blue-grass seed,	14			14		14										
Apples, dried,	22	25			28	24		28								22
Peaches, dried,	32	33			28	33		28								22
Coarse salt,	56	50	85	50		50				70		50			50	56
Fine salt,	56	50	62	50		50				70		50			50	56
Potatoes,	60			60		60			60	60	60			60		
Peas,	60									60						60
Beans,	62	56		60		60				60		60				60
Castor Beans,	46			46		46										
Onions,	57			57		57				50	50					
Corn Meal,				50							50					
Mineral Coal,				70												

To reduce cubic feet to bushels, struck measure, divide the cubic feet by 56 and multiply by 45.

PROPORTION OF ALCOHOL IN LIQUORS.

TABLE, showing the proportion of alcohol in 100 parts, each,
of the following liquors.

Designation.	Parts in 100.	Designation.	Parts in 100.
Scotch Whiskey.....	54.32	Sherry	19.17
Irish Whiskey.....	53.9	Claret.....	15.1
Rum.....	53.68	Champagne.....	13.8
Brandy.....	53.39	Gooseberry.....	11.84
Gin	51.6	Elder.....	8.79
Port.....	22.9	Ale.....	6.87
Madeira.....	22.27	Porter.....	4.2
Currant.....	20.55	Cider.....	9.8 to 5.2
Teneriffe.....	19.79	<i>Prof. Brande.</i>	

NUTRITIVE VALUE OF CERTAIN CROPS.

If we suppose an acre to yield the following quantities of the usually cultivated crops, the weight of dry starch and gum, of gluten, albumen, casein, &c., of oil or fat, and of saline matter, reaped in each crop, will be represented nearly by the following numbers:—

DESIGNATION.	Bushels.	lbs.	Bark or woody fibre.	Starch, sugar, and gum.	Gluten, albumen, and casein.	Oil.	Saline matter.
Wheat.....	25	1500	225	825	180	45	30
Barley.....	35	1800	270	1080	230	50	50
Oats.....	50	2100	420	1050	300	100	75
Peas.....	25	1600	130	800	380	54	48
Beans.....	25	1600	160	640	420	40	50
Indian Corn.....	30	1800	100	1260	220	130	80
Potatoes.....	12 tons	27000	1080	4800	540	45	240
Turnips.....	30 "	67000	1340	6000	1000	200	450
Wheat Straw.....	1½ "	3000	1500	900	40	80	150
Meadow Hay.....	1½ "	3400	1020	1360	240	120	220
Clover Hay.....	2 "	4500	1120	1800	420	200	400
Cabbage.....	20 "	45000	430	2300	1300	180	600

Johnston.

NOTE.—From the above table it appears that the acre which, by cropping with wheat, would yield a given weight of starch, sugar, and gum, would, when cropped with barley or oats, yield *one-fourth* more of these substances—with potatoes, about *four times* as much, and with turnips *eight times* the same quantity. In other words, the piece of ground which, when sown with wheat, will maintain one man, would support one and a quarter if sown with barley or oats, four with potatoes, and eight with turnips—in so far as the nutritive power of these crops depends on the starch, sugar, and gum they contain.

PERCENTAGE OF OIL IN SEEDS, GRAIN, &c.

Oil per cent. in different seeds, grain, &c.

Oil per cent.		Oil per cent.	
Linseed.....	11 to 22 say 17	Oats.....	5 to 8 say 6½
Hempseed.....	14 “ 25 “ 19	Indian corn.....	5 “ 9 “ 7
Rapeseed.....	40 “ 70 “ 55	Wheat bran.....	3 “ 5 “ 4
White mustard....	36 “ 38 “ 37	Potatoes, turnips,	
Sweet almond....	40 “ 54 “ 47	and cabbage.....	1½
Bitter almond.....	28 “ 46 “ 37	Wheat-straw.....	2 “ 3½ “ 3
Turnip seed.....	40 “ 50 “ 45	Oat-straw.....	4
Wheat flour.....	2 “ 4 “ 3	Meadow hay.....	2 “ 5 “ 3½
Barley.....	2 “ 3 “ 2½	Clover hay.....	3 “ 5 “ 5

QUANTITIES OF SEED REQUIRED TO THE ACRE, &c.

TABLE, showing the quantity of garden seeds required to plant a given space.

Designation.	Space and quantity of seeds.
Asparagus.....	1 oz. produces 1000 plants, and requires a bed 12 ft. sq.
“ Roots..	1000 plant a bed 4 feet wide 225 feet long.
Eng. Dwarf Beans	1 quart plants from 100 to 150 feet of row.
French “	1 “ “ 250 or 350 feet of row.
Beans, pole, large	1 “ “ 100 hills.
“ “ small	1 “ “ 300 hills, or 250 feet of row.

QUANTITY OF SEED REQUIRED.

Designation.	Space and quantity of seeds.
Beets.....	10 lbs. to the acre; 1 oz. plants 150 feet of row.
Broccoli and Kale	1 oz. plants 2,500 plants, and requires 40 sq. ft. of ground.
Cabbage.....	Early sorts same as brocoli, and require 60 sq. ft. ground.
Cauliflower.....	The same as cabbage.
Carrot.....	1 oz. to 150 of row.
Celery.....	1 oz. gives 7000 plants, and requires 8 sq. feet of ground.
Cucumber.....	1 oz. for 150 hills.
Cress.....	1 oz. sows a bed 16 feet square.
Egg Plant.....	1 oz. gives 2000 plants.
Endive.....	1 oz. gives 3000 plants, and requires 80 feet of ground.
Leek.....	1 oz. gives 2000 plants, and requires 60 feet of ground.
Lettuce.....	1 oz. " 7000 " and requires seed bed of 120 feet.
Melon.....	1 oz. for 120 hills.
Nasturtium.....	1 oz. sows 25 feet of row.
Onion.....	1 oz. " 200 " "
Okra.....	1 oz. " 200 " "
Parsley.....	1 oz. " 200 " "
Parsnip.....	1 oz. " 250 " "
Peppers.....	1 oz. gives 2500 plants.
Peas.....	1 quart sows 120 feet of row.
Pumpkin.....	1 oz. to 50 hills.
Radish.....	1 oz. to 100 feet.
Salsify.....	1 oz. to 150 feet of row.
Spinage.....	1 oz. to 200 feet of row.
Squash.....	1 oz. to 75 hills.
Tomato.....	1 oz. gives 2500 plants, requiring seed bed of 80 feet.
Turnip.....	1 oz. to 2000 feet.
Water Melon....	1 oz. to 50 hills.

TABLE, showing the quantity of seed required to the acre.

Designation.	Quantity of seed.	Designation.	Quantity of seed.
Wheat.....	1½ to 2 bush.	Broom Corn.....	1 to 1½ bush.
Barley.....	1½ to 2½ "	Potatoes.....	5 to 10 "
Oats.....	2 to 4 "	Timothy.....	12 to 24 quarts
Rye.....	1 to 2 "	Mustard.....	8 to 20 "
Buckwheat.....	¾ to 1½ "	Herd Grass.....	12 to 16 "
Millet.....	1 to 1½ "	Flat Turnip.....	2 to 3 lbs.
Corn.....	½ to 1 "	Red Clover.....	10 to 16 "
Beans.....	1 to 2 "	White Clover.....	3 to 4 "
Peas.....	2½ to 3½ "	Blue Grass.....	10 to 15 "
Hemp.....	1 to 1½ "	Orchard Grass.....	20 to 30 "
Flax.....	¾ to 2 "	Carrots.....	4 to 5 "
Rice.....	2 to 2½ "	Parsnips.....	6 to 8 "

TABLE, showing the quantity per acre when planted in rows or drills.

Broom Corn.....	1 to 1½ bush.	Onions.....	4 to 5 lbs.
Beans.....	1½ to 2 "	Carrots.....	2 to 2½ "
Peas.....	1½ to 2 "	Parsnips.....	4 to 5 "
		Beets.....	4 to 6 "

DEPTH OF SOWING WHEAT.

Wheat may be sowed too shallow as well as too deep. The depth must vary with the soil. A thinner covering is required in a close, thick, heavy soil, than in one light, gravelly, and sandy. Experiments made with wheat give the following results:—

Seeds sown to the depth of.....				Appeared above ground in	No. of plants that came up.
..... $\frac{1}{2}$ inch.				11 days.	$\frac{7}{8}$
"	"	"	"	12 "	all.
"	"	"	"	18 "	$\frac{7}{8}$
"	"	"	"	20 "	$\frac{6}{8}$
"	"	"	"	21 "	$\frac{3}{8}$
"	"	"	"	22 "	$\frac{1}{8}$
"	"	"	"	23 "	$\frac{1}{8}$

PROPORTIONS OF WEIGHT TO BULK.

TABLE, showing the weight per cubic foot of various substances, and the number of cubic feet required to make a ton of each.

Material.	Lbs. per cubic ft.	Cub. feet per ton.	Material.	Lbs. per cubic ft.	Cub. feet per ton.
METALS.			STONE, ETC.		
Cast Iron.....	454	4.93	Glass.....	180	12.44
Wrought Iron.....	485	4.62	Sand.....	95	23.56
Steel.....	490	4.6	Slate.....	167	13.4
Copper, cast.....	549	4.08	WOOD.		
Copper, wrought.....	557	4.02	Ash.....	48	46.
Brass.....	524	4.03	Beach.....	46	43.7
Lead.....	709	3.15	Cedar.....	35	64.
Silver.....	654		Elm.....	44	51.
Tin.....	456	4.9	Mahogany, Spanish.....	57	39.3
Gold.....	1203		Oak, English.....	53	43.
Zinc.....	439	5.	White Oak, American.....	45	49.
Platinum.....	1218		Live Oak.....	70	33.
Mercury.....	848	2.64	Pine, Pitch.....	43	51.6
White Lead.....	198	11.	" Yellow.....	38	59.
STONE, ETC.			" White.....	34	66.
Granite.....	165	13.5	Poplar.....	46	45.
Limestone.....	165	13.5	MISCELLANEOUS.		
Marble.....	171	13.1	Water, fresh.....	62.5	35.8
Paving Stone.....	151	14.8	" salt.....	64.5	34.8
Sand Stone.....	130	17.	Air*.....	.07529	
Brick.....	130	18.7	Steam†.....	.03689	
Chalk.....	174	12.8	Cork.....	15.	149.3
Clay.....	125	18.	Olive oil.....	57.	39.3
			Tallow.....	59.	

* At the level of the sea.

† Not under pressure.

CORN—PORK.

According to the Patent Office Reports, and the results of numerous experiments, 1 bushel of corn weighing 56 lbs. will produce $10\frac{1}{2}$ lbs. of pork. Throwing off $\frac{1}{2}$ to come at the net weight, gives $8\frac{2}{3}$ lbs. of pork as the product of 1 bushel of corn, or 1 lb. of pork as the product of $6\frac{2}{3}$ lbs. of corn. $3\frac{1}{2}$ lbs. of cooked corn-meal makes 1 lb. of pork. Assuming that it requires $6\frac{2}{3}$ lbs. of corn to make 1 lb. of pork (exclusive of the labor of feeding and taking care of hogs), the relation which the price of corn bears to that of pork is exhibited in the following

TABLE, showing the price of pork per lb. at different prices per bushel for corn.

Corn per bush. Cents.		Pork per pound. Cents.		Corn per bush. Cents.		Pork per pound. Cents.
12 $\frac{1}{2}$	1.50	38	4.52
15	1.78	40	4.76
17	2.	42	5.
20	2.38	45	5.35
22	2.62	50	5.95
25	2.96	55	6.54
30	3.57	60	7.14
33	3.92	65	7.74
35	4.	70	8.57

By reversing the above table we have the price of corn per bushel at different prices per lb. for pork. The use of the above table is obvious. For example, should corn be

selling for 50 cents per bushel and pork for only 5 cents per lb., it would be most profitable to sell the corn; but should corn be selling for 40 cents per bushel and pork for 6 cents per lb., it would be most profitable to reduce the corn to pork, and sell the latter.

To find the price of pork per lb., taking the price of corn per bushel as the datum.

RULE.—Divide the price of a bushel of corn by 8.40 (the number of lbs. of pork produced by a bushel of corn), and the quotient will be the answer.

EXAMPLE.—When corn is 20 cents per bushel, what should be the price of pork per lb.?

SOLUTION.—20.00 cents, \div 8.40 lbs., = 2.38 cents. *Ans.*

To find the price of corn per bushel, taking the price of pork per lb. as the datum.

RULE.—Multiply the price of a lb. of pork by 8.40 (the number of lbs. of pork produced by a bushel of corn), and the product will be the answer.

EXAMPLE.—What should be the price of corn per bushel when pork is selling at $4\frac{1}{2}$ cents per lb.

SOLUTION.—4.50 cents, \times 8.40 lbs., = 37.8 cents. *Ans.*

NOTE.—The foregoing table and rules must not be taken as *invariably* correct. It requires but little reflection to satisfy the farmer that the proportions and results exhibited by them must be influenced by many conditions and causes, such as the sample of corn used, the constitution and breed

as well as the age of the animal, its condition, powers of digestion, habits, health, &c. The very nature of the subject precludes the possibility of *exactly* defining the results and proportions. At best we can only have some *general, average* results and rules. The foregoing is deemed a safe general average.



LIFE AND INCREASE OF ANIMALS.



To keep hens in winter.

Provide—

1. A comfortable roost;
2. Plenty of sand, gravel and ashes, *dry*, to play in;
3. A box of lime;
4. Boiled meat, chopped fine, every two or three days;

5. Corn and oats, which will be best if boiled tender ;
6. All the crumbs and potato parings ;
7. Water, neither cold nor blood-warm.

This treatment has proved quite successful in a great many cases where the formula has been strictly adhered to, and hens which without it gave no eggs, with it immediately laid one each, on an average, every two days.

TABLE, *showing the period of reproduction and gestation of domestic animals.*

DESIGNATION.	Proper age for reproduction.	Period of the power of reproduction in years.	No. of Females for one Male.	PERIOD OF GESTATION AND INCUBATION		
				Shortest period, days.	Mean period, days.	Longest period, days.
Mare,.....	4 years.	10 to 12		322	347	419
Stallion.....	5 "	12 to 15	20 to 30			
Cow,.....	3 "	10 to 14		240	283	321
Bull,.....	3 "	8 to 10	30 to 40			
Ewe,.....	2 "	6		146	154	161
Ram,.....	2 "	7	40 to 50			
Sow,.....	1 "	6		109	115	143
Boar,.....	1 "	6	6 to 10			
She Goat,....	2 "	6		150	156	163
He Goat,....	2 "	5	20 to 40			
She Ass,.....	4 "	10 to 12		365	380	391
He Ass,.....	5 "	12 to 15				
She Buffalo, ..		8		281	308	335
Bitch,.....	2 "	8 to 9		55	60	63
Dog,.....	2 "	8 to 9				
She Cat,.....	1 "	5 to 6		48	50	56
He Cat,.....	1 "	9 to 10	5 to 6			
Doe Rabbit, ..	6 months	5 to 6		20	28	35
Buck Rabbit, ..	6 "	5 to 6	30			
Cock,.....	6 "	5 to 6	12 to 15			
Hen,.....		3 to 5		19	21	24
Turkey,.....				24	26	30
Duck,.....				28	30	32
Goose,.....				27	30	33
Pigeon,.....				16	18	20
Pea Hen,.....				25	28	30
Guinea Hen, ..				20	23	25
Swan,.....				40	42	45

Growth and life of animals.

	Man	grows	for 20 years, and lives	90 or 100 years.
The Camel	"	"	8 "	"
The Horse	"	"	5 "	"
The Ox	"	"	4 "	"
The Lion	"	"	4 "	"
The Dog	"	"	2 "	"
The Cat	"	"	1 1/2 "	"
The Hare	"	"	1 "	"
The Guinea pig	"	"	7 months, and lives	6 or 7 "

A TABLE showing at one view when *Forty Weeks* (the period of gestation in a cow) will expire, from any day throughout the year.

Jan.	Oct.	Feb.	Nov.	March	Dec.	April.	Jan.	May.	Feb.	June.	March.
1	8	1		1	6	1	6	1	5	1	8
2	9		9	2	7	2	7	2	6	2	9
3	10	2	10	3	8	3	8	3	7	3	10
4	11	3	11	4	9	4	9	4	8	4	11
5	12	4	12	5	10	5	10	5	9	5	12
6	13	5	13	6	11	6	11	6	10	6	13
7	14	6	14	7	12	7	12	7	11	7	14
8	15	7	15	8	13	8	13	8	12	8	15
9	16	8	16	9	14	9	14	9	13	9	16
10	17	9	17	10	15	10	15	10	14	10	17
11	18	10	18	11	16	11	16	11	15	11	18
12	19	11	19	12	17	12	17	12	16	12	19
13	20	12	20	13	18	13	18	13	17	13	20
14	21	13	21	14	19	14	19	14	18	14	21
15	22	14	22	15	20	15	20	15	19	15	22
16	23	15	23	16	21	16	21	16	20	16	23
17	24	16	24	17	22	17	22	17	21	17	24
18	25	17	25	18	23	18	23	18	22	18	25
19	26	18	26	19	24	19	24	19	23	19	26
20	27	19	27	20	25	20	25	20	24	20	27
21	28	20	28	21	26	21	26	21	25	21	28
22	29	21	29	22	27	22	27	22	26	22	29
23	30	22	30	23	28	23	28	23	27	23	30
24	31	23		24	29	24	29	24	28	24	31
	Nov.	24	Dec.	25	30	25	30	March		April.	
25	1	25	1	26	31	26	31	25	1	25	1
26	2	26	2		Jan.		Feb.	26	2	26	2
27	3	27	3	27	1	27	1	27	3	27	3
28	4	28	4	28	2	28	2	28	4	28	4
29	5	29	5	29	3	29	3	29	5	29	5
30	6			30	4	30	4	30	6	30	6
31	7			31	5			31	7	31	7

Table continued.

July.	April	Aug.	May.	Sept.	June.	Oct.	July.	Nov.	Aug.	Dec.	Sept
1	7	1	8	1	8	1	8	1	8	1	7
2	8	2	9	2	9	2	9	2	9	2	8
3	9	3	10	3	10	3	10	3	10	3	9
4	10	4	11	4	11	4	11	4	11	4	10
5	11	5	12	5	12	5	12	5	12	5	11
6	12	6	13	6	13	6	13	6	13	6	12
7	13	7	14	7	14	7	14	7	14	7	13
8	14	8	15	8	15	8	15	8	15	8	14
9	15	9	16	9	16	9	16	9	16	9	15
10	16	10	17	10	17	10	17	10	17	10	16
11	17	11	18	11	18	11	18	11	18	11	17
12	18	12	19	12	19	12	19	12	19	12	18
13	19	13	20	13	20	13	20	13	20	13	19
14	20	14	21	14	21	14	21	14	21	14	20
15	21	15	22	15	22	15	22	15	22	15	21
16	22	16	23	16	23	16	23	16	23	16	22
17	23	17	24	17	24	17	24	17	24	17	23
18	24	18	25	18	25	18	25	18	25	18	24
19	25	19	26	19	26	19	26	19	26	19	25
20	26	20	27	20	27	20	27	20	27	20	26
21	27	21	28	21	28	21	28	21	28	21	27
22	28	22	29	22	29	22	29	22	29	22	28
23	29	23	30	23	30	23	30	23	30	23	29
24	30	24	31		July.	24	31	24	31	24	30
	May.		June	24	1		Aug.		Sept.		Oct.
25	1	25	1	25	2	25	1	25	1	25	1
26	2	26	2	26	3	26	2	26	2	26	2
27	3	27	3	27	4	27	3	27	3	27	3
28	4	28	4	28	5	28	4	28	4	28	4
29	5	29	5	29	6	29	5	29	5	29	5
30	6	30	6	30	7	30	6	30	6	30	6
31	7	31	7			31	7			31	7

AGE OF ANIMALS.

To find the age of a horse.

The colt is born with 12 grinders. When 4 front teeth have made their appearance the colt is 12 days old, and when the next 4 appear it is four weeks old. When the corner teeth appear it is eight months old, and when the latter have attained the height of the front teeth it is a year old. The two year old colt has the kernel (the dark substance in the middle of the tooth's crown) ground or worn out of all the front teeth. In the third year the middle front teeth are being shifted, and when three years old these are substituted for the horse teeth. In the fourth year the next 4 are shifted, and in the fifth year the corner teeth are shifted. In the sixth year the kernel is worn out of the middle front teeth, and the bridle teeth have now attained their full growth. At seven years a hook has been formed on the corner teeth of the upper jaw: the kernel of the teeth next at the middle is worn out, and the bridle teeth begin to wear off. At eight years of age the kernel is worn out of all the lower front teeth, and begins to decrease in the middle upper fronts. In the ninth year the kernel has wholly disappeared from the upper middle front teeth, the hook on the corner teeth has increased in size, and the bridle teeth loose

their point. In the tenth year the kernel has worn out of the teeth next to the middle fronts of the upper jaw, and in the eleventh year the kernel has entirely disappeared from the corner teeth of the same jaw. At twelve years the crowns of all the front teeth in the lower jaw have become triangular, and the bridle teeth are much worn down. As the horse advances in age the gums shrink away from the teeth, which appear long and narrow, and the kernels become changed into darkish points. Gray hairs increase in the forehead and the chin becomes angular.

A modification of the foregoing, much more scientific or systematic, and probably quite as reliable, is the classification of Pessina, a distinguished veterinary surgeon of Germany.

Its principles may be distinctly understood by reference to the accompanying cuts, A, B, C, and D.

A, represents the corner tooth of a young horse; the oth-




FIG. A.

er nippers vary very little from this one in their construction and form.

The top of the tooth is long from side to side, and the extreme lower end is long from front to rear. The manner in

which the shape changes as we go farther down the tooth is represented in figure B, where cross sections at different sections are shown.




FIG. B.

The horse's tooth is worn away by use, and its upper surface assumes the form of these different sections consecutively, according to the extent to which it has been worn off. Of course, this only forms a general rule by which to judge of the age of a horse. Cribbiters, horses feeding chiefly on very old dry hay, and oats mixed with grit, and horses which are continually gnawing their mangers, will have their teeth worn away faster than will those which are fed on grass and moistened, cut, and ground feed, and which keep their teeth to themselves when they are not eating.

Pessina's table of indications of age is correct for the average of horses, and in all cases is sufficiently so for general purposes.

We quote the following from Herbert's hints to horse-keepers:—

“At *five* years the corners are 'up even with the other teeth; the mark is entirely worn out from the middle nippers, and partly worn from the next pair (fig. C).




FIG. C.

“At *six* years the mark is almost gone from the second pair, and the *outer edge* of the corner teeth is worn down.

“At *seven* years the mark is entirely gone from the second pair, and the edges of the corner teeth are worn somewhat flat.

“At *eight* years the teeth of the lower jaw are worn entirely flat, the mark having disappeared from all of them. The form of the surface of the tooth has become oval, and the central enamel is long from side to side, and is near to the front of the tooth.

“At *nine* years the middle nippers are rounded on the inner side, the oval of the second pair and of the corner teeth becomes broader, the central enamel is nearer to the inner side, and the marks have disappeared from the teeth of the upper jaw.

“At *ten* years the second pair are rounded on the inner side, and the central enamel is very near to the inner side.

"At *eleven* years the corner teeth are rounded, and the central enamel becomes very narrow.

"At *twelve* years the nippers are all rounded, and the central enamel has entirely disappeared from the lower jaw; but it may still be seen in the upper jaw.

"At *thirteen* years the middle nippers commence to assume a triangular form in the lower jaw, and the central enamel has entirely disappeared from the corner teeth of the upper jaw.

"At *fourteen* years the middle nippers have become triangular, and the second pair are assuming that form; the central enamel has diminished in the middle nippers of the upper jaw.

"At *fifteen* years the second pair have become triangular (fig. D); the central enamel is still visible in the upper jaw.




FIG. D.

"At *sixteen* years all of the teeth in the lower jaw have become triangular, and the central enamel is entirely removed from the second pair in the upper jaw.

"At *seventeen* years the sides of the triangle of the middle nippers are all of the same length; the central enamel has entirely disappeared from the upper teeth.

"At *eighteen* years the sides of the triangle of the middle nippers are longer at the sides of the teeth than in front.

"At *nineteen* years the middle nippers become flattened from side to side and long from front to rear.

"At *twenty* years the second pair assume the same form.

"At *twenty-one* years all of the teeth of the lower jaw have become flattened from side to side; the greatest diameter having become exactly the reverse of what it was in youth."

TO FIND THE AGE OF CATTLE.

In the *cow* the horn is often regarded as affording, by the number of its rings, a criterion of the animal's age. The horn of a heifer remains smooth or unprotuberant till the expiration of the second year of its life. A circle of thicker matter, or sort of horny button then begins to be formed, which is completed in another year; the next year this circle or button moves from the head, or is impelled by the cylindric growth of the horn, and another circle or button begins to be formed, which after another twelve-month is also impelled outward, and so on year after year of the whole life of the animal, so that by counting the number of rings on the cow's horns, and adding 2 to their number, its age is arrived at.

The rings on the *bull's* horns do not begin to appear until he is five years old, so that to arrive at his age we must add 5 to the number of rings. The horn of the ox is so very

strongly modified by his peculiar condition, as to be totally unlike that of the bull, the rings scarcely appearing at all.

The above rule would enable one to tell the age of the animal with unerring certainty, were the growth of the horns in each animal uniform and the rings distinct, which is not always the case; the rings often being confused and indistinct, and the growth of the horns varying in different animals. Besides, knavish cattle dealers often rasp off several of the rings of old and unsalable cows, and so smooth the rest of the horns as to make them look in keeping with their pretensions.

A safer rule is afforded by the teeth. At birth the two centre teeth (front) protrude through the gum; at the end of the second week the second pair appear; at the end of the third week the third pair, and at the end of the fourth week the fourth and last pair. The wearing of these teeth now constitutes the only guide for the next three months, at the expiration of which time all these (which are called the "milk teeth") begin to diminish in size and shrink away from each other, which process continues until the animal is two years old, when the new teeth begin to push out the slender remnants of the old and shrunken ones. At the end of the second year the first two permanent teeth appear in front; at three years the second pair are well up: at four the third pair, and at five years the fourth and last pair, have appeared, and the central pair are beginning to become worn down: at six years the last pair are full sized: at seven

years the dark line with bony boundery appears in all the teeth, and a broad circular mark appears within the central pair : at eight years this mark appears in all the teeth : at nine years a process of absorption and shrinkage, similar to that which reduced the first teeth, begins to take place in the central pair ; at ten it begins with the second pair ; at eleven with the third pair, at twelve with the fourth pair. The age of the animal, after this period is attained, is determined by the degree of shrinkage and wearing away of all the teeth in the order of their appearance, until the fifteenth year, when scarcely any teeth remain.


To ascertain the age of sheep.

The age of sheep may be known by the front teeth, which are 8 in number, and appear the first year all of a size. In the second year the two middle ones fall out and are supplanted by two large ones. During the third year a small tooth appears on each side. In the fourth year the large teeth are six in number. In the fifth year all the front teeth are large, and in the sixth year the whole begin to get worn.

To tell the age of goats.

The age of goats is ascertained by their teeth in the same manner that of the sheep is, and by the annular rings on their horns.

COMPUTE WEIGHT OF CATTLE.



For cattle of a girth of from 5 to 7 feet, allow 23 lbs. to the superficial foot.

For cattle of a girth of from 7 to 9 feet, allow 31 lbs. to the superficial foot.

For small cattle and calves of a girth of from 3 to 5 feet, allow 16 lbs. to the superficial foot.

For pigs, sheep, and all cattle measuring less than 3 feet girth, allow 11 lbs. to the superficial foot.

RULE.—Ascertain the girth in inches back of the shoulders, and the length in inches from the square of the buttock to a point even with the point of the shoulder-blade. Multiply the girth by the length, and divide the product by 144 for the superficial feet, and then multiply the superficial feet by

the number of lbs. allowed as above for cattle of different girths, and the product will be the number of lbs. of beef, veal, or pork in the four quarters of the animal. To find the number of stone divide the number of lbs. by 14.

EXAMPLE.—What is the computed weight of beef in a steer, whose girth is 6 feet 4 inches, and length 5 feet 3 inches?

SOLUTION.—76 inches, girth, \times 63 inches, length, $=4788 \div 144=33\frac{1}{4}$ square feet, $\times 23=764\frac{3}{4}$ lbs., or $54\frac{5}{8}$ stone. *Ans.*

NOTE.—When the animal is but half fattened a deduction of 14 lbs. in every 280, or one stone in every 20 must be made; and if very fat, one stone for every 20 must be added.


Where great numbers of cattle are annually bought and sold under circumstances that forbid ascertaining their weight with positive accuracy, the compute weight may be thus taken with approximate exactness—at least with as much accuracy as is necessary in the aggregate valuation of stock. No rules or tables can, however, be at all times implicitly relied on, as there are many circumstances connected with the build of the animal, the mode of fattening, its condition, breed, &c., that will influence the measurement, and consequently the weight. A person skilled in taking the compute weight of stock soon learns, however, to make allowances for all these circumstances.

The following table is compiled from two English works on the subject:—

Girth. ft. in.	Length. ft. in.	Renton's Table. stone. lb.	Cary's Table. stone. lb.
5 0	3 6	21 0	21 00
5 0	4 0	24 0	24 00
5 0	3 9	27 1	27 00
5 6	4 0	34 4	34 07
6 0	4 6	38 8	38 11
6 0	5 0	43 1	43 00
6 6	4 6	45 9	45 07
6 6	4 9	48 0	48 00
7 0	5 6	64 6	64 07
7 0	6 0	70 5	70 03
8 0	6 6	99 8	99 12
8 0	7 0	107 5	107 06



FOOD OF ANIMALS.



TABLE, showing the comparative difference between good hay and the substances mentioned below, as food for stock—being the results of experiments.

10 lbs. of hay are equal to		10 lbs. of hay are equal to	
8 to 10 lbs.	clover hay.	30 to 35 lbs.	mangold wurtzel.
45 to 50 "	green clover.	45 to 50 "	turnips.
40 to 50 "	wheat straw.	20 to 30 "	cabbage.
20 to 40 "	barley straw.	3 to 5 "	peas and beans.
20 to 40 "	oat straw.	5 to 6 "	wheat.
10 to 15 "	pea straw.	5 to 6 "	barley.
20 to 25 "	potatoes.	4 to 7 "	oats.
25 to 30 "	carrots (red).	5 to 7 "	Indian corn.
40 to 45 "	" (white).	2 to 4 "	oil cake.

NOTE.—In the use of the above table much of course will depend upon the quality of the sample, the age and constitution of the animal, and the form in which the food is administered. Much also depends upon a *change of food*, and a *due admixture* of the different kinds.

TABLE, *showing the comparative difference between good hay and the articles mentioned below, as food for stock—being the mean of experiment and theory.*

100 lbs. of hay are equal to	100 lbs. of hay are equal to
275 lbs. green Indian corn.	54 lbs. rye.
442 " rye straw.	46 " wheat.
360 " wheat "	59 " oats.
164 " oats "	45 " peas and beans mixed.
180 " barley "	64 " buckwheat.
153 " pea "	57 " Indian corn.
200 " buckwheat straw.	63 " acorns.
201 " raw potatoes.	105 " wheat bran.
175 " boiled "	109 " rye "
339 " mangold wurtzel.	167 " wheat, pea, and oat chaff.
504 " turnips.	179 " rye and barley, mixed.
300 " carrots.	

NOTE.—It must be borne in mind that the nutritive effects of food upon the animal are varied by numberless causes, such as the animal's power of digestion and appropriation, its condition, shelter, air, water, exercise, &c. But all else being equal, the nutritive qualities of the articles mentioned are in the above proportions.

The results of numerous experiments, reported by individuals and Agricultural Associations, show, that each 100 lbs. of live weight of the animal requires of hay or its equivalent, per day, as follows:—

Working horses.....	3.08 lbs.
" oxen.....	2.40 "

Fatting oxen.....	5.00 lbs.
" " when fat.....	4.00 "
Milch cows.....	from 2.25 to 2.40 "
Dry ".....	2.42 "
Young growing cattle.....	3.08 "
Steers.....	2.84 "
Pigs.....	3.00 "
Sheep.....	3.00 "
Elephant*.....	3.12 "

In the ox, the daily loss of muscle or tissue requires that he should consume 20 to 24 ounces of gluten or albumen, which will be supplied by any of the following weights of vegetable food :—

Meadow hay.....	20 lbs	Turnips.....	120 lbs.
Clover hay.....	16 "	Cabbage.....	70 "
Oat straw.....	110 "	Wheat or other white grain.....	11 "
Pea straw.....	12 "	Beans or peas.....	6 "
Potatoes.....	60 "	Oil cake.....	4 "
Carrots.....	70 "		

Or instead of any one of these, a mixture of several may be given with the best results. But if the due proportion of nitrogenous food be not given, the ox will lose his muscular strength and will generally fail. So with growing and fattening stock of every description; the proportion of each of the kinds of food required by the animal must, in practice, be adjusted to the purpose for which it is fed.

It is not strictly correct that this or that kind of vegetable is more fitted to sustain animal life simply because of the large proportion of *nitrogen* or *gluten* it contains; it is wisely provided, however, that, along with this nitrogen, all

* Mr. Barnum's elephant, weighing 4700 lbs., was found to consume 100 lbs. of hay and 1 bushel of oats per day.

plants contain a certain proportion of starch or sugar, and of saline or earthy matter—all of which are required in a mixture which will most easily sustain an animal in a healthy condition ; so that the proportion of nitrogen in a substance may be considered as a rough practical index of the proportion of the more important saline and earthy ingredients also.

TABLE, *showing the effects produced by an equal quantity of the following substances, as food for sheep.*

Lbs.	Designation.	Increased weight of living animal in Lbs.	Produced Wool. Lbs.	Produced Tallow. Lbs.
1000	potatoes, raw with salt.....	46½	6½	12½
"	" " without salt.....	44	6½	11½
"	mangel-wurtzel, raw.....	38½	5½	6½
"	wheat.....	155	14	59½
"	oats.....	146	10	42½
"	barley.....	136	11½	60
"	peas.....	134	14½	41
"	rye, with salt.....	133	14	35
"	" without salt.....	90	12	43
"	corn meal, wet.....	129	13½	17½
"	buckwheat.....	120	10	33

NOTE.—The above are the results of numerous experiments by De Raumer.



DECREASE AND EXPECTATION OF LIFE.

TABLE, showing the decrement and expectation of human life.

Age.	Persons Liv- ing.	Decrement of Life.	Expectation of Life in Years and Decimals.	Age.	Persons Liv- ing.	Decrement of Life.	Expectation of Life in Years and Decimals.	Age.	Persons Liv- ing.	Decrement of Life.	Expectation of Life in Years and Decimals.
At birth.	4893	1264	28.1	34	1772	38	30.24	68	772	37	12.43
1	3629	174	"	35	1737	35	23.22	69	735	37	"
2	3355	188	"	36	1702	35	"	70	698	37	10.03
3	3167	132	"	37	1667	35	"	71	601	37	"
4	3035	84	"	38	1632	35	"	72	624	37	"
5	2951	58	40.87	39	1597	35	"	73	567	37	"
6	2893	55	"	40	1562	35	26.04	74	549	37	"
7	2838	47	"	41	1527	35	"	75	511	37	7.83
8	2791	40	"	42	1492	35	"	76	474	37	"
9	2751	36	"	43	1457	35	"	77	437	37	"
10	2715	28	39.27	44	1423	34	"	78	400	37	"
11	2687	27	"	45	1396	27	23.92	79	363	37	"
12	2660	27	"	46	1369	27	"	80	326	35	5.85
13	2633	27	"	47	1342	27	"	81	291	34	"
14	2606	27	"	48	1315	27	"	82	257	34	"
15	2579	42	36.10	49	1288	27	"	83	223	34	"
16	2557	43	"	50	1258	27	21.16	84	193	34	"
17	2494	43	"	51	1261	27	"	85	155	21	4.57
18	2451	43	"	52	1234	27	"	86	134	21	"
19	2408	43	"	53	1207	27	"	87	113	21	"
20	2365	43	34.21	54	1180	27	"	88	92	20	"
21	2322	42	"	55	1153	27	18.25	89	72	20	"
22	2280	42	"	56	1126	27	"	90	52	8	3.73
23	2238	42	"	57	1099	27	"	91	44	7	"
24	2196	42	"	58	1072	27	"	92	37	7	"
25	2154	40	32.32	59	1045	27	"	93	30	7	"
26	2114	38	"	60	1018	27	15.43	94	23	7	"
27	2076	38	"	61	991	27	"	95	16	6	1.62
28	2038	38	"	62	964	27	"	96	10	5	"
29	2000	38	"	63	937	27	"	97	5	3	"
30	1962	38	30.24	64	910	27	"	98	2	1	"
31	1924	38	"	65	883	37	12.43	99	1	1	"
32	1886	38	"	66	846	37	"				
33	1848	38	"	67	809	37	"				

The above table, originally compiled by Dr. Wigglesworth, of New England, after many years of careful observation and statistical research, exhibits the average yearly decrease

of life out of a given number born, and the expectation of reaching a certain age deduced from that decrease as the *datum*. Among the many similar tables that have been constructed, it is perhaps the most accurate. It received the cautious scrutiny and revision of the Supreme Court of Massachusetts, and was adopted by it (*see Easterbrook vs. Hopgood*, 10 *Mass. Reports*, 313) as the rule in estimating the value of life estates.

EXPLANATION.—Opposite the age of the individual, under the column headed “Expectation of Life, &c.,” will be found the additional number of years he may reasonably expect to live. Thus a man 40 years of age may reasonably expect to live 26.04 years longer.

For the purpose of comparison with observations in Europe, St. Maur’s Table is subjoined, taken from observations in Paris and the country around it.

ST. MAUR’S TABLE.

Of 24,000 born

17,540	attain to	2 years.	9,544	attain to	30 years.
15,162	“	“ 3 “	8,770	“	“ 35 “
14,177	“	“ 4 “	7,729	“	“ 40 “
13,477	“	“ 5 “	7,008	“	“ 45 “
12,968	“	“ 6 “	6,197	“	“ 50 “
12,562	“	“ 7 “	5,375	“	“ 55 “
12,255	“	“ 8 “	4,564	“	“ 60 “
12,015	“	“ 9 “	3,450	“	“ 65 “
11,861	“	“ 10 “	2,544	“	“ 70 “
11,405	“	“ 15 “	1,507	“	“ 75 “
10,909	“	“ 20 “	807	“	“ 80 “
10,259	“	“ 25 “	291	“	“ 85 “

103 attain to 90 years.	23 attain to 96 years.
71 " " 91 "	18 " " 97 "
63 " " 92 "	16 " " 98 "
47 " " 93 "	8 " " 99 "
40 " " 94 "	6 or 7 " " 100 "
33 " " 95 "	

EXPLANATION.—To ascertain by the above table what probability there is that a man of a given age will attain to any other age, make the number opposite the latter age the numerator and the number opposite the former age the denominator, and the fraction will express the probability sought for.

EXAMPLE.—What probability is there that a man of 30 will attain the age of 70 years?

SOLUTION.—Opposite 70 find $2,544 = 318$

" 30 " $9,544 = 1193$ Ans. That is to say, he has 318 chances out of 1193 of living to 70.

COMPOUND INTEREST.

TABLE, showing the amount of \$1 for any number of years from 1 to 24, at 5 and 6 per cent., compound interest.

Years.	5 per cent.	6 per cent.	Years.	5 per cent.	6 per cent.
1	1.05	1.06	13	1.88564	2.13292
2	1.1025	1.1236	14	1.97993	2.26090
3	1.15763	1.19101	15	1.07892	2.39655
4	1.21550	1.26247	16	1.18287	2.54035
5	1.27628	1.33822	17	1.29201	2.69277
6	1.34009	1.41851	18	1.41661	2.85433
7	1.40710	1.50363	19	1.52695	3.02559
8	1.47745	1.59384	20	1.6329	3.20713
9	1.55132	1.68947	21	1.78596	3.39956
10	1.62889	1.79084	22	1.92526	3.60353
11	1.71033	1.89829	23	1.07152	3.81974
12	1.79585	2.01219	24	1.22509	4.04893

EXPLANATION.—Opposite the number of years in the column under the rate per cent., will be found the amount of \$1, with the compound interest included for the time given. Should the amount of any given sum with the compound interest at a given rate per cent. for a given time be required, multiply the amount found in the column under the given rate per cent., and opposite the given time, by the sum at interest so given, and the product will be the answer.

EXAMPLE.—What will be the amount of \$150 at compound interest at the rate of 5 per cent. for 10 years?

SOLUTION.— $1.62889 \times 150 = \$244.33.35$. *Ans.*

ANNUITIES.

TABLE, *showing the present worth of \$1 annuity at 5 and 6 per cent. compound interest for any number of years from 1 to 34.*

Year.	5 per cent.	6 per cent.	Years.	5 per cent.	6 per cent.
1	0.95238	0.94339	18	11.68958	10.82760
2	1.87941	1.83339	19	12.08532	11.15711
3	2.72325	2.67401	20	12.46221	11.46992
4	3.5495	3.46510	21	12.82115	11.76407
5	4.3948	4.21236	22	13.16300	12.04158
6	5.07569	4.91732	23	13.48807	12.30338
7	5.70037	5.58438	24	13.79864	12.55035
8	6.46821	6.20979	25	14.09394	12.78335
9	7.10782	6.80169	26	14.37518	13.00316
10	7.72173	7.3608	27	14.64303	13.21053
11	8.30641	7.88687	28	14.89813	13.40616
12	8.86325	8.38384	29	15.14107	13.59072
13	9.39257	8.8568	30	15.37245	13.76483
14	9.89864	9.29493	31	15.59281	13.92908
15	10.37966	9.71225	32	15.80268	14.08398
16	10.83777	10.10589	33	16.00255	14.22917
17	11.27407	10.47726	34	16.19290	14.36613

For explanation and example see Compound Interest above.

INTEREST TABLE.

For convenience in finding the interest from one to \$5000 from one to 2000 days, at 6 or 7 per cent., we give the following

TABLE.*

\$	1da.	2	3	4	5	6	7	8	9	10	20	30	40	50	60	70	80	90	100	200
1 {	6 per ct.	.000	.001	.001	.001	.001	.001	.001	.001	.002	.003	.005	.007	.008	.010	.012	.013	.015	.017	.038
7 per ct.	.001	.001	.001	.001	.001	.001	.001	.002	.002	.002	.004	.006	.008	.010	.012	.014	.016	.017	.019	.039
2 {	.000	.001	.001	.001	.002	.002	.002	.003	.003	.003	.007	.011	.013	.017	.020	.023	.027	.030	.033	.067
.000	.001	.001	.002	.002	.002	.002	.003	.003	.003	.004	.008	.012	.016	.019	.023	.027	.031	.035	.039	.078
3 {	.000	.001	.001	.002	.002	.003	.003	.004	.004	.005	.010	.015	.020	.025	.030	.035	.040	.045	.050	.100
.001	.001	.002	.002	.003	.003	.004	.005	.005	.006	.006	.012	.017	.023	.029	.035	.041	.047	.052	.058	.117
4 {	.001	.001	.002	.003	.003	.004	.005	.005	.006	.007	.013	.021	.027	.033	.040	.047	.053	.060	.067	.133
.001	.002	.002	.003	.004	.005	.005	.006	.006	.007	.008	.016	.023	.031	.039	.047	.054	.062	.070	.078	.156
5 {	.001	.002	.003	.004	.005	.006	.007	.007	.008	.009	.017	.025	.033	.042	.050	.058	.067	.075	.083	.167
.001	.002	.003	.004	.005	.006	.007	.008	.009	.010	.011	.019	.029	.038	.049	.058	.068	.078	.087	.097	.194
6 {	.001	.002	.003	.004	.005	.006	.007	.008	.009	.010	.020	.030	.040	.050	.060	.070	.080	.090	.100	.200
.001	.002	.003	.005	.006	.007	.008	.009	.010	.012	.012	.023	.035	.047	.058	.070	.082	.093	.105	.117	.228
7 {	.001	.002	.003	.005	.006	.007	.008	.009	.010	.012	.023	.035	.047	.058	.070	.082	.093	.105	.117	.228
.001	.003	.004	.006	.007	.008	.009	.010	.011	.012	.014	.027	.041	.054	.068	.082	.095	.109	.122	.136	.272
8 {	.001	.003	.004	.006	.007	.008	.009	.011	.012	.013	.027	.040	.053	.067	.080	.093	.107	.120	.133	.267
.002	.003	.005	.006	.008	.009	.011	.012	.014	.016	.016	.031	.047	.062	.078	.093	.109	.124	.140	.156	.311
9 {	.001	.003	.004	.006	.007	.009	.010	.012	.013	.015	.030	.045	.060	.075	.090	.105	.120	.135	.150	.300
.002	.003	.005	.007	.009	.010	.012	.014	.016	.017	.017	.035	.052	.070	.087	.105	.122	.140	.157	.175	.351
10 {	.002	.003	.005	.007	.008	.010	.012	.013	.015	.017	.033	.050	.067	.083	.100	.117	.133	.150	.167	.333
.002	.004	.006	.008	.010	.012	.014	.016	.017	.019	.019	.039	.058	.078	.097	.117	.135	.156	.175	.194	.389
20 {	.003	.007	.010	.013	.017	.020	.023	.027	.030	.033	.067	.100	.133	.167	.200	.233	.267	.300	.333	.667
.004	.008	.012	.016	.019	.023	.027	.031	.035	.039	.048	.078	.117	.156	.194	.233	.272	.311	.350	.389	.778

INTEREST TABLE.

221

30	.005	.010	.015	.020	.025	.030	.035	.040	.045	.050	.100	.150	.200	.250	.300	.350	.400	.450	.500	1.000
	.006	.012	.017	.023	.029	.035	.041	.047	.052	.058	.117	.175	.233	.292	.350	.408	.467	.525	.583	1.167
40	.007	.013	.020	.027	.033	.040	.047	.053	.060	.067	.133	.200	.267	.333	.400	.467	.533	.600	.667	1.333
	.008	.016	.023	.031	.039	.047	.054	.062	.070	.078	.156	.233	.311	.389	.467	.544	.622	.700	.778	1.556
50	.008	.017	.025	.033	.042	.050	.058	.067	.075	.083	.167	.250	.333	.417	.500	.583	.667	.750	.833	1.667
	.010	.019	.029	.039	.049	.058	.068	.078	.087	.097	.194	.292	.389	.486	.583	.681	.778	.875	.972	1.944
60	.010	.020	.030	.040	.050	.060	.070	.080	.090	.100	.200	.300	.400	.500	.600	.700	.800	.900	1.000	2.000
	.012	.025	.035	.047	.058	.070	.082	.093	.105	.117	.233	.350	.467	.583	.700	.817	.933	1.050	1.167	2.333
70	.012	.023	.035	.047	.058	.070	.082	.093	.105	.117	.233	.350	.467	.583	.700	.817	.933	1.050	1.167	2.333
	.014	.027	.041	.054	.068	.082	.095	.109	.122	.136	.272	.408	.544	.681	.817	.953	1.089	1.225	1.361	2.722
80	.013	.027	.040	.053	.067	.080	.093	.107	.120	.133	.267	.400	.533	.667	.800	.933	1.067	1.200	1.333	2.667
	.016	.031	.047	.062	.078	.093	.109	.124	.140	.156	.311	.467	.622	.778	.933	1.089	1.244	1.400	1.556	3.111
90	.015	.030	.045	.060	.075	.091	.105	.120	.135	.150	.300	.450	.600	.750	.900	1.050	1.200	1.350	1.500	3.000
	.017	.035	.052	.070	.087	.105	.122	.140	.157	.175	.350	.525	.700	.875	1.050	1.225	1.400	1.575	1.750	3.500
100	.017	.033	.050	.067	.083	.100	.117	.133	.150	.167	.333	.500	.667	.833	1.000	1.167	1.333	1.500	1.667	3.333
	.019	.039	.058	.078	.097	.117	.136	.156	.175	.194	.389	.583	.778	.972	1.167	1.361	1.556	1.750	1.944	3.889
200	.033	.067	.100	.133	.167	.200	.233	.267	.300	.333	.667	1.000	1.333	1.667	2.000	2.333	2.667	3.000	3.333	6.644
	.039	.078	.117	.156	.194	.233	.272	.311	.350	.389	.778	1.167	1.556	1.944	2.333	2.722	3.111	3.500	3.889	7.777
300	.050	.100	.150	.200	.250	.300	.350	.400	.450	.500	1.000	1.500	2.000	2.500	3.000	3.500	4.000	4.500	5.000	10.000
	.058	.117	.175	.233	.292	.351	.408	.467	.525	.583	1.167	1.750	2.333	2.917	3.500	4.083	4.667	5.250	5.833	11.666
400	.067	.133	.200	.267	.333	.400	.467	.533	.600	.667	1.333	2.000	2.667	3.333	4.000	4.667	5.333	6.000	6.667	13.333
	.078	.156	.233	.311	.389	.467	.544	.622	.700	.778	1.556	2.333	3.111	3.889	4.667	5.444	6.222	7.000	7.777	15.555
500	.083	.167	.250	.333	.417	.500	.583	.667	.750	.833	1.667	2.500	3.333	4.167	5.000	5.833	6.667	7.500	8.333	16.667
	.097	.194	.292	.389	.486	.583	.681	.778	.875	.972	1.944	2.917	3.889	4.861	5.833	6.805	7.778	8.750	9.722	19.444

EXPLANATION.—To find from the above table the interest of a given amount for a given number of days, refer to the square opposite the number of dollars at the left, and the number of days at the top. For \$5000 remove the decimal point one place to the right.

* NOTES.—1. On account of space the table is condensed as much as possible; but in practice any interest table may be used, so that the interest on any amount for any number of days may be taken at a glance, and the calculation may be greatly facilitated. This table is prepared at \$30 d. per year.

2. Accounts may be averaged by any per cent., but the cash balance can only be found by using the legal rate per cent.

TABLE OF SIMPLE INTEREST, AT SEVEN PER CENT.,

For each Day to a Month, from \$1 to \$100.

Amount	DAYS IN THE MONTH.																							
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
40	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
50	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
60	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
70	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
80	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
90	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
100	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

EXPLANATION.—Find the amount in the left hand column then follow the line of figures until you come to the column giving the number of days at the head and you have the amount of days to one month; then by adding together, you have the interest of as many months as required, on amounts from one dollar to one hundred.

TABLE OF SIMPLE INTEREST, AT SIX PER CENT.,

For each Day to a Month, from \$1 to \$100.

Amount.	DAYS IN THE MONTH.																														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
7	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
20	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
30	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
40	1	1	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2
50	1	1	2	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3	3
60	1	1	2	3	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
70	1	1	2	3	4	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
80	1	1	2	3	4	5	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6
90	2	2	3	4	5	6	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7
100	2	2	3	4	5	6	7	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8

EXPLANATION.—Find the amount in the left hand column then follow the line of figures until you come to the column giving the number of days at the head and you have the amount of days to one month; then by adding together, you have the interest of as many months as required on amounts from one dollar to one hundred.

TABLE OF WAGES AT \$3.00 to \$25.00 PER MONTH OF 26 WORKING DAYS,

For any number of days from 1 to 26.

Days	\$3.	\$4.	\$5.	\$6.	\$7.	\$8.	\$9.	\$10.	\$11.	\$12.	\$13.	\$14.	\$15.	\$16.	\$17.
1	.12	.15	.19	.23	.27	.31	.35	.38	.42	.46	.50	.54	.58	.62	.66
2	.23	.31	.39	.46	.54	.62	.69	.77	.85	.92	1.00	1.08	1.15	1.23	1.31
3	.35	.46	.57	.69	.81	.92	1.04	1.15	1.27	1.38	1.50	1.62	1.73	1.85	1.96
4	.46	.62	.77	.93	1.07	1.23	1.38	1.54	1.69	1.85	2.00	2.16	2.31	2.46	2.62
5	.58	.77	.96	1.15	1.34	1.54	1.73	1.92	2.12	2.31	2.50	2.69	2.88	3.08	3.27
6	.69	.83	1.11	1.38	1.62	1.85	2.08	2.31	2.54	2.77	3.00	3.23	3.46	3.69	3.92
7	.81	1.07	1.31	1.62	1.88	2.15	2.42	2.69	2.96	3.23	3.50	3.77	4.04	4.31	4.58
8	.92	1.21	1.54	1.84	2.15	2.46	2.77	3.08	3.38	3.69	4.00	4.31	4.62	4.92	5.23
9	1.04	1.38	1.73	2.07	2.42	2.77	3.12	3.46	3.81	4.15	4.50	4.85	5.19	5.54	5.88
10	1.15	1.54	1.92	2.31	2.69	3.08	3.46	3.85	4.23	4.62	5.00	5.38	5.77	6.15	6.54
11	1.27	1.69	2.12	2.54	2.96	3.38	3.81	4.23	4.65	5.08	5.50	5.92	6.35	6.77	7.19
12	1.38	1.84	2.31	2.77	3.23	3.69	4.15	4.62	5.08	5.54	6.00	6.46	6.92	7.38	7.85
13	1.50	2.00	2.50	3.00	3.50	4.00	4.50	5.00	5.50	6.00	6.50	7.00	7.50	8.00	8.50
14	1.62	2.15	2.69	3.23	3.77	4.31	4.85	5.38	5.92	6.46	7.00	7.54	8.08	8.62	9.15
15	1.73	2.31	2.88	3.46	4.04	4.62	5.19	5.77	6.35	6.92	7.50	8.08	8.62	9.23	9.81
16	1.85	2.46	3.08	3.69	4.31	4.92	5.54	6.16	6.77	7.38	8.00	8.62	9.23	9.85	10.46
17	1.96	2.61	3.27	3.92	4.58	5.23	5.88	6.54	7.19	7.85	8.50	9.15	9.81	10.46	11.12
18	2.08	2.77	3.46	4.15	4.84	5.51	6.23	6.92	7.62	8.31	9.00	9.69	10.38	11.08	11.77
19	2.19	2.93	3.62	4.38	5.11	5.85	6.58	7.31	8.04	8.77	9.50	10.23	10.96	11.69	12.42
20	2.31	3.07	3.81	4.61	5.38	6.16	6.92	7.69	8.46	9.23	10.00	10.77	11.54	12.31	13.08
21	2.42	3.23	4.04	4.84	5.65	6.46	7.27	8.08	8.88	9.69	10.50	11.31	12.12	12.92	13.73
22	2.54	3.38	4.23	5.07	5.92	6.77	7.61	8.46	9.31	10.15	1.00	11.85	12.69	13.54	14.38
23	2.65	3.54	4.42	5.31	6.19	7.08	7.96	8.85	9.73	10.62	11.50	12.38	13.27	14.15	15.04
24	2.77	3.69	4.61	5.54	6.46	7.38	8.31	9.23	10.15	11.08	12.00	12.92	13.85	14.77	15.69
25	2.89	3.84	4.81	5.77	6.73	7.69	8.65	9.62	10.58	11.54	12.50	13.46	14.42	15.38	16.35
26	3.00	4.00	5.00	6.00	7.00	8.00	9.00	10.00	11.00	12.00	13.00	14.00	15.00	16.00	17.00

Table continued.

Days	\$18.	\$19.	\$20.	\$21.	\$22.	\$23.	\$24.	\$25.
1	.69	.73	.77	.81	.84	.88	.92	.96
2	1.38	1.46	1.54	1.62	1.70	1.77	1.84	1.92
3	2.08	2.19	2.31	2.42	2.54	2.65	2.74	2.88
4	2.77	2.92	3.08	3.23	3.38	3.54	3.70	3.85
5	3.46	3.65	3.85	4.04	4.24	4.42	4.62	4.81
6	4.15	4.38	4.62	4.85	5.08	5.31	5.54	5.77
7	4.85	5.12	5.38	5.65	5.92	6.19	6.46	6.73
8	5.54	5.85	6.16	6.46	6.76	7.08	7.38	7.69
9	6.23	6.58	6.92	7.27	7.62	7.95	8.30	8.65
10	6.92	7.31	7.69	8.08	8.46	8.85	9.24	9.62
11	7.62	8.04	8.46	8.88	9.30	9.73	10.16	10.58
12	8.31	8.77	9.23	9.69	10.16	10.62	11.08	11.54
13	9.00	9.50	10.00	10.50	11.00	11.50	12.00	12.50
14	9.69	10.23	10.77	11.31	11.84	12.38	12.92	13.46
15	10.38	10.96	11.54	12.12	12.70	13.27	13.84	14.42
16	11.08	11.69	12.31	12.92	13.54	14.15	14.74	15.38
17	11.77	12.42	13.08	13.73	14.38	15.04	15.70	16.35
18	12.46	13.15	13.85	14.54	15.24	15.92	16.62	17.31
19	13.15	13.88	14.62	15.35	16.08	16.81	17.54	18.27
20	13.85	14.62	15.38	16.15	16.92	17.69	18.46	19.23
21	14.54	15.35	16.16	16.96	17.76	18.58	19.38	20.19
22	15.23	16.08	16.92	17.77	18.62	19.46	20.30	21.15
23	15.92	16.81	17.69	18.58	19.45	20.35	21.24	22.11
24	16.62	17.54	18.46	19.38	20.30	21.23	22.16	23.08
25	17.31	18.27	19.23	20.19	21.16	22.12	23.08	24.04
26	18.00	19.00	20.00	21.00	22.00	23.00	24.00	25.00

EXPLANATION.—The column on the left hand of the table shows the number of days: and the rate per month is seen at the top of the page.

EXAMPLE.—To find the amount of 19 days' work, at \$11 per month: find 19 in the column of days; then move to the right, on the same line, till you come under \$11 (rate per month), and you find \$8.04—the answer.

In all cases, the amount will be found directly under the price per month, and at the right of the given time.

In this table, the wages are cast at 26 working days per 10*

month. For a fraction of a day, take an equal part of the amount for one day, and for rates less than \$3 per month, half what is shown for twice the amount.

Should it be desired to ascertain the wages per day for any given sum per month above \$25, it can be done by adding to or doubling the above amounts. Thus for \$30 per month, take 20 and 10 in the above table and add them; for \$37 per month, take 20 and 10 and 7, and add them; for \$50, take 25 and double it; for \$75 per month, take 25 and triple it, &c.

KEEPING ACCOUNTS.

Blank account books, designed for keeping simple ledger accounts, are generally of two kinds, viz. : Those in which the *Dr.* and *Cr.* sides of the account are on the same page, and those in which they are on opposite pages. We give below samples of each, with the mode of keeping the account.

Page 72		WILLIAM WILSON.	<i>Dr.</i>	<i>Cr.</i>
1861.				
Jan.	12	To 18 bus. pota'toes, at 50 cts.....	\$9 00	\$
"	20	" 1 ton hay, at \$8.....	8 00	
"	25	By cash on account.....		10 00
Feb.	7	To 1 yoke steers sold you this day.....	80 00	
"	15	By 2000 feet pine boards, at \$10. m.....		20 00
"	20	To 30 bus. oats, at 30 cts.....	9 00	
Mar.	5	" 40 " corn, at 50 cts.....	20 00	
April	3	By 1 pair boots for S m.....		4 00
"	10	" 50 lbs sugar at 8 cts.....		4 00
May	12	" 10 " coffee at 15 cts.		1 50
June	7	To 3 cords wood at \$2 50.....	7 50	
"	20	By cash on account.....		50 00
July	1	By balance of account charged below.....		44 00
			133 50	133 50
July	1	To balance of account	44 00	

Page 72.		WM. WILSON.	<i>Dr.</i>	Page 73.	WM WILSON.	<i>Cr.</i>
1861.						
Jan.	12	To 18 bus. pota'toes		Jan.	25	By cash on account, \$10 00
		at 50 cts.	\$9 00	Feb'y	15	By 2000 feet pine
"	20	To 1 ton hay at \$8	8 00			boards, at \$10 M 20 00
Feb.	7	To 1 yoke steers sold		April	3	By 1 pair boots for
		you this day....	80 00			Sam..... 4 00
"	20	To 30 bus oats, at		"	10	By 50 lbs. sugar, at
		30 cts	9 00			8 cts..... 4 00
Mar.	5	To 40 bus. corn, at		May	12	By 10 lbs. coffee, at
		50 cts.....	20 00			15 cts..... 1 50
June	7	To 3 cords wood, at		June	20	By c sh on account, 50 00
		\$2 50.....	7 50	July	1	By balance of acct
			33 50			charged..... 44 00
July	1	To balance of act....	44 00			133 50

*Form of a Bill of the foregoing.*WILLIAM WILSON, *Dr.*

1861.	IN ACCOUNT WITH THOMAS BUNN, <i>Cr.</i>	
January 12,	To 18 bus. potatoes, at 50 cts.....	\$9 00
“ 20,	“ 1 ton hay, at \$8.....	8 00
February 7,	“ 1 yoke steers	80 00
“ 20,	“ 30 bus. oats, at 30 cts.....	9 00
March 5,	“ 40 bus. corn, at 50 cts.....	20 00
June 7,	“ 3 cords wood, at \$2 50.....	7 50

1861	<i>Cr.</i>	\$133 50
January 25,	By cash on account.....	\$10 00
Feb'y 15,	“ 2000 ft. lumber, at \$10 M	20 00
April 3,	“ 1 pair boots for Sam....	4 00
“ 10,	“ 50 lbs. sugar, at 8 cts....	4 00
May 12,	“ 10 lbs. coffee, at 15 cts..	1 50
June 20,	“ cash on account.....	50 00
July 1,	To balance.....	\$44 00

NOTE.—Since the whole science of book-keeping rests upon *charges* and *credits*, if you, once for all, get what is a charge and what is a credit clearly fixed in your mind, and fully understand when you ought to charge and when you ought to credit, you will have little difficulty in keeping your accounts straight, simple, and satisfactory.

When you let your neighbor, or he with whom you deal, have anything from you, it is a *charge* against him, and you must charge him with it on the debit side of the account; but whenever you receive anything from him, it is a *credit*, and you must credit him with it on the credit side of the

account. Thus you "charge" for what you give, and "credit" for what you receive. He with whom you deal does likewise—charging you with what he gives you, and crediting you with what he receives from you. Hence his charges against you will correspond with your credits to him, and his credits to you will correspond with your charges against him.

In like manner, should it be desired to keep an account with a certain field, or meadow, or cow, the name is entered at the top of the page and in the index, just as an individual's, and what you give to it, the labor it costs you, &c., you charge to it, and what it yields you you credit to it. In this way a farmer can keep an account with each of his fields or altogether, with each of his cows or with the herd, with each of his pigs or altogether, with each of his sheep or with the whole flock, &c.

The word "*To*" prefixed to an entry indicates a charge or debit; the word "*By*" indicates a credit.

Each entry should be made on the day the transaction took place.

The account should be cast and balanced at least once every six months, and if not settled the balance brought down, as above, when the account may be continued.

BOOK-KEEPING BY DOUBLE ENTRY.

Book-keeping by *double entry* is that form of keeping accounts in which two entries are made in the Ledger for

every one in the Day-Book ; one a charge, or *debit*, and the other a *credit*. Thus you not only charge the party who *receives* from you, but you credit that department of your business from which, whatever it is, is received. You keep an account with as many different departments of your business as you deem necessary. A farmer might keep an account with his herd, with wheat, rye, corn, grass, hay, and other crops, or different fields, separately or together, under the head of "Farm." Where the time required can be spared, we think it desirable to keep accounts by double entry with every department of a business, down to a very minute detail, because where books are kept by this system, you can turn to any account and ascertain at a glance its condition ; that is, how much money you have spent on it, and how much it has returned you, and what balance is for or against it. The books necessary to be used in keeping accounts by this system are two, the Day-Book and Ledger. A third, called a Journal, is sometimes used intermediary between the Day-Book and Ledger ; but we consider it much more trouble than benefit, and therefore think best entirely to dispense with it.

The Day-Book is ruled with two dollar and cent columns on the right hand side, and one column on the left hand side, in which the page of the Ledger is entered when the account is transferred to the Ledger.

The Ledger is generally ruled, as in the example given below ; the name of the account is written across the top of

the page, and if the transactions will probably be numerous other pages following may be reserved to continue the account upon when the first page is full.

It is customary with a person keeping books by this method to have an account with "Cash," with his family, and if he takes and gives notes, with "Bills Receivable," and "Bills Payable." We will give below a sample of transactions entered in the Day-Book and carried to the Ledger. If I sold, October 1st, to John Brown, twenty bushels of apples, at 75 cents per bushel, and was to deliver them to him for \$1, and on October 5th, bought of him five barrels of flour, for family use, at \$4 per barrel, which he was to deliver gratis, my entries in the Day-Book would be as follows, supposing I kept accounts with the departments mentioned:—

Page 1.

CENTERVILLE, OCT. 1st, 1861.

5	JOHN BROWN, <i>Dr.</i>				
	Sold him 20 bus. apples, at 75 cts per bus.	\$15 00			
	Cartage.....	1 00	16 00		
	<i>Cr.</i>				
6	ORCHARD,			15 00	
7	TEAMING			1 00	
	— Oct. 5 —				
10	FAMILY EXPENSE. <i>Dr.</i>				
	Bought of John Brown 5 bbls flour, at \$4 per bbl	20 00			
	<i>Cr.</i>				
5	JOHN BROWN.....			20 00	

The Ledger accounts of the above would be as follows:—

Page 5.
Cr.

<i>Dr.</i>		JOHN BROWN.			
1861.			1861		
October	1	16 00	Oct.	5	20 00

TEAMING.										Page 7. Cr.
Dr.						1861 Oct.	5			1 100

ORCHARD.										Page 6. Cr.
Dr.						1861. Oct.	1			1 00

FAMILY EXPENSE.										Page 10. Cr.
Dr.	1861. Oct.	5				1	20	00		

In the Day-Book, in the right hand dollar and cent column, the credits are entered ; in the left hand, the debits, as shown.

In the Ledger, the half of the page to the left of the centre is devoted to debits ; to the right, to credits. The column to the left of the dollar and cent column in the Ledger is where the page of the Day-Book from which the entry is taken is noted.

The form which we have given above is, perhaps, the simplest in which books can be kept by double entry, consequently the best. No difficulty will be experienced in this system of keeping books, after one has already fixed in his mind what is a charge or debit, and what is a credit, as explained above. Some remarks may not, however, be unne-

cessary in this connection, to show *what* to credit and *what* to charge, under certain circumstances. If you give a man a note for the balance of his account, you debit his account and credit Bills Payable. When you pay the note, you debit Bills Payable and credit Cash. If you receive a note for balance of account, you credit the man's account and debit Bills Receivable. When the note is paid, you credit Bills Receivable and debit Cash. In the first entry in the above example, it may be well to say, you do not give credit to the man who drives the wagon, or to the wagon for its use. These are legitimate charges against Teaming. At the proper time you credit the man his wages, and charge or debit Teaming for it (or that portion of his time in which he has been engaged teaming), &c.

Some businesses require an Interest account to be kept; of course, from our previous remarks, any one who finds it necessary will see the proper way to keep it.

It is necessary, in connection with the Day-Book and Ledger, to keep a Cash-Book and Bill-Books, where a person does a credit business. The Cash-Book, to keep a record of the receipts and disbursements of cash, which should be balanced every night (if any cash has been spent or received during the day), and the money counted; the balance on hand and the balance shown by the book should correspond; if they do not, something has been omitted. If you have on hand more than the balance calls for, you have received money which has not been entered on the debit side of the

account. If you have too little, you have spent money for which the account has not been credited.

The Bill-Books are to keep a record of notes received and notes paid out. The Bills Payable book records the following facts: The date of the note, the time it is to run, the date of falling due, to whom it was given, in whose favor it was made, and the amount it was made for. The Bills Receivable book records: Who made the note, in whose favor it was made, how long it has to run, when it is due, and the amount it is for. When notes are paid or received, these facts should, of course, be properly noted in the Day-Book.

When accounts are first opened it is best to take an inventory of property of all kinds on hand, charging each department with which you intend to keep an account with that portion which it requires, and crediting an account for the same which shall represent all your "Stock in Trade." This account is usually called "Stock." Then, at the time you wish to close up your accounts to ascertain your profits and losses, you take another inventory, and give your departmental accounts credit for what property they have on hand, charging the general stock account for the same; the balance of this account (*i. e.*, the difference between the footing of the debit and credit columns) then shows how much more or less property you have on hand than when you commenced business. If the credit side exceeds the debit, of course you have more property; and if the debit

exceeds the credit, of course you have less than when you began. Then the balance of each departmental account (all proper charges having been entered, and its share of property on hand credited) will show how much it has made or lost. These balances are then usually carried to a general account, called "Profit and Loss;" those having a credit balance are charged that amount, and Profit and Loss is credited; and those having a debit balance are credited that amount, and Profit and Loss is charged for it. This being done with the Departmental accounts and the General Stock accounts, with the Cash accounts, and the Bills Payable and Bills Receivable accounts, and Profit and Loss having been also charged for bad debts—and the parties owing them having been credited therefor—the balance of that account shows the Profit and Loss of the business. Some parties do not credit the accounts of persons who owe bad debts, and charge Profit and Loss; but, after making up the Profit and Loss account, draw it off on a sheet of paper, and account for them there. Others open an account called "Suspense," to which they credit the amount of the several bad debts (specifying them in the Day-Book), and charge Profit and Loss. This method prevents the accounts of bad debtors appearing closed on your Ledger. After you have made up your books as directed, it is best to make a balance sheet, which will show at a glance what departments have made money, what lost, who owes you, and who you owe. After this, the several departments should be charged back again

This we think of no advantage, and it increases the labor and trouble. When you render a bill from the account, you must necessarily turn to the Day-Book to ascertain the particulars, and the mere page of the Day-Book is sufficient for this purpose. The less accounts are complicated the easier they are kept, and the less liable are mistakes to be made.

No erasures, scratching out, or interlineations should be suffered. If a wrong entry be made, or an entry made wrongly, let it be explained by a counter entry on the other side of the account, or overscored in such a manner that the mistake can be seen. All erasures, blotting out, scratching, &c., tend to throw suspicion upon the honesty of the account.

Books of "Original Entries" are only an aid of the memory, and he who keeps them should be able to swear that the entries were made on the day they purport to have been. He may not be able to recollect the various entries, but if it was his invariable custom to make them on the day of the transaction, they stand in place of his memory—they are not, however, evidence of the *delivery* of the goods.

Form of a Receipt in full.

NEW YORK, July 1st, 1861.

Received of Thomas Brown the sum of forty-four dollars,
in full of all accounts up to this date.

\$44 00.

WILLIAM WILSON.

Form of a Check.

\$150 00.

NEW YORK, July 1st, 1861.

Please pay William Wilson, or order, one hundred and fifty dollars, and charge to the account of

THOMAS ANDERSON.

To the Southold Savings Bank.

Form of a Due-Bill.

NEW YORK, July 1st, 1861.

Due William Wilson, or order, on settlement this day, one hundred and fifty dollars.

\$150 00.

THOMAS ANDERSON.

Form of a Promissory Note.

NEW YORK, July 1st, 1861.

Four months after date I promise to pay William Wilson, or order, one hundred and fifty dollars; value received.

\$150 00.

THOMAS ANDERSON.

Another form.

NEW YORK, July 1st, 1861.

On the 1st day of April next, I promise to pay William Wilson, or order, one hundred and fifty dollars; value received.

\$150 00.

THOMAS ANDERSON.

Form of a Promissory Note with Surety.

NEW YORK, July 1st, 1861.

Sixty days after date, we, or either of us, promise to pay William Wilson, or order, one hundred and fifty dollars; value received.

THOMAS ANDERSON, (Principal.)

\$150 00.

JOHN JONES, (Surety.)

Form of a Draft or Bill of Exchange.

\$150 00.

BUFFALO, July 1st, 1861.

Ten days after sight, pay William Wilson, or order, one hundred and fifty dollars, value received, and charge the same to account of

Yours, &c.,

THOMAS ANDERSON.

To William Allen, New York.

NOTES.—A due-bill bears interest from its date; a promissory note not until after it is due, unless so expressed on its face.

NEGOTIABILITY.—The words, “or order,” “or bearer,” are necessary to make a check, a due-bill, a promissory note, a bill of exchange, &c., *negotiable*; that is, to enable the holder of it to trade and pass it to another.

When the words “or bearer” are introduced, the instrument may then pass from hand to hand, like a bank-bill, without endorsement; but when the words “*or order*” are used, the instrument must be endorsed by the original holder of it.

ENDORSEMENT.—Endorsing a note is writing your name across the back of it. Endorsements are of two kinds, an endorsement in blank or general endorsement, and a special endorsement.

An *endorsement in blank* is the original holder's simply writing his name across the back of it. The succeeding holders of it may or may not, also, endorse it. If each or

any of them do, they also become severally bound for its payment.

A *special endorsement* is made by writing across the back of it, before endorsing it, the words, "Pay to the order of [name of party to whom it is passed]," which *limits* the payment of it to that party, or *his* orders, and so forth.

ACCEPTANCE.—When a draft or bill of exchange is made upon a third party (as in the above form), the latter is not in any way bound by it until he accepts it, which he does when it is presented to him for acceptance, by writing across the face of it the word "*accepted*," with the date, and signing his name thereunder. He is then a party to the bill, and bound for its payment at maturity.

PROTEST.—Protest is the notice required by law to be given to the endorsers of promissory notes, and the makers and endorsers of bills of exchange, of their dishonor, that is, of their non-acceptance or non-payment.

If the drawee, or person to whom a bill of exchange is directed, refuses to accept it on presentation, notice must be immediately given to the maker of it.

If he accepts it, and afterwards fails to pay it at maturity, notice must immediately be given to the maker.

If the maker of a promissory note fails to pay it at maturity, notice must immediately be given to all the endorsers.

A check is a draft at sight, and if not paid, must be protested.

It is a general rule that all *guarantors* of commercial paper must be immediately notified of its dishonor.

It is, of course, not necessary to protest a due-bill, or a promissory note, which is still held by the person to whom it was originally given.

When a note is made payable "*on demand*," it is necessary to make a demand before it will bear interest or can be sued for.

U. S. BONDS.

Interest is calculated on U. S. bonds and on the public debt at 365 days to the year, and is due semi-annually. In England interest is calculated in the same way, and the legal rate is 5 per cent.

By Five-Twenties is meant the 6 per cent. gold-bearing bonds of the United States, which are to mature in 20 years, but which the Government, by giving due notice, can pay in gold any time after five years from the date of issue.

The old five-twenties were the first issued. They bear date May 1, 1862, and are redeemable after May 1, 1867, and payable May 1, 1882. The new "five-twenties" were issued Nov. 1, 1864, July 1, 1865, and Nov. 1, 1865.

By Ten-Forties is meant the 5 per cent. gold-bearing bonds which are to mature in 40 years, but which may be paid by the Government at any time after 10 years.

By Seven-Thirties is meant a *currency* loan, which matures in 3 years, at which time they may be changed for the *five-twenty* 7 per cent. bonds, bearing interest in gold. The name is derived from the rate of interest, it being 7.3 per cent. The "First series" bear date Aug. 15, 1864. The "Second series" bear date June 15, 1865, and are convertible June 15, 1868. The "Third series" bear date July 5, 1865. On this issue the Government reserves the right to

pay the interest at 6 per cent. in gold, instead of 7.30 per cent. in currency.

By Six per cents. of '81 is meant the 6 per cent. gold-bearing bonds which cannot be redeemed by Government, except by purchase, until after maturity.

RELATIVE VALUE OF GOLD AND CURRENCY.

To ascertain the value in gold of a "greenback" dollar or National currency, at the different quotations of gold :

RULE.—Divide \$1 by the quoted value of \$1 in gold ; the result will be *the value* of a dollar in currency.

EXAMPLE.—When gold is 33 per cent. premium what is the value of \$1 in currency? $\$1.00 \div \$1.33 = .7522$.

NOTE.—In the following table the decimals are carried to mills and tenths of a mill.

TABLE, *showing the greenback value of \$1 at the different quotations of gold. When gold is at*

.01 pr. ct. prem. a greenback dollar is worth. .99	.12 pr. ct. prem. a greenback dollar is worth. .8929
.02..... .9803	.13..... .885
.03..... .9708	.14..... .8771
.04..... .9615	.15..... .8695
.05..... .9523	.16..... .862
.06..... .9433	.17..... .8564
.07..... .9355	.18..... .8474
.08..... .9259	.19..... .8403
.09..... .9174	.20..... .8333
.10..... .909	.21..... .8264
.11..... .9009	.22..... .8279

.23 pr. ct. prem. a greenback dollar is worth. .813	.37 pr. ct. prem. a greenback dollar is worth.. .7308
.24..... .8064	.38..... .7246
.25..... .80	.39..... .7194
.26..... .7928	.40..... .7142
.27..... .7874	.41..... .7092
.28..... .7812	.42..... .7042
.29..... .7751	.43..... .6993
.30..... .7692	.44..... .6944
.31..... .7633	.45..... .6896
.32..... .7575	.46..... .6849
.33..... .7522	.47..... .2162
.34..... .7462	.48..... .6758
.35..... .7409	.49..... .6716
.36..... .7353	.50..... .6666

NOTE.—The highest quotation of gold at the New York Stock Exchange during the war was 285, July 11th, 1864. A dollar currency was then worth 35 cents. Gold in Richmond, Va., reached 4400, Feb. 4th, 1865. A dollar in Confederate currency was worth .02 $\frac{1}{4}$ cents.

ENGLISH BONDS AND CONSOLS.

Exchequer Bills are English bonds, similar to those of the U. S. The rates of interest vary from 5 to 3 per cent., and while the Government pays the interest, it cannot be required to refund the principal.

Consols are several English securities consolidated by act of Parliament. The rate of interest is 3 per cent.

The Stock Exchange is an association for the purpose of buying and selling stocks.

A Broker is a person who executes orders for those who are not members of the exchange.

A Jobber deals in stock on his own account. A "stag," or "outsider," is a broker who is not a member of the exchange.

A Bull is one who buys stock to be delivered to him at a future time, with the intention of selling it, in the meantime, at a higher price before he is obliged to receive it.

A Bear is one who sells stock that he does not own, to be delivered at a future date, hoping in the meantime to buy it at a less price. A "lame duck" is one who is unable to fulfil his contracts, and hence is expelled from the exchange.

"Selling Short" is applied to sales of stock which the seller does not own, deliverable at a future time, generally not exceeding 60 days. The *bears* usually "sell short." The buyer pays interest for over 3 days.

"Seller's Option" gives the seller the privilege of delivering the stock at any time before the time specified for delivery.

"Buyer's Option" gives the purchaser the privilege of claiming the delivery of the stock at any time before the time specified for delivery.

STOCK QUOTATIONS.

From N. Y. Herald.

Sales.		
12000 Am. G.....	148½	\$12000 gold at 43½ per cent. premium.
12000 U. S. 6's. '81 cou.....	112½	{ \$12000 U. S. 6 per cent. coupon bonds, maturing 1881, at 12½ per cent. premium.
10000 U. S. 5-20 Reg. '62.....	104½	
		U. S. 5-20 Registered Bonds issued in 1862.

40000 Tr'y. N. 7-30 2d s.....	107	Treasury notes at 7.3 per cent. interest, second series.
100 N. Y. Cen. 7's, '65-'76.....	120	100 Shares of New York Central R.R. 7 per cent. bonds issued in 1865, and maturing in 1876.
500 Hud. R. 7's 1st M.....	101	Hudson R. 7 per cent. first mortgage bonds.
200 " " 2d M. S. F.....	104	Hudson R. 7 per cent. second mortgage sinking fund.
100 E. R.R. 2 d's.....	51½	Erie R.R. sold at 2 days' credit at 51½ cents per dollar.
100 " " b. 5 w. n.....	51	Erie R.R. to be delivered before 5 days without notice.
1000 C. and Am. 6's, '89.....	108½	Camden and Amboy 6 per cent. maturing in 1889.
100 Mich. C. 6's, b. 15 and int.....	107½	Mich. Cent. R.R. Stock to be delivered before 15 days with interest.
100 U. S. Tr. 7-30 int. F. & A.....	111½	U. S. Treasury notes 7.3 per cent. interest paid in February and August.
1000 Erie pref. s., w. n.....	88	Erie Preferred Stock without notice.
100 Penn. 6's, int. off.....	104	Pennsylvania 6 per cent. stock, the last interest of which has been paid.
500 E. R.R. b. u.....	23	Erie R.R. stock, "Buyer's Option," when to call for the stock.
300 " " s. u.....	31	Erie R.R. stock, "Seller's Option," when to deliver the stock.

SUCCESS IN BUSINESS.

SHORT CREDITS.

Short credit has much to do with the amount of profits in business. The difference between long and short credits will be seen by the following table, showing the amount of \$100 in ten years.

	Am't at 3 pr. ct.	Am't at 5 pr. ct.	Am't at 8 pr. ct.	Am't at 10 pr. ct.
If turned over every 3 months,	\$326.20	\$703.99	\$2172.45	\$4525.92
" " " 6 "	180.61	265.32	466.09	672.75
" " " 8 "	155.79	207.89	317.21	417.72
" " " 12 "	134.39	162.88	215.89	259.37
" " " 2 years,	115.92	127.62	146.93	161.05
" " " 5 "	106.09	110.25	116.64	121.00

SMALL PROFITS.

By the above table it will be seen that quick sales and

small profits are more desirable than large profits and long credits. It must be considered, however, before reducing profits, whether the sales can be increased so as to compensate for the reduction of profits.

ECONOMY IN EXPENSE.

Many a young man in business fails to succeed, owing to a want of economy in expense. *All expense must be deducted from the profits.* "Fortunes are spent by trifles." "A penny saved is worth two earned."

2	cents per day in ten years will amount to \$100.85 at 7 p. c. compound int.				
5	"	"	"	252.14	" "
25	"	"	"	1260.71	" "
50	"	"	"	2521.42	" "
100	"	"	"	5042.84	" "
\$2	"	"	"	10085.68	" "

MARKING GOODS.

It is customary among merchants to use a private mark to denote the cost and selling price of goods. Any word or phrase containing ten different letters is selected, and used to represent figures, as "White sugar," "Misfortune," &c., thus:

w	h	i	t	e	s	u	g	a	r
1	2	3	4	5	6	7	8	9	0

An extra letter called a "repeater" is generally used to prevent the repetition of a figure, as *x*, *y*, or *z*, &c.; thus 388 would be represented by *igz*, using *z* as a repeater, instead of *igg*. The object is to prevent a clue being given to

the key-word. Any mark or character may be used to represent a figure instead of a letter. Fractions may be written thus, $473\frac{2}{3} = twi\frac{h}{2}$; or by an arbitrary mark, thus, \circ may represent $\frac{2}{3}$, then $473\frac{2}{3} = twi\circ$. Sometimes the cost mark is written below a line. and the selling price above; thus,

4.62 *tsh* sell. pr.

— = —

3.24 *iht* cost.



THE OTIS PATENT LIGHTNING ROD,
AS APPLIED TO THE N. Y. STATE ARSENAL.
M. WELLS, Electrician, 112 Broadway, N. Y.

LIGHTNING-RODS.

The humid gases, generated by the heating and sweating of the hay, which immediately follows its accumulation in closely-packed masses, offers a strong attraction to electricity, just at the time when it is most abundant. It is an object of peculiar importance to the farmer to guard his buildings, at such times, with properly constructed lightning-rods; and they are a cheap mode of insurance against fire from this cause, as the expense is trifling and the security great.

As an example of the more elaborate style of rods, we show in the accompanying cut the manner in which Otis' Patent Lightning-Rods have been applied to the New York State Arsenal.

To construct a lightning-rod.

Take round or square soft iron $\frac{3}{4}$ of an inch in diameter, in pieces of convenient length; connect the pieces by splitting one end and flattening and inserting the other, and fasten with a rivet or screw, so that the rod preserves its uniform thickness throughout. Or, the pieces may be connected in a more perfect manner, although not often so convenient, by having a male screw cut on one end of the pieces and a female screw on the other, and simply screwing them together as the rod is raised; care being taken that the

pieces are brought in contact at the outer edge, so as to form a united surface. If a square rod is used, notch the corners with a single downward stroke of a cold chisel, at intervals of two or three inches. No part of the rod should be painted, as its efficiency would be greatly impaired. Let the upper extremity consist of one finely drawn point of copper or silver, or well gilded iron, to prevent rusting. Let the lower part of the rod, at the surface of the ground, terminate in two or three flattened divergent branches, leading several feet outwardly from the building, and buried at a depth which reaches perpetual moisture, in a bed of charcoal. Attach the rod to the building by clasps protruding three or four inches and containing glass rings or funnels for the rod to pass through. The rod must not touch the building nor the iron clasps, but only the glass; because, the latter being a non-conductor of electricity, in the event of the rods being struck by lightning, the charge is conducted harmlessly to the ground, having no point of contact with a conductor by which it might be led into the building. Upon reaching the top of the building the rod should be conducted to the centre of the ridge, and the end should then be raised to a height equal to one-half of the distance to the end of the ridge. If the roof is irregular in height, of course judgment must be used in fixing the point where the end of the rod rises above the roof, bearing in mind this important consideration—that the rod protects objects at twice the distance of its height above any point in a line perpendicular to its upper termination.

The conducting power of bodies is in the ratio of their surfaces. Hence a bundle of wires, ribbons, or tubes of metal, are more efficient than an equal quantity of solid, round, or square rods.

The *conductors* of electricity in the order of their power are, copper, silver, gold, iron, tin, lead, zinc, platinum, charcoal, black lead, strong acids, soot and lampblack, metallic ores, metallic oxides, dilute acids, saline solutions, animal fluids, sea-water, fresh water, ice, living vegetables, living animals, flame, smoke, vapor and humid gases, salts, rarified air, dry earth and massive minerals.

The *non-conductors* in their order are, shellac, amber, resins, sulphur, wax, asphaltum, glass, all vitrified bodies, raw silk, bleached silk, dyed silk, wool, hair, feathers, dry paper, parchment and leather, baked wood and dry vegetables.

The question of the utility of lightning-rods is not clearly decided; and certainly very grave doubts exist as to the usefulness of the various complicated patent devices which are hawked about the country, under the sanction of splendid testimonials.

Dr. Franklin's theory was,—and he claimed to have proved it by having drawn the electricity harmlessly from a cloud over his kite-string,—that the value of the lightning-rod consists, not in its ability to receive shocks, but in the fact that it taps the surcharged clouds and conveys the electricity quietly to the earth.

Based upon this theory, there has recently been advanced an idea that seems sensible. It is to substitute a piece of galvanized telegraph wire for the kite-string, a pointed rod of iron at the top of the building for the kite, and another rod driven into the ground for the key in Dr. Franklin's hand.

The iron at the top should project five or six feet above the roof, and if the ridge-pole is more than twenty feet long, there should be two or more of these, all to be connected with each other and with the rod in the ground by simple wire.

This plan has the great merit of being cheap and within the reach of all—and, so far as anything is actually known of the subject, is as good as the more elaborate and expensive ones.

PRESSURE OF EARTH AGAINST WALLS.

To find the pressure of the different kinds of earths, filling, &c., against walls, it is necessary first to ascertain the line or angle of rupture, or *natural slope*, the earth would assume but for the resistance of the wall. This natural slope differs with the different kinds of earths. Assuming that the earth is level with the top of the wall, the line of rupture for the different kinds of earths, filling, &c., will be as follows:—

A bank of *vegetable earth* will rupture on the surface at a distance from the top of the wall of three-fifths the height of the wall.

A bank of *sand* will rupture at two-thirds the height of the wall.

A bank of *unhewn stone*, at one-seventh the height of the wall.

A bank of *rubble* at two-fifths the height of the wall.

A bank of *brick*, with a bank of vegetable earth behind it, will rupture at a distance of about one-sixth the height of the wall.

A bank of *clay*, well rammed, will rupture at a distance of three-sixteenths the height of the wall.

The strongest horizontal stress against the wall is at half the angle which the natural slope makes with it; hence:

The greatest pressure for a bank of *vegetable earth* will be at three-tenths the height of the wall from the bottom.

For a bank of *sand*, at one-third the height of the wall.

For a bank of *rubble*, at one-fifth the height of the wall.

For a bank of *unhewn stone*, at one-fourteenth the height of the wall.

For a bank of *brick*, at one-twelfth the height of the wall.

For a bank of *clay*, at three-thirty-seconds the height of the wall.

Walls should therefore be built proportionably strong to these heights to sustain the different pressures. If the bank is liable to be saturated with water the wall should be doubled in strength.

FRACTIONS—DECIMALS.

A fraction is one or more parts of a unit, and is expressed by fractional characters, thus, $\frac{1}{2}$, $\frac{1}{4}$, $\frac{3}{4}$; or by decimals, thus, .5, .25, .75.

When expressed by fractional characters, the upper figure is called the *numerator*, because it numbers or gives value to the fraction, by showing how many parts of the whole number into which the unit is divided is taken; and the lower figure is called the *denominator*, because it names the number of parts into which the unit is divided. Thus, $\frac{3}{8}$ means that the unit is divided into 8 parts, and that 3 out of the 8 are taken, &c.

When expressed by a decimal, the decimal number shows that so many parts of the unit are taken, the unit itself being impliedly divided into as many parts as will correspond with the decimal number, and still retain its ratio to it. Thus, .5 means $\frac{5}{10}$, .25 means $\frac{25}{100}$, .125 means $\frac{125}{1000}$, &c.

To reduce fractions to decimals.

Divide the numerator by the denominator, adding cyphers as required.

EXAMPLE.—What are the decimals of $\frac{1}{2}$, $\frac{3}{4}$, and $\frac{7}{8}$?

SOLUTION.— $10 \div 2 = .5$, $300 \div 4 = .75$, $7000 \div 8 = .875$. *Ans.*

To add decimals.

Add as in common addition, setting the whole numbers

or integers directly under each other from the decimal point to the left, and the decimals from the decimal point to the right, as in the following example:—

$$\begin{array}{r}
 12.75 \\
 24.027 \\
 14.5 \\
 16.1278 \\
 \hline
 67.4048
 \end{array}$$

To subtract decimals.

Set the whole numbers and decimals under each other, as directed above, and proceed as in common subtraction, as in the following example:—

$$\begin{array}{r}
 75.15 \\
 28.875 \\
 \hline
 46.275
 \end{array}$$

To multiply decimals.

Set the figures and multiply as in common multiplication, and point off in the product as many decimals as there are decimal places in the multiplier and multiplicand, as in the following example:—

$$\begin{array}{r}
 23.25 \\
 22.15 \\
 \hline
 11625 \\
 2325 \\
 4650 \\
 4650 \\
 \hline
 514.9875
 \end{array}$$

To divide decimals.

Proceed as in common division, and point off to the right in the quotient as many decimals as the decimal places in the dividend exceed the decimal places in the divisor, as in the following example :—

$$\begin{array}{r}
 2.48 \overline{) 129.952} \quad [52.4 \\
 \underline{124 } \\
 5 5 \\
 \underline{4 6} \\
 96
 \end{array}$$

Useful decimals.

$\frac{1}{16}$.0625	$\frac{1}{8}$.333+	$\frac{1}{8}$.1111+
$\frac{2}{16}$ ($\frac{1}{8}$)	.125	$\frac{2}{8}$.666+	$\frac{2}{8}$.2222+
$\frac{3}{16}$.1875	$\frac{3}{8}$.2	$\frac{3}{8}$.3333+
$\frac{4}{16}$ ($\frac{1}{4}$)	.25	$\frac{4}{8}$.4	$\frac{4}{8}$.4444+
$\frac{5}{16}$.3125	$\frac{5}{8}$.6	$\frac{5}{8}$.5555+
$\frac{6}{16}$ ($\frac{3}{8}$)	.375	$\frac{6}{8}$.8	$\frac{6}{8}$.6666+
$\frac{7}{16}$.4375	$\frac{7}{8}$.1428+	$\frac{7}{8}$.7777+
$\frac{8}{16}$ ($\frac{1}{2}$)	.5	$\frac{8}{8}$.2856+	$\frac{8}{8}$.8888+
$\frac{9}{16}$.5625	$\frac{9}{8}$.4285+	$\frac{9}{8}$.1
$\frac{10}{16}$ ($\frac{5}{8}$)	.625	$\frac{10}{8}$.5712+	$\frac{10}{8}$ ($1\frac{1}{4}$)	.2
$\frac{11}{16}$.6875	$\frac{11}{8}$.7141+	$\frac{11}{8}$.3
$\frac{12}{16}$ ($\frac{3}{4}$)	.75	$\frac{12}{8}$.8569+	$\frac{12}{8}$ ($1\frac{1}{2}$)	.4
$\frac{13}{16}$.8125			$\frac{13}{8}$.7
$\frac{14}{16}$ ($\frac{7}{8}$)	.875			$\frac{14}{8}$.9
$\frac{15}{16}$.9375			$\frac{15}{8}$	

FACTS ABOUT PRINTING AND BOOK- MAKING.

The following are the different styles of type ordinarily used in book-printing:—

PICA.

Springs are weakened by use, but recover their strength if laid by.

SMALL PICA.

Metals have five degrees of lustre—splendent, shining, glistening, glimmering, and dull.

LONG PRIMER.

The hardness of metals is as follows: Iron, Platinum, Copper, Silver, Gold, Tin, Lead.

BOURGEOIS.

A fall of 1-10 of an inch a mile will produce a current in rivers.

BREVIEW.

Melted snow produces about 1-8 of its bulk of water.

MINION.

Silica is the basis of the mineral world, and carbon of the organized.

NONPAREIL.

Sound passes in water at a velocity of 4703 feet per second, and in air 1100 feet, at a temperature of 33°.

AGATE.

At the depth of 45 feet, the temperature of the earth is uniform throughout the year.

PEARL.

The weight of a cubic foot of air is 527.01 grains, or 1 205 ounces, avoirdupois.

NOTE.—*Diamond* is smaller than pearl—*Emerald* still smaller.

We do not apologize for giving the above and the few following facts about printing, because that art has become so universally used by all classes that it is of practical importance to disseminate information in regard to it.

The specimens given above are called Roman; CAPITALS and SMALL CAPITALS belong legitimately with this style. *Italics* are cast to accompany it, to give emphasis to certain parts of the matter being composed, or set up. Italic figures and small capitals of italic are not made. Many other styles of type, such as Black Letter, Script, Church Text, Clarendon, Title, Ionic, Full Face, &c., are cast, and are ordinarily used to display certain lines in Job Printing, and are consequently called job type.

Printers generally charge for the setting of type, or, as they technically term it, the composition of matter, by the number of *ems* it contains. An *em* is the square of the body of the type; they measure the matter composed by multiplying the number of ems or lines it is in length by the number of ems or lines it is in width. Nonpareil is half the size in body of Pica, consequently 4 ems of Nonpareil equal 1 of Pica. Agate is half Small Pica. Pearl is half Long Primer.

In 1 square inch there are.....	36	ems Pica.
“ “ “	50	“ Small Pica.
“ “ “	56 $\frac{1}{4}$	“ Long Primer.
“ “ “	72 $\frac{1}{4}$	“ Bourgeois.
“ “ “	87	“ Brevier.
“ “ “	113 $\frac{3}{4}$	“ Minion.

In 1 square inch there are.....	144	ems	Nonpareil.
“ “ “	200 $\frac{1}{2}$	“	Agate.
“ “ “	225	“	Pearl.

That is according to the type in the office where this book is printed ; different foundrymen vary the sizes of type slightly, so that the above is not a perfectly accurate guide in measuring the number of ems in a page or book ; still it is sufficiently so to give a very close approximation to what any printer would measure. In using the above to calculate in ems the contents of a page or book, be particular to calculate *square inches*, not *inches square*. The price of type-setting in New York varies with the different printers. Generally the price for book composition is from 80c. to \$1.00 per 1000 ems. Much figure-work is charged extra, so also is an extra charge made where a very narrow column is set. Pearl is charged extra on account of its smallness. The price given is for plain matter.

Presswork is charged for by the *token*, which is 250 impressions of the press. Prices vary so much per token, according to the quality of the work and the number of impressions, that it is next to impossible to give an idea of it that will benefit the reader. Plain book-work, in editions of 1000 to 2000 copies, is charged usually at 50c. to \$1.00 per token.

SIZES OF BOOKS.

The various sizes of books were named from the number of folds that were made of a sheet of paper 19 inches by 24,

which, at the time the sizes of books acquired their names, was the largest sheet manufactured. Thus, a sheet of that size folded once, making 2 leaves or 4 pages, was called a folio volume; folded twice, making 4 leaves or 8 pages, was called a quarto volume; folded four times, making 8 leaves or 16 pages, was called an octavo; folded six times, making 12 leaves or 24 pages, was called a duodecimo, &c. They are written thus: 2fo., 4to, 8vo, 16mo, &c.

Afterwards, when the sheets came to be manufactured larger, books continued to be designated as above, but were distinguished from the above sizes by giving the new sheets names, and prefixing the name of the sheet to the above. Thus, a sheet 22 inches by 28 was called "Royal," and hence books printed on it were called royal folio, royal quarto, royal octavo, &c.

TABLE, *showing the number of leaves and pages from the folding of a sheet.*

	Folds.	Leaves.	Pages.
Sheet of 19 inches by 24.	2fo.....	2	4
	4to.....	4	8
	8vo.....	8	16
	12mo.....	12	24
	16mo.....	16	32
	18mo.....	18	36
	24mo.....	24	48
	32mo.....	32	64

NOTE.—The foldings, leaves, and pages of the royal sheet, &c., are the same as the above, but the sheet being larger of course the leaves and pages are larger.

STRENGTH OF MATERIALS.

Tensile strength.

Tensile strength is the amount of cohesion existing between the atoms of a mass, or the tenacity with which the fibres or particles of a body resist separation. The tensile strength of a body is therefore in proportion to the number of its fibres, or to the area of its section.

TABLE, *showing the weight in lbs. necessary to tear asunder one square inch of the following substances.*

METALS.		WOODS	
Designation.	Wt. in lbs.	Designation.	Wt. in lbs.
Copper, wrought.....	34,000	Ash	16,000
“ cast	19,000	Beech	11,500
“ wire	61,200	Birch	15,000
Gold, cast.....	20,000	Box	20,000
Iron, “	27,000	Cedar.....	11,400
Iron Wire	103,000	Chesnut	10,500
“ best bar	72,000	Cypress	6,000
“ medium bar.....	60,000	Elm	13,400
“ inferior “	30,000	Fir, strongest.....	12,000
Lead.....	880	“ American	8,800
Platinum wire	53,000	Lig. vitæ	11,800
Silver, cast.....	40,000	Locust	20,500
Steel	120,000	Mahogany	21,000
Tin, block	5,000	Maple	10,500
Zinc, cast	3,500	Oak, American, white..	11,500
Brass	42,000	“ seasoned.....	13,600
		Pine, Pitch.....	12,000
		Poplar	7,000
		Sycamore	13,000
		Walnut	7,000
		Willow	13,000

To find the tensile strength.

RULE.—Multiply the area of the transverse section in inches, by the weight given in the preceding table, and the product will be the strength in lbs.

EXAMPLE.—What is the tensile strength of a seasoned white oak scantling 2 inches by 3?

SOLUTION.— $2 \times 3 = 6$, area of transverse section, $\times 13,600 = 81,600$ lbs. *Ans.*

EXAMPLE SECOND.—What is the tensile strength of a round poplar stick 3 inches in diameter?

SOLUTION.—7.068, area of circle (*vide* table of the areas of circles), $\times 7,000 = 49,476$ lbs. *Ans.*

EXAMPLE THIRD.—What is the tensile strength of the best bar iron, 2 inches broad by 1 inch thick?

SOLUTION.— $2 \times 1 = 2$, area of transverse section, $\times 72,000 = 144,000$ lbs. *Ans.*

NOTE.—The above gives the *maximum tensile strength* of the materials, or the *utmost strain* they are capable of sustaining when drawn lengthwise. But it is to be borne in mind that the *practical value* is about *one-fourth* of the above.

TABLE, showing the strength of iron wire rope and hempen cable.

	Circumference of Wire Rope in inches.	Trade Number.	Circumference of Hemp Rope, of equal strength, in inches.	Breaking weight in tons of 2,000 lbs.
Fine Wire, ..	6.02	1	15½	74.
	6.20	2	14½	65.
	5.44	3	13	54.
	4.90	4	12	43.6
	4.50	5	10½	35.
	3.91	6	9½	27.2
	3.86	7	8	20.2
	2.98	8	7	16.
	2.56	9	6	11.4
	2.45	10	5	8.64
	4.45	11	10½	36.
	4.00	12	10	30.
	3.63	13	9½	25.
	3.26	14	8½	20.
	2.98	15	7½	16.
	2.68	16	6½	12.3
	2.40	17	5½	8.8
Coarse Wire, }	2.12	18	5	7.6
	1.9	19	4.75	5.8
	1.63	20	4.	4.09
	1.53	21	3.3	2.83
	1.31	22	2.80	2.13
	1.23	23	2.46	1.65
	1.11	24	2.2	1.38
	0.94	25	2.04	1.03
	0.88	26	1.75	0.81
	0.78	27	1.50	0.56

J. A. Roebling, C. E.

STRENGTH OF CABLES, ROPES, AND HAWSERS.

To find the strength of cables.

RULE.—Multiply the square of the circumference in inches by 120, and the product is the weight in lbs. the cable will bear with safety.

EXAMPLE.—What weight will a cable 6 inches in circumference bear with safety?

SOLUTION.— $6^2 = 36 \times 120 = 4320$ lbs. *Ans.*

To find the strength of ropes and hawsers.

RULE.—Multiply the square of the circumference in inches by 200, and it gives the weight in lbs. the rope will bear with safety.

TABLE, *showing what weight a hemp rope will bear with safety.*

Circumference.	lbs.	Circumference.	lbs.	Circumference.	lbs.
1	200	3 $\frac{1}{4}$	2450.	6	7200.
1 $\frac{1}{2}$	312.5	3 $\frac{3}{4}$	2812.5	6 $\frac{1}{2}$	7812.5
1 $\frac{3}{4}$	450.	4	3200.	6 $\frac{3}{4}$	8450.
1 $\frac{1}{2}$	612.5	4 $\frac{1}{4}$	3612.5	6 $\frac{1}{2}$	9112.5
2	800.	4 $\frac{1}{2}$	4050.	7	9800.
2 $\frac{1}{4}$	1012.5	4 $\frac{3}{4}$	4512.5	7 $\frac{1}{4}$	10512.5
2 $\frac{1}{2}$	1250.	5	5000.	7 $\frac{1}{2}$	11250.
2 $\frac{3}{4}$	1512.5	5 $\frac{1}{4}$	5512.5	7 $\frac{3}{4}$	12012.5
3	1800.	5 $\frac{1}{2}$	6050.	8	12800.
3 $\frac{1}{4}$	2112.5	5 $\frac{3}{4}$	6612.5		

STRENGTH OF METAL AND WOODEN RODS.

A rod having an area of the 1000th part of a square inch, made of the following materials, will sustain weights as follows:—

Designation.	Lbs.	Designation.	Lbs.
Cast steel.....	134	Tin.....	5
Best wrought iron....	70	Lead.....	2
Cast iron.....	19	Oak.....	12
Copper.....	19	Beach.....	12 $\frac{1}{2}$
Platinum.....	16	Ash.....	14
Silver.....	11	White Pine.....	11
Gold.....	9		

HEMPEN CORDS.

Hempen cords when twisted will support the following weights to the *square inch* of their section :—

Diameter.	Lbs.	Diameter.	Lbs.
$\frac{1}{4}$ to 1 inch.....	8746	3 to 5 inches.....	5345
1 to 3 inches.....	6800	5 to 7 inches.....	4860

NOTE.—A square inch of hemp fibres will support a weight of 9200 lbs.

The *maximum* strength of a good hemp rope is 6400 lbs. to the square inch. Its *practical* value not more than one-half this strain. Before breaking it stretches from $\frac{1}{8}$ to $\frac{1}{4}$, and its diameter diminishes from $\frac{1}{4}$ to $\frac{1}{8}$.

The strength of manilla is about $\frac{1}{2}$ that of hemp. White ropes are $\frac{1}{3}$ more durable.

LATERAL OR TRANSVERSE STRENGTH.

TABLE, showing the transverse strength of timber, 1 foot long and 1 inch square : Weight suspended from one end.

Materials.	Breaking weight. Lbs.	Greatest deflection. inches.	Weight borne with safety. Lbs.	Value for general use. Lbs.
White oak, seasoned	240.	9.	196.	40.
Chesnut, “	170.	1.8	115.	65.
Yellow pine, “	150.	1.7	100.	62.
White “	135.	1.4	95.	64.
Ash, “	175.	2.4	105.	77.
Hickory, “	270.	8.	200.	50.

TABLE, *showing the transverse strength of iron—square bar, 2 inches by 12 inches long: Weight suspended from one end.*

Material.	Breaking weight, Lbs.	Weight borne safely Lbs.	Value.	Value for general use.
Cast iron.	5781	4000	400	290

Round, 3 inches diameter by 12 inches long: Weight suspended from end.

Material.	Breaking weight, Lbs.	Weight borne with safety. Lbs.	Value.	Value for general use
Cast iron.	12000	8000	240	175

NOTE.—The strength of a projecting beam is only *one-fourth* of what it would be if supported at both ends, and only *one-sixth* of what it would be if *fixed* at both ends. The former is to the latter as 2 is to 3.

To find the transverse strength when the bar or beam is fixed at one end and the load applied at the other.

RULE.—Multiply the value in the preceding table by the breadth, and square of the depth in inches, and divide the product by the length in feet. The quotient will be the weight in lbs.

EXAMPLE.—What weight will a seasoned white oak beam 4 inches square and projecting 36 inches sustain?

SOLUTION.— $4 \times 4^2 \times 40 = 2560 \div 3$ feet, projection, $= 853\frac{1}{3}$ lbs. *Ans.*

EXAMPLE 2d.—What weight will a cast iron bar 2 inches square and projecting 4 feet sustain?

SOLUTION.— $2 \times 2^2 \times 400 = 3200 \div 4 = 800$ lbs. *Ans.*

NOTE.—When the beam is loaded uniformly throughout its length the result must be doubled.

When the bar or beam is fixed at both ends and the weight applied in the middle.

RULE.—Multiply the value in the preceding table by six times the breadth, and the square of the depth in inches, and divide the product by the length in feet.

EXAMPLE.—What weight will an ash beam 8 inches deep by 10 broad and 10 feet long sustain in the middle, when fixed at the ends?

SOLUTION.— $77 \times 60 \times 8^2 = 295680 \div 10 = 29568$ lbs. *Ans.*

EXAMPLE 2d.—What weight will a cast iron bar 2 inches square and 4 feet long support, when applied in the middle, the ends being fixed?

SOLUTION.— 400×12 , six times breadth, $\times 2^2 = 19200 \div 4 = 4800$ lbs. *Ans.*

NOTE.—When the weight is equally distributed along its entire length, the above results must be doubled.

When the bar or beam is supported at both ends and the weight applied in the middle.

RULE.—Multiply the value in the preceding table by the square of the depth, and four times the breadth in inches, and divide the product by the length in feet.

EXAMPLE.—What weight will a white pine beam 8 inches broad by 6 deep and 6 feet long carry when applied in the middle, the ends being supported?

SOLUTION.— $64 \times 6^2 \times 32 = 24576 \div 6 = 4129 + \text{lbs.}$ *Ans.*

EXAMPLE 2d.—What weight will a cast iron bar 2 inches square and 60 inches between the supports carry?

SOLUTION.— $400 \times 2^2 \times 8 = 12800 \div 5 \text{ feet} = 2560 \text{ lbs.}$ *Ans.*

TABLE, showing the resistance of materials to crushing.

Designation.	Crushing weight per square inch.	
	In lbs.	In tons of 2 000 lbs.
WOODS.		
Ash.....	8,683	4.3
Beech, well seasoned.....	19,363	9.6
Birch, " ".....	11,663	5.8
Cedar.....	5,863	2.9
Elder.....	9,973	4.9
Elm, well seasoned.....	10,331	5.1
Fir, (spruce,).....	6,819	3.4
Mahogany.....	8,198	4.09
Oak.....	5,982	2.9
Pine, pitch.....	6,790	3.3
" yellow.....	5,445	2.7
Poplar.....	5,124	2.5
Sycamore, highly seasoned.....	12,101	6.
Walnut.....	7,227	3.6
Willow.....	6,128	3.06
METALS.		
Brass, yellow.....	10,304	5.15
Iron, cast.....	98,000	49.
" bar.....	40,000	20.
" boiler plate.....	32,000	16.
MINERALS.		
Brick, common.....	800	0.40
" fire.....	1,700	0.85
Brickwork.....	612	0.306
Chalk.....	334	0.16
Granite.....	11,000	5.50

STRENGTH OF ICE.

Ice 2 inches thick will bear men on foot.

" 4 " " " " horseback.

Ice 6 inches thick will bear cattle and teams with light loads.

“ 8 “ “ “ teams with heavy loads.

“ 10 “ “ will sustain a pressure of 1000 lbs. per square foot.

This supposes the ice to be sound throughout its whole thickness, without “snow-ice.”

WEIGHT OF SQUARE ROLLED IRON.



From $\frac{1}{16}$ inch to 12 inches, and 1 foot in length.

Size in inches.	Weight in pounds.	Size in inches.	Weight in pounds.	Size in inches.	Weight in pounds.	Size in inches.	Weight in pounds.
$\frac{1}{16}$.013	2.	13.529	4.	64.700	7.	190.136
$\frac{1}{8}$.053	2.	15.263	4.	68.448	7.	203.024
$\frac{3}{16}$.118	2.	17.112	4.	72.305	8.	216.836
$\frac{1}{4}$.211	2.	19.066	4.	76.264	8.	230.068
$\frac{5}{16}$.475	2.	21.120	4.	80.333	8.	244.220
$\frac{3}{8}$.845	2.	23.292	5.	84.489	8.	258.800
$\frac{7}{16}$	1.320	2.	25.560	5.	88.784	9.	273.792
$\frac{1}{2}$	1.901	2.	27.939	5.	93.168	9.	289.220
$\frac{9}{16}$	2.588	3.	30.416	5.	97.657	9.	305.056
$\frac{5}{8}$	3.380	3.	33.010	5.	102.240	9.	321.332
$\frac{3}{4}$	4.278	3.	35.704	5.	106.953	10.	337.920
$1.$	5.280	3.	38.503	5.	111.756	10.	355.136
$1.$	6.390	3.	41.408	5.	116.671	10.	372.672
$1.$	7.604	3.	44.418	6.	121.664	10.	390.628
$1.$	8.926	3.	47.534	6.	132.040	11.	408.910
$1.$	10.352	4.	50.756	6.	142.816	11.	427.812
$1.$	11.883	4.	54.084	6.	154.012	11.	447.024
		4.	57.517	7.	165.632	11.	466.684
		4.	61.055	7.	177.672	12.	486.656

EXAMPLE.—What is the weight of a bar of rolled iron $1\frac{1}{2}$ inches square and 12 inches long?

In column 1st find $1\frac{1}{2}$, and opposite to it is 7.604 pounds, which is 7 lbs. and $\frac{604}{1000}$ of a lb. If the lesser denomination of ounces is required, the result is obtained as follows: Multiply the remainder by 16, pointing off the decimals as in multiplication of decimals, and the figures remaining on the left of the point indicate the number of ounces.

$$\begin{array}{r} \text{Thus, } \frac{604}{1000} \text{ of a lb.} = .604 \\ \phantom{\text{Thus, } \frac{604}{1000} \text{ of a lb.} = } 16 \\ \hline \phantom{\text{Thus, } \frac{604}{1000} \text{ of a lb.} = } 9.664 \text{ ounces.} \end{array}$$

The weight, then, is 7 lbs. $9.\frac{664}{1000}$ ounces.

If the weight for less than a foot in length was required, the readiest operation is this:

EXAMPLE.—What is the weight of a bar $6\frac{1}{4}$ inches square and $9\frac{3}{4}$ inches long?

In column 5th, opposite to $6\frac{1}{4}$, is 132.040, which is the weight for a foot in length.

$$\begin{array}{rcl} 6\frac{1}{4} \times 12 \text{ inches} & = & 132.040 \\ \hline 6. \text{ " is } \frac{1}{2} & = & 66.020 \\ 3. \text{ " is } \frac{1}{2} \text{ of } 6 & = & 33.010 \\ .\frac{1}{2} \text{ " is } \frac{1}{8} \text{ of } 3 & = & 6.5016 \\ .\frac{1}{4} \text{ " is } \frac{1}{2} \text{ of } \frac{1}{2} & = & 2.7508 \\ \hline 9.\frac{3}{4} & = & 108.\frac{2824}{10000} \text{ pounds.} \end{array}$$


WEIGHT OF ROUND ROLLED IRON.

From $\frac{1}{8}$ inch to 12 inches diameter, and 1 foot in length.

Diam't'r in inches	Weight in pounds.	Diam't'r in inches	Weight in pounds.	Diam't'r in inches	Weight in pounds.	Diam't'r in inches	Weight in pounds.
$\frac{1}{8}$	0.10	$2\frac{1}{8}$	11.988	$4\frac{1}{8}$	53.760	$7\frac{3}{4}$	159.456
$\frac{1}{4}$.041	$2\frac{1}{4}$	13.440	$4\frac{1}{4}$	56.788	8.	169.856
$\frac{3}{8}$.093	$2\frac{3}{8}$	14.975	$4\frac{3}{8}$	59.900	$8\frac{1}{4}$	180.696
$\frac{1}{2}$.165	$2\frac{1}{2}$	16.688	$4\frac{1}{2}$	63.094	$8\frac{3}{4}$	191.808
$\frac{5}{8}$.273	$2\frac{5}{8}$	18.293	5.	66.752	$8\frac{1}{2}$	203.260
$\frac{3}{4}$.463	$2\frac{3}{4}$	20.076	$5\frac{1}{8}$	69.731	9.	215.040
$\frac{7}{8}$.663	$2\frac{7}{8}$	21.944	$5\frac{1}{4}$	73.172	$9\frac{1}{4}$	227.152
1.	1.043	3.	23.888	$5\frac{3}{8}$	76.700	$9\frac{3}{4}$	239.600
$1\frac{1}{8}$	1.493	$3\frac{1}{8}$	25.926	$5\frac{1}{2}$	80.304	$9\frac{1}{2}$	252.376
$1\frac{1}{4}$	2.032	$3\frac{1}{4}$	28.040	$5\frac{3}{4}$	84.001	10.	266.288
$1\frac{1}{2}$	2.654	$3\frac{1}{2}$	30.240	$5\frac{7}{8}$	87.776	$10\frac{1}{4}$	278.924
$1\frac{3}{4}$	3.360	$3\frac{3}{4}$	32.512	6.	91.634	$10\frac{3}{4}$	282.988
$1\frac{7}{8}$	4.172	$3\frac{7}{8}$	34.886	$6\frac{1}{8}$	95.552	$10\frac{7}{8}$	306.800
2.	5.019	$4\frac{1}{8}$	37.332	$6\frac{1}{4}$	103.704	11.	321.216
$2\frac{1}{8}$	5.972	$4\frac{1}{4}$	39.864	$6\frac{3}{8}$	112.160	$11\frac{1}{4}$	336.004
$2\frac{1}{4}$	7.010	4.	42.464	$6\frac{1}{2}$	120.960	$11\frac{3}{4}$	351.104
$2\frac{3}{8}$	8.128	$4\frac{3}{8}$	45.174	7.	130.048	$11\frac{7}{8}$	366.536
$2\frac{1}{2}$	9.333	$4\frac{1}{2}$	47.952	$7\frac{1}{8}$	139.544	12.	382.208
$2\frac{3}{4}$	10.616	$4\frac{3}{4}$	50.815	$7\frac{1}{4}$	149.328		

The application of this table is precisely similar to that of the preceding one.

MASONRY.



A perch of stone is 24.75 cubic feet ; when built in the wall, 22 cubic feet make 1 perch, $2\frac{3}{4}$ cubic feet being allowed for the mortar and filling.

Three pecks of lime and four bushels of sand to a perch of wall.

To find the number of perches of stone in walls.

RULE.—Multiply the length in feet by the height in feet, and that by the thickness in feet, and divide the product by 22, and the quotient will be the number of perches of stone in the wall.

EXAMPLE.—How many perches of stone contained in a wall 40 feet long, 20 feet high, and 18 inches thick?

SOLUTION.—40 feet, length, \times 20 ft., height, \times $1\frac{1}{2}$ feet, thick, $=1200 \div 22 = 54.54$ perches. *Ans.*

NOTE.—To find the number of perches of *masonry*, divide the product, as above, by 24.75, instead of 22.

Brick-work.

The dimensions of common bricks are from $7\frac{3}{4}$ to 8 inches long, by $4\frac{1}{4}$ wide, and $2\frac{1}{2}$ thick. Front bricks are $8\frac{1}{4}$ inches long, by $4\frac{1}{2}$ wide, and $2\frac{1}{2}$ thick.

The usual size of fire bricks is $9\frac{1}{2}$ inches long, by $4\frac{3}{8}$ wide, by $2\frac{3}{8}$ thick.

Twenty common bricks to a cubic foot when laid; 15 common bricks to a foot of 8-inch wall when laid.

To find the number of common bricks in a wall.

RULE.—Multiply the length of the wall in feet by the height in feet, and that by its thickness in feet, and that again by 20, and the product will be the number of bricks in the wall.

EXAMPLE.—How many common bricks in a wall 40 feet long by 20 feet high and 12 inches thick?

SOLUTION.—40 ft., length, \times 20 ft., height, \times 1 ft., thick, \times 20 $=16000$. *Ans.*

NOTE.—For walls 8 ins. thick, multiply the length in feet by the height in feet, and that by 15, and the product will be the number of bricks in the wall.

When the wall is perforated by doors and windows, or other openings, find the sum of their cubic feet by severally multiplying their lengths and widths and thicknesses in feet together, and deducting the whole from the cubic contents of the wall, including the openings, before multiplying by 20 or 15, as above.

Laths.

Laths are $1\frac{1}{4}$ to $1\frac{1}{2}$ inches wide by 4 feet long, are usually set $\frac{1}{4}$ inch apart, and a bundle contains 100.

THE MECHANICAL POWERS.

The mechanical powers are three in number, namely: the LEVER, the INCLINED PLANE, and the PULLEY. The wheel and the axle is a *revolving* lever; the wedge is a *double inclined* plane, and the screw is a *revolving inclined* plane.

THE LEVER.

To find the length of the longest arm of the lever; the weight to be raised, the power to be applied, and the length of the shortest arm of the lever being given.

RULE.—Multiply the weight by its distance from the fulcrum and divide the product by the power, and the quotient is the distance from the fulcrum the power must be applied, or, the longest arm of the lever.

EXAMPLE.—Given, a weight of 900 lbs., distant 2 feet from the fulcrum, to be raised by a force or power of 75 lbs.; required, the length of the longest arm of the lever.

SOLUTION.—900 lbs., the weight, \times 2 feet, distance from fulcrum, $= 1800 \div 75$ lbs., the power, $= 24$ feet. *Ans.*

To find the length of the shortest arm of the lever; the weight to be raised, the power to be applied, and the length of the longest arm of the lever being given.

RULE.—Multiply the power by its distance from the fulcrum, and divide the product by the weight, and the quotient is the distance the weight must be placed from the fulcrum, or, the shortest arm of the lever.

EXAMPLE.—What distance must a weight of 800 lbs. be placed from the fulcrum, to be raised by a power of 150 lbs. placed 8 feet from the fulcrum?

SOLUTION.—150 lbs., the power, $\times 96$ inches, its distance from the fulcrum, $= 14400 \div 800$ lbs., the weight, $= 18$ inches.

Ans.

To find the power required to raise a given weight; the distances of the weight and the power from the fulcrum being given.

RULE.—Multiply the weight by its distance from the fulcrum and divide the product by the distance of the power from the fulcrum.

EXAMPLE.—What power will raise a weight of 600 lbs. 20 inches from the fulcrum, applied 8 feet from the fulcrum?

SOLUTION.—600 lbs., weight, $\times 20$ inches, distance of weight from fulcrum, $= 12000 \div 96$ inches, distance of power from fulcrum, $= 125$ lbs. *Ans.*

To find the weight, at a given distance from the fulcrum, a given power at a given distance from the fulcrum will raise.

RULE.—Multiply the power by its distance from the ful-

crum and divide the product by the distance of the weight from the fulcrum.

EXAMPLE.—What weight will a power of 250 lbs. 10 feet from the fulcrum raise, the weight placed 20 inches from the fulcrum?


SOLUTION.—250 lbs., the power, \times 120 inches, its distance from the fulcrum, $= 30000 \div 20$ inches, distance of weight from fulcrum, $= 1500$ lbs. *Ans.*

The GENERAL RULE, therefore, for ascertaining the relation of power to weight in a lever, is: the power applied, multiplied by its distance from the fulcrum, is equal to the weight multiplied by its distance from the fulcrum.

The pressure upon the fulcrum equals the sum of the weight and power.

NOTE.—It must be remembered that, according to the foregoing rules and examples, the weight and force are made by the introduction of the lever to equal or balance each other. Hence, to get at their *practical value*, we must either shorten the short arm, or lengthen the long arm of the lever, add to the power, or deduct from the weight, to such an extent as each may judge for himself expedient under the circumstances.

THE INCLINED PLANE.



To find the power or force required to raise a given weight up an inclined plane of a given length and height.

RULE.—As the length of the plane is to its height, so is the weight to the power.

EXAMPLE.—Required the power necessary to raise 1500 lbs. up an inclined plane 20 feet long and 8 feet high?

SOLUTION.—As $20 : 8 :: 1500 : 600$ lbs. *Ans.*

To find the height of an inclined plane when its length and base are given.

RULE.—Subtract the square of the base from the square of the length, and the square root of the remainder is the height.

EXAMPLE.—Given an inclined plane, the length of which is 40 feet and base 38: required, its height?

SOLUTION.— 1600 , square of length, — 1444 , square of base, = $\sqrt{156} = 12.49$ feet. *Ans.*

To find the length when its base and height are given.

RULE.—Add the squares of the height and the base, and the square root of their sum will be the length.

EXAMPLE.—What is the length of an inclined plane the base of which is 20 feet and its height 12?

SOLUTION.—400, square of base, + 144, square of height,
 $= \sqrt{544} = 23.32$ feet. *Ans.*

To find the base when the length and height are given.

RULE.—Subtract the square of the height from the square of the length, and the square root of the remainder will be the base.

EXAMPLE.—What is the base of an inclined plane, whose height is 10 feet, and length 25?

SOLUTION.—625, square of length, — 100, square of height, $= \sqrt{525} = 22.91$ feet. *Ans.*

To find the pressure of a weight on an inclined plane when raised by its equivalent power.

RULE.—As the length is to the weight, so is the base to the pressure.

EXAMPLE.—What is the pressure of 1000 lbs. on an inclined plane, the length of which is 80 feet and the base 70?

SOLUTION.—80 feet, length, : 1000 lbs., :: 70 feet, base,
 : 875 lbs. *Ans.*

NOTES.—When the line of direction of the power is parallel to the plane, the power is least and the pressure least.

When the power does not run parallel to the plane, draw a line perpendicular to the direction of the power's action from the end of the base line (at the back of the plane), and the intersection of this line on the length will determine the length and height of the base.

THE WHEEL AND THE AXLE.

The power multiplied by the radius of the wheel is equal to the weight multiplied by the radius of the axle.

As the radius of the wheel is to the radius of the axle, so is the effect to the power.

To find the weight a given tractile force or power will move on a wheel and axle of given radii.

RULE.—Multiply the tractile or drawing force by the radius of the wheel, and divide the product by the radius of the axle.

EXAMPLE.—What weight will a tractile force of 250 lbs. draw on a wheel (or wheels) of a radius of 3 feet: radius of axle 4 inches?

SOLUTION.—250 lbs., tractile force, \times 36 inches, radius of wheel, $= 9000 \div 4$ inches, radius of axle, $= 2250$ lbs. *Ans.*

To find the tractile force required to move a given weight on a wheel and axle of given radii.

RULE.—Multiply the weight by the radius of the axle and divide the product by the radius of the wheel.

EXAMPLE.—Required, the tractile force necessary to draw 2000 lbs. on a wheel of $2\frac{1}{2}$ feet radius, and axle of 3 inches radius?

SOLUTION.—2000 lbs., weight, \times 3 inches, radius of axle, $= 6000 \div 30$ inches, radius of wheel, $= 200$ lbs. *Ans.*

To find the radius required for a wheel to move a given weight by a given force on a given radius of axle.

RULE.—Multiply the weight by the radius of the axle and divide the product by the force.

EXAMPLE.—What radius must a wheel have, the radius of whose axle is 4 inches, to move a weight of 1320 lbs. by a force of 220 lbs?

SOLUTION.—1320 lbs., weight, \times 4 inches, radius of axle, $= 5280 \div 220$ lbs., tractile force, $= 24$ inches. *Ans.*

To find the radius of an axle required to move a given weight by a given force, on a wheel of a given radius.


RULE.—Multiply the force by the radius of the wheel and divide the product by the weight.

EXAMPLE.—A weight of 1200 lbs. is to be moved on a wheel of 4 feet radius by a force of 150 lbs.: What must be the radius of the axle?

SOLUTION.—150 lbs., force, \times 48 inches, radius of wheel, $= 7200 \div 1200$ lbs. $= 6$ inches. *Ans.*

NOTE.—It will be observed that, according to the above rules, illustrated by the foregoing examples, the power or force of traction and the weight or load are equivalents; that is to say, the one is, by the interposition of the wheel and axle, made to counterpoise the other. To find their easy *practical value*, deduct $\frac{1}{4}$ from the weight, or add $\frac{1}{8}$ to the tractile force.

THE WEDGE.



To find the force necessary to separate two bodies from one another in a direction parallel to the back of the wedge.

RULE.—As the length of the wedge is to half its back, so is the resistance to the force.

EXAMPLE.—The length of the back of a double wedge is 6 inches, and its length through the middle 12 inches. Required, the force necessary to separate a substance having a resistance of 200 lbs. ?

SOLUTION.—12 inches, length, : 3 inches, back, :: 200 lbs., resistance, : 50 lbs. *Ans.*

To find the requisite force when only one of the bodies is movable.


RULE.—As the length of the wedge is to its back, so is the resistance to the force.

EXAMPLE.—What power applied to the back of a wedge will raise a weight of 20,000 lbs.; the wedge being 6 inches deep and 100 long on its base?

SOLUTION.—100 inches, length, : 6 inches, depth, :: 20,000 lbs., weight, : 1200 lbs. *Ans.*

NOTE.—The power of the wedge increases as its length increases, or as the thickness of its back decreases.

THE SCREW.



The screw is a revolving inclined plane, or an inclined plane wound round a cylinder. Hence, when its length and its pitch, or height, are ascertained, the same rules that govern the inclined plane apply to the screw.

To find the length of the inclined plane of a screw.

RULE.—Add the square of the circumference of the screw to the square of the pitch, or distance between the threads, and take the square root of the same, which will be the length of the plane. The height is the distance between the consecutive threads.

EXAMPLE.—What is the length of the inclined plane of a screw of 12 inches circumference and 1 inch pitch?

SOLUTION.— $12^2 + 1^2 = 145$ and $\sqrt{145} = 12.04159$ inches.

Ans.

NOTE.—It will be observed that the length of the plane as given in the above example is the length of only *one turning of the screw*, or the length of once round the circumference, which, in ascertaining the power of a screw, is all that is necessary to be known of the length. The entire length of the plane and the entire height of the screw have nothing to do with its power. A single section, comprising one revolution of the plane or the cylinder, is enough.

To find the power required to raise a given weight by means of a screw of given dimensions.

RULE.—As the length of the inclined plane is to the pitch, or height of it, so is the weight to the power.

EXAMPLE.—What is the power requisite to raise 9000 lbs. by a screw 15 inches circumference, and $1\frac{1}{2}$ inches pitch?


SOLUTION.— $15^2 + 1\frac{1}{2}^2 = 227\frac{1}{4}$ and $\sqrt{227\frac{1}{4}} = 15.62$ inches, length, then 15.62 inches, length, : $1\frac{1}{2}$ inches, pitch, :: 9000 lbs., weight, : 864.27 lbs. Ans.

NOTE.—When a wheel or capstan is applied to turn the screw, the length of the lever is the radius of the circle described by the handle of the wheel or capstan bar, and half the diameter of the screw is the radius of the axle.

When the screw is turned by a wheel, a crank, or capstan,

find the power of the wheel, crank, or capstan by means of the rules given under "The Wheel and the Axle," and deduct the force thus acquired from the force necessary to drive the screw in raising the weight alone. The remainder is the force required to raise the weight by the combined power of the screw and the lever.

THE PULLEY.



When only one cord or rope is used.

To find the force necessary to raise a given weight by means of a pulley of a given number of sheaves, &c.

RULE.—Divide the weight to be raised by the number of parts of the rope engaged in supporting the lower or movable block.

EXAMPLE.—What is the force required to raise 600 lbs. by means of a lower block containing six sheaves: rope fastened to the upper block?

SOLUTION.— $2 \times 6 = 12$; then, $600 \div 12 = 50$ lbs. *Ans.*

EXAMPLE 2d.—What force when the rope is fastened to the lower block?

SOLUTION.— $6 \times 2 + 1 = 13$; then $600 \div 13 = 46.16$ lbs. *Ans.*


When more than one rope is used.

In a *Spanish Burton*, where there are two ropes, two movable pulleys, and one fixed and one stationary pulley, with the ends of one rope fastened to the support and upper movable pulley, and the ends of the other fastened to the lower block and the power, the weight is to the power as 5 to 1.

In one where the ends of one rope are fastened to the support and the power, and the ends of the other to the lower and upper blocks, the weight is to the power as 4 to 1.


DEFINITIONS OF MATHEMATICAL FORMS.

FIG. 1.



Parallel Lines are everywhere equally distant; as, A B and C D.

FIG. 2.




An *Angle* is the difference of direction between two lines which meet; as, A D E. The point of meeting is called the vertex of the angle, and when the angle is named the letter at the vertex is placed second; as, C D E.

FIG. 3.



A *Right Angle* is formed when a straight line meeting another makes two equal angles; as, A D C and C D B.

FIG. 4.




An *Acute Angle* is one less than a right angle; as, E B D, Fig. 3.

An *Obtuse Angle** is one greater than a right angle; as, A D E, Fig. 4.

A *Surface* has two dimensions—length and breadth.

A *Triangle* is a figure having three sides; as, A B C, Fig. 5.

FIG. 5.



The *Altitude* of a triangle is the perpendicular distance of the vertex from the line of the base; as, B C is the altitude of the triangle A B C, Fig. 5.

A *Right-Angle Triangle* is a triangle having a right angle;

* As the right angle contains 90° , it follows that the acute angle contains less, and the obtuse angle more, than 90° .

as, A C B, Fig. 5. The side opposite the right angle is called the hypotenuse; as, A B.

A *Parallelogram* is a four-sided figure whose opposite sides are parallel; as, Fig. 6.

FIG. 6.




A *Rectangle* is a parallelogram whose angles are right angles; as, Fig. 7.

FIG. 7.



A *Square* is a rectangle the sides of which are equal. Fig. 8.


FIG. 8.



A *Trapezoid* is a four-sided figure having but two of its sides parallel; as, A B C D, Fig. 9.

The *Altitude* of a Parallelogram, a Rectangle, a Square or a Trapezoid is the perpendicular distance between the base and the line of the parallel side opposite the base; as, E F, Fig. 9.


FIG. 9.



A *Circle* is a plane surface bounded by a line, every point of which is equally distant from a point called the centre; as, A B C D, Fig. 10.

The *Circumference* of a circle is the line by which it is bounded; as, A B C D, Fig. 10.

FIG. 10.



The *Diameter* of a circle is a straight line passing through the centre and terminating in the circumference; as, D E B, Fig. 10.

The *Radius* of a circle is the distance from the centre to the circumference; as, E F.

FIG. 11.



A *Solid* has three dimensions—length, breadth, and thickness ; as, Fig. 11.

FIG. 12.



A *Prism* is a solid whose sides are parallelograms, and whose ends are equal and similar ; as, Fig. 12.

FIG. 13.




When the ends of a prism are triangular, it is called a *triangular prism* ; as, Fig. 12.

FIG. 14.



When the ends of a prism are square, it is called a *square prism* ; as, Fig. 13.

FIG. 15.



When the ends of a prism are hexagonal, it is called a *hexagonal prism* ; as, Fig. 14.

FIG. 16.



When the ends of a prism are circular, it is called a *cylinder* ;* as, Fig. 15.

FIG. 17.



When all the sides of a rectangular prism are square, it is called a *cube* ; as, Fig. 16.

A *Pyramid* is a solid, the base of which is a plane rectilinear figure, and having sides which are triangles whose vertices meet at a point at the top called the vertex of the pyramid. Fig. 17.

FIG. 18.



The *Altitude* of a pyramid or a cone is the perpendicular distance from the vertex to the plane of the base ; as, Fig. 17.

A *Cone* is a solid, the base of which is a circle, and which tapers uniformly to a point at the top called a vertex. Fig. 18.

* A cylinder is a regular polygon, or prism, with an infinite number of sides.

A *Frustum* of a pyramid or a cone is the part that remains after cutting off the top by a plane parallel to the base.

Fig. 19 represents the frustum of a pyramid.

Fig. 20 represents the frustum of a cone.

An *Ellipse* is a plane curve such that the sums of the distances of any points in the bounding line from two points within called the foci are always equal.

The line A B passing through the foci is called the major diameter; and the diameter perpendicular to A B at its centre is called the minor diameter.

A *Sphere* is a solid, bounded by a convex surface, every point of which is equally distant from a point within called the centre; as, Fig. 22.

A *Spheroid* is a solid, generated by the revolution of an ellipse about one of its diameters. If the ellipse revolves about its major diameter the spheroid is called *prolate*. If about its minor diameter the spheroid is called *oblate*.

Fig. 23 represents a *prolate spheroid*.

Fig. 24 represents an *oblate spheroid*.

FIG. 19



FIG. 20.



FIG. 21.



FIG. 22.




FIG. 23.



FIG. 24.



$$7 : 22 = c$$

$$\begin{array}{r} 2 \\ 7 \overline{) 14} \\ \underline{14} \\ 0 \end{array}$$

$$20 \setminus 2 = 60 \setminus 8 = 40 \quad [5]$$

CIRCLES.

To find the circumference of a circle.

RULE 1.—Multiply the diameter by 3.1416, and the product will be the circumference.

RULE 2.—Or, as 7 is to 22 so is the diameter to the circumference.

EXAMPLE.—What is the circumference of a circle whose diameter is 25 ?

SOLUTION.— $25 \times 3.1416 = 78.54$. *Ans.* By Rule 2.— $7 : 22 :: 25 : 78.5$. *Ans.*

To find the diameter of a circle.

RULE 1.—Divide the circumference by 3.1416, and the quotient will be the diameter.

RULE 2.—Or, as 22 is to 7, so is the circumference to the diameter.

EXAMPLE.—What is the diameter of a circle whose circumference is 69.11 ?

SOLUTION.— $69.11 \div 3.1416 = 22$. *Ans.* By Rule 2.— $22 : 7 :: 69.11 : 22$. *Ans.*

To find the area of a circle.

RULE 1.—Multiply the square of the diameter by .7854, or the square of the circumference by .07958, and the product will be the area.

RULE 2.—Or, multiply half the circumference by half the diameter.

RULE 3.—Or, as 14 is to 11, so is the square of the diameter to the area.

RULE 4.—Or, as 88 is to 7, so is the square of the circumference to the area.

To find the side of an equal square containing the same area as a given circle.

RULE.—Take the square root of the area, which will be the side of the equal square.

To find the solidity of a sphere.

RULE.—Multiply the cube of the diameter by .5236, and the product is the solidity.

EXPLANATION AND USE OF THE FOLLOWING TABLE.

In the left hand column will be found the diameter of the circle; in the next column to the right will be found its corresponding circumference; in the third to the right will be found the area, and in the right hand column will be found the length of the side of an equal square containing the same area.

EXAMPLE.—What is the side of a square having the same area as a circle of $64\frac{1}{2}$ inches diameter?

SOLUTION.—Find $64\frac{1}{2}$ in the left-hand column, and opposite it to the right, under the heading "*Side of Equal Square,*" will be found 57.101, the length of the side. *Ans.*

TABLE, showing the Areas of Circles and the Sides of their equivalent Squares, from 1 to 100.

Diam.	Circum.	Area.	Side of equal square.	Diam.	Circum.	Area.	Side of equal square.
$1\frac{1}{2}$.3926	.01227	.110	$11\frac{1}{2}$	36.91	108.43	10.413
$1\frac{3}{4}$.7854	.04908	.221	12	37.69	113.09	10.634
$1\frac{5}{4}$	1.570	.1963	.443	$12\frac{1}{2}$	38.48	117.85	10.856
$1\frac{3}{2}$	2.356	.4417	.663	$12\frac{3}{4}$	39.27	122.71	11.077
1	3.141	.7854	.886	$12\frac{5}{4}$	40.05	127.67	11.299
$1\frac{1}{4}$	3.927	1.227	1.107	13	40.84	132.73	11.520
$1\frac{1}{2}$	4.712	1.767	1.329	$13\frac{1}{4}$	41.62	137.88	11.742
$1\frac{3}{4}$	5.497	2.404	1.550	$13\frac{3}{4}$	42.41	143.13	11.964
2	6.283	3.141	1.772	$13\frac{5}{4}$	43.19	148.48	12.185
$2\frac{1}{4}$	7.068	3.976	1.994	14	43.98	153.93	12.407
$2\frac{1}{2}$	7.854	4.908	2.215	$14\frac{1}{4}$	44.76	159.48	12.628
$2\frac{3}{4}$	8.639	5.939	2.437	$14\frac{3}{4}$	45.55	165.13	12.850
3	9.424	7.068	2.658	$14\frac{5}{4}$	46.33	170.87	13.071
$3\frac{1}{4}$	10.21	8.295	2.880	15	47.12	176.71	13.293
$3\frac{1}{2}$	10.99	9.621	3.101	$15\frac{1}{4}$	47.90	182.65	13.514
$3\frac{3}{4}$	11.78	11.044	3.323	$15\frac{3}{4}$	48.69	188.69	13.736
4	12.56	12.566	3.544	$15\frac{5}{4}$	49.48	194.82	13.958
$4\frac{1}{4}$	13.35	14.186	3.766	16	50.26	201.06	14.179
$4\frac{1}{2}$	14.13	15.904	3.988	$16\frac{1}{4}$	51.05	207.39	14.401
$4\frac{3}{4}$	14.92	17.720	4.209	$16\frac{3}{4}$	51.83	213.82	14.622
5	15.70	19.635	4.431	$16\frac{5}{4}$	52.62	220.35	14.844
$5\frac{1}{4}$	16.49	21.647	4.652	17	53.40	226.98	15.065
$5\frac{1}{2}$	17.27	23.758	4.874	$17\frac{1}{4}$	54.19	233.70	15.287
$5\frac{3}{4}$	18.06	25.967	5.095	$17\frac{3}{4}$	54.97	240.52	15.508
6	18.84	28.274	5.317	$17\frac{5}{4}$	55.76	247.45	15.730
$6\frac{1}{4}$	19.63	30.679	5.538	18	56.54	254.46	15.952
$6\frac{1}{2}$	20.42	33.183	5.760	$18\frac{1}{4}$	57.33	261.58	16.173
$6\frac{3}{4}$	21.20	35.784	5.982	$18\frac{3}{4}$	58.11	268.80	16.395
7	21.99	38.484	6.203	$18\frac{5}{4}$	58.90	276.11	16.616
$7\frac{1}{4}$	22.77	41.282	6.425	19	59.69	283.52	16.838
$7\frac{1}{2}$	23.56	44.178	6.646	$19\frac{1}{4}$	60.47	291.03	17.059
$7\frac{3}{4}$	24.34	47.173	6.868	$19\frac{3}{4}$	61.26	298.64	17.281
8	25.13	50.265	7.089	$19\frac{5}{4}$	62.04	306.35	17.502
$8\frac{1}{4}$	25.91	53.456	7.311	20	62.83	314.16	17.724
$8\frac{1}{2}$	26.70	56.745	7.532	$20\frac{1}{4}$	63.61	322.06	17.946
$8\frac{3}{4}$	27.48	60.131	7.754	$20\frac{3}{4}$	64.40	330.06	18.167
9	28.27	63.617	7.976	$20\frac{5}{4}$	65.18	338.16	18.389
$9\frac{1}{4}$	29.05	67.200	8.197	21	65.97	346.36	18.610
$9\frac{1}{2}$	29.84	70.882	8.419	$21\frac{1}{4}$	66.75	354.65	18.832
$9\frac{3}{4}$	30.63	74.662	8.640	$21\frac{3}{4}$	67.54	363.05	19.053
10	31.41	78.539	8.862	$21\frac{5}{4}$	68.32	371.54	19.275
$10\frac{1}{4}$	32.20	82.516	9.083	22	69.11	380.13	19.496
$10\frac{1}{2}$	32.98	86.590	9.305	$22\frac{1}{4}$	69.90	388.82	19.718
$10\frac{3}{4}$	33.77	90.762	9.526	$22\frac{3}{4}$	70.68	397.60	19.940
11	34.55	95.033	9.748	$22\frac{5}{4}$	71.47	406.49	20.161
$11\frac{1}{4}$	35.34	99.402	9.970	23	72.25	415.47	20.383
$11\frac{1}{2}$	36.12	103.86	10.191	$23\frac{1}{4}$	73.04	424.55	20.604

Diam.	Circum.	Area.	Side of equal square.	Diam.	Circum.	Area.	Side of equal square.
23 $\frac{1}{2}$	73.82	433.73	20.826	36	113.	1017.8	31.504
23 $\frac{3}{4}$	74.61	443.01	21.047	36 $\frac{1}{4}$	113.8	1032.0	32.125
24	75.39	452.39	21.269	36 $\frac{1}{2}$	114.6	1046.3	32.347
24 $\frac{1}{4}$	76.18	461.86	21.491	36 $\frac{3}{4}$	115.4	1060.7	32.568
24 $\frac{1}{2}$	76.96	471.43	21.712	37	116.2	1075.2	32.790
24 $\frac{3}{4}$	77.75	481.10	21.934	37 $\frac{1}{4}$	117.	1089.7	33.011
25	78.54	490.87	22.155	37 $\frac{1}{2}$	117.8	1104.4	33.233
25 $\frac{1}{4}$	79.32	500.74	22.377	37 $\frac{3}{4}$	118.6	1119.2	33.455
25 $\frac{1}{2}$	80.10	510.70	22.598	38	119.3	1134.1	33.676
25 $\frac{3}{4}$	81.89	520.77	22.820	38 $\frac{1}{4}$	120.1	1149.0	33.898
26	81.68	530.93	23.041	38 $\frac{1}{2}$	120.9	1164.1	34.119
26 $\frac{1}{4}$	82.46	541.18	23.263	38 $\frac{3}{4}$	121.7	1179.3	34.341
26 $\frac{1}{2}$	83.25	551.54	23.485	39	122.5	1194.5	34.562
26 $\frac{3}{4}$	84.03	562.00	23.706	39 $\frac{1}{4}$	123.3	1209.9	34.784
27	84.82	572.55	23.928	39 $\frac{1}{2}$	124.	1225.4	35.005
27 $\frac{1}{4}$	85.60	583.20	24.149	39 $\frac{3}{4}$	124.8	1240.9	35.227
27 $\frac{1}{2}$	86.39	593.95	24.371	40	125.6	1256.6	35.449
27 $\frac{3}{4}$	87.17	604.80	24.592	40 $\frac{1}{4}$	126.4	1272.3	35.670
28	87.96	615.75	24.814	40 $\frac{1}{2}$	127.2	1288.2	35.892
28 $\frac{1}{4}$	88.75	626.79	25.035	40 $\frac{3}{4}$	128.	1304.2	36.113
28 $\frac{1}{2}$	89.53	637.94	25.257	41	128.8	1320.2	36.335
28 $\frac{3}{4}$	90.32	649.18	25.479	41 $\frac{1}{4}$	129.5	1336.4	36.556
29	91.10	660.52	25.700	41 $\frac{1}{2}$	130.3	1352.6	36.778
29 $\frac{1}{4}$	91.89	671.95	25.922	41 $\frac{3}{4}$	131.1	1369.0	36.999
29 $\frac{1}{2}$	92.67	683.49	26.144	42	131.9	1385.4	37.221
29 $\frac{3}{4}$	93.46	695.12	26.365	42 $\frac{1}{4}$	132.7	1401.9	37.443
30	94.24	706.86	26.586	42 $\frac{1}{2}$	133.5	1418.6	37.664
30 $\frac{1}{4}$	95.03	718.69	26.808	42 $\frac{3}{4}$	134.3	1435.3	37.886
30 $\frac{1}{2}$	95.81	730.61	27.029	43	135.	1452.2	38.107
30 $\frac{3}{4}$	96.60	742.64	27.251	43 $\frac{1}{4}$	135.8	1469.1	38.329
31	97.38	754.76	27.473	43 $\frac{1}{2}$	136.6	1486.1	38.550
31 $\frac{1}{4}$	98.17	766.99	27.694	43 $\frac{3}{4}$	137.4	1503.3	38.772
31 $\frac{1}{2}$	98.97	779.31	27.916	44	138.2	1520.5	38.993
31 $\frac{3}{4}$	99.74	791.73	28.137	44 $\frac{1}{4}$	139.	1537.8	39.215
32	100.0	804.24	28.359	44 $\frac{1}{2}$	139.8	1555.2	39.437
32 $\frac{1}{4}$	101.3	816.86	28.580	44 $\frac{3}{4}$	140.5	1572.8	39.658
32 $\frac{1}{2}$	102.1	829.57	28.802	45	141.3	1590.4	39.880
32 $\frac{3}{4}$	102.8	842.39	29.023	45 $\frac{1}{4}$	142.1	1608.1	40.101
33	103.6	855.30	29.245	45 $\frac{1}{2}$	142.9	1625.9	40.323
33 $\frac{1}{4}$	104.4	868.30	29.467	45 $\frac{3}{4}$	143.7	1643.8	40.544
33 $\frac{1}{2}$	105.2	881.41	29.688	46	144.5	1661.9	40.766
33 $\frac{3}{4}$	106.	894.61	29.910	46 $\frac{1}{4}$	145.2	1680.0	40.987
34	106.8	907.92	30.131	46 $\frac{1}{2}$	146.	1698.2	41.209
34 $\frac{1}{4}$	107.5	921.32	30.353	46 $\frac{3}{4}$	146.8	1716.5	41.431
34 $\frac{1}{2}$	108.3	934.82	30.574	47	147.6	1734.9	41.652
34 $\frac{3}{4}$	109.1	948.41	30.796	47 $\frac{1}{4}$	148.4	1753.4	41.874
35	109.9	962.11	31.017	47 $\frac{1}{2}$	149.2	1772.0	42.095
35 $\frac{1}{4}$	110.7	975.90	31.239	47 $\frac{3}{4}$	150.	1790.7	42.317
35 $\frac{1}{2}$	111.5	989.80	31.461	48	150.7	1809.5	42.538
35 $\frac{3}{4}$	112.3	1003.7	31.682	48 $\frac{1}{4}$	151.5	1828.4	42.760

Diam.	Circum.	Area.	Side of equal square.	Diam.	Circum.	Area.	Side of equal square.
48 $\frac{1}{4}$	152.3	1847.4	42.982	61	191.6	2922.4	54.059
48 $\frac{3}{4}$	153.1	1866.5	43.203	61 $\frac{1}{4}$	192.4	2946.4	54.281
49	153.9	1885.7	43.425	61 $\frac{1}{2}$	193.2	2970.5	54.502
49 $\frac{1}{4}$	154.7	1905.	43.646	61 $\frac{3}{4}$	193.9	2994.7	54.724
49 $\frac{1}{2}$	155.5	1924.4	43.868	62	194.7	3019.0	54.946
49 $\frac{3}{4}$	156.2	1943.9	44.089	62 $\frac{1}{4}$	195.5	3043.4	55.167
50	157.	1963.5	44.311	62 $\frac{1}{2}$	196.3	3067.9	55.389
50 $\frac{1}{4}$	157.8	1983.1	44.532	62 $\frac{3}{4}$	197.1	3092.5	55.610
50 $\frac{1}{2}$	158.6	2002.9	44.754	63	197.9	3117.2	55.832
50 $\frac{3}{4}$	159.4	2022.8	44.976	63 $\frac{1}{4}$	198.7	3142.0	56.053
51	160.2	2042.8	45.197	63 $\frac{1}{2}$	199.4	3166.9	56.275
51 $\frac{1}{4}$	161.	2062.9	45.419	63 $\frac{3}{4}$	200.2	3191.9	56.496
51 $\frac{1}{2}$	161.7	2083.0	45.640	64	201.	3216.9	56.718
51 $\frac{3}{4}$	162.5	2103.3	45.862	64 $\frac{1}{4}$	201.8	3242.1	56.940
52	163.3	2123.7	46.083	64 $\frac{1}{2}$	202.6	3267.4	57.161
52 $\frac{1}{4}$	164.1	2144.1	46.305	64 $\frac{3}{4}$	203.4	3292.8	57.383
52 $\frac{1}{2}$	164.9	2164.7	46.526	65	204.2	3318.3	57.604
52 $\frac{3}{4}$	165.7	2185.4	46.748	65 $\frac{1}{4}$	204.9	3343.8	57.826
53	166.5	2206.1	46.970	65 $\frac{1}{2}$	205.7	3369.5	58.047
53 $\frac{1}{4}$	167.2	2227.0	47.191	65 $\frac{3}{4}$	206.5	3395.3	58.269
53 $\frac{1}{2}$	168.	2248.0	47.413	66	207.3	3421.2	58.490
53 $\frac{3}{4}$	168.8	2269.0	47.634	66 $\frac{1}{4}$	208.1	3447.1	58.712
54	169.6	2290.2	47.856	66 $\frac{1}{2}$	208.9	3473.2	58.934
54 $\frac{1}{4}$	170.4	2311.4	48.077	66 $\frac{3}{4}$	209.7	3499.3	59.155
54 $\frac{1}{2}$	171.2	2332.8	48.299	67	210.4	3525.6	59.377
54 $\frac{3}{4}$	172.	2354.2	48.520	67 $\frac{1}{4}$	211.2	3552.0	59.598
55	172.7	2375.8	48.742	67 $\frac{1}{2}$	212.	3578.4	59.820
55 $\frac{1}{4}$	173.5	2397.4	48.964	67 $\frac{3}{4}$	212.8	3605.0	60.041
55 $\frac{1}{2}$	174.3	2419.2	49.185	68	213.6	3631.6	60.263
55 $\frac{3}{4}$	175.1	2441.0	49.407	68 $\frac{1}{4}$	214.4	3658.4	60.484
56	175.9	2463.0	49.628	68 $\frac{1}{2}$	215.1	3685.2	60.706
56 $\frac{1}{4}$	176.7	2485.0	49.850	68 $\frac{3}{4}$	215.9	3712.2	60.928
56 $\frac{1}{2}$	177.5	2507.1	50.071	69	216.7	3739.2	61.149
56 $\frac{3}{4}$	178.2	2529.4	50.293	69 $\frac{1}{4}$	217.5	3766.4	61.371
57	179.	2551.7	50.514	69 $\frac{1}{2}$	218.3	3793.6	61.592
57 $\frac{1}{4}$	179.8	2574.1	50.736	69 $\frac{3}{4}$	219.1	3821.0	61.814
57 $\frac{1}{2}$	180.6	2596.7	50.958	70	219.9	3848.4	62.035
57 $\frac{3}{4}$	181.4	2619.3	51.179	70 $\frac{1}{4}$	220.6	3875.9	62.257
58	182.2	2642.0	51.401	70 $\frac{1}{2}$	221.4	3903.6	62.478
58 $\frac{1}{4}$	182.9	2664.9	51.622	70 $\frac{3}{4}$	222.2	3931.3	62.700
58 $\frac{1}{2}$	183.7	2687.8	51.844	71	223.	3959.2	62.922
58 $\frac{3}{4}$	184.5	2710.8	52.065	71 $\frac{1}{4}$	223.8	3987.1	63.143
59	185.3	2733.9	52.287	71 $\frac{1}{2}$	224.6	4015.1	63.365
59 $\frac{1}{4}$	186.1	2757.1	52.508	71 $\frac{3}{4}$	225.4	4043.2	63.586
59 $\frac{1}{2}$	186.9	2780.5	52.730	72	226.1	4071.5	63.808
59 $\frac{3}{4}$	187.7	2803.9	52.952	72 $\frac{1}{4}$	226.9	4099.8	64.029
60	188.4	2827.4	53.173	72 $\frac{1}{2}$	227.7	4128.2	64.251
60 $\frac{1}{4}$	189.2	2851.0	53.395	72 $\frac{3}{4}$	228.5	4156.7	64.473
60 $\frac{1}{2}$	190.	2874.7	53.616	73	229.3	4185.3	64.694
60 $\frac{3}{4}$	190.8	2898.5	53.838	73 $\frac{1}{4}$	230.1	4214.1	64.916

Diam.	Circum.	Area.	Side of equal square.	Diam.	Circum.	Area.	Side of equal square.
73 $\frac{1}{2}$	230.9	4242.9	65.137	86	270.1	5808.8	76.215
73 $\frac{3}{4}$	231.6	4271.8	65.359	86 $\frac{1}{2}$	270.9	5842.6	76.437
74	232.4	4300.8	65.580	86 $\frac{3}{4}$	271.7	5876.5	76.658
74 $\frac{1}{4}$	233.2	4329.9	65.802	86 $\frac{3}{4}$	272.5	5910.5	76.880
74 $\frac{1}{2}$	234.	4359.1	66.023	87	273.3	5944.6	77.101
74 $\frac{3}{4}$	234.8	4388.4	66.245	87 $\frac{1}{4}$	274.1	5978.9	77.323
75	235.6	4417.8	66.467	87 $\frac{1}{2}$	274.8	6013.2	77.544
75 $\frac{1}{4}$	236.4	4447.3	66.688	87 $\frac{3}{4}$	275.6	6047.6	77.766
75 $\frac{1}{2}$	237.1	4476.9	66.910	88	276.4	6082.1	77.987
75 $\frac{3}{4}$	237.9	4506.6	67.191	88 $\frac{1}{4}$	277.2	6116.7	78.209
76	238.7	4536.4	67.353	88 $\frac{1}{2}$	278.	6151.4	78.431
76 $\frac{1}{4}$	239.5	4566.3	67.574	88 $\frac{3}{4}$	278.8	6186.2	78.652
76 $\frac{1}{2}$	240.3	4596.3	67.796	89	279.6	6221.1	78.874
76 $\frac{3}{4}$	241.1	4626.4	68.017	89 $\frac{1}{4}$	280.3	6256.1	79.095
77	241.9	4656.6	68.239	89 $\frac{1}{2}$	281.1	6291.2	79.317
77 $\frac{1}{4}$	242.6	4686.9	68.461	89 $\frac{3}{4}$	281.9	6326.4	79.538
77 $\frac{1}{2}$	243.4	4717.3	68.682	90	282.7	6361.7	79.760
77 $\frac{3}{4}$	244.2	4747.7	68.904	90 $\frac{1}{4}$	283.5	6397.1	79.981
78	245.	4778.3	69.125	90 $\frac{1}{2}$	284.3	6432.6	80.203
78 $\frac{1}{4}$	245.8	4809.0	69.347	90 $\frac{3}{4}$	285.1	6468.2	80.425
78 $\frac{1}{2}$	246.6	4839.8	69.568	91	285.8	6503.8	80.646
78 $\frac{3}{4}$	247.4	4870.7	69.790	91 $\frac{1}{4}$	286.6	6539.6	80.868
79	248.1	4901.6	70.011	91 $\frac{1}{2}$	287.4	6575.5	81.089
79 $\frac{1}{4}$	248.9	4932.7	70.233	91 $\frac{3}{4}$	288.2	6611.5	81.311
79 $\frac{1}{2}$	249.7	4963.9	70.455	92	289.	6647.6	81.532
79 $\frac{3}{4}$	250.5	4995.1	70.676	92 $\frac{1}{4}$	289.8	6683.8	81.754
80	251.3	5026.5	70.898	92 $\frac{1}{2}$	290.5	6720.0	81.975
80 $\frac{1}{4}$	252.1	5058.0	71.119	92 $\frac{3}{4}$	291.3	6756.4	82.197
80 $\frac{1}{2}$	252.8	5089.5	71.341	93	292.1	6792.9	82.419
80 $\frac{3}{4}$	253.6	5121.2	71.562	93 $\frac{1}{4}$	292.9	6829.4	82.640
81	254.4	5153.0	71.784	93 $\frac{1}{2}$	293.7	6866.1	82.862
81 $\frac{1}{4}$	255.2	5184.8	72.005	93 $\frac{3}{4}$	294.5	6902.9	83.083
81 $\frac{1}{2}$	256.	5216.8	72.227	94	295.3	6939.7	83.305
81 $\frac{3}{4}$	256.8	5248.8	72.449	94 $\frac{1}{4}$	296.	6776.7	83.526
82	257.6	5281.0	72.670	94 $\frac{1}{2}$	296.8	7013.8	83.748
82 $\frac{1}{4}$	258.3	5313.2	72.892	94 $\frac{3}{4}$	297.6	7050.9	83.970
82 $\frac{1}{2}$	259.1	5345.6	73.113	95	298.4	7088.2	84.191
82 $\frac{3}{4}$	259.9	5378.0	73.335	95 $\frac{1}{4}$	299.2	7122.5	84.413
83	260.7	5410.6	73.566	95 $\frac{1}{2}$	300.	7163.0	84.634
83 $\frac{1}{4}$	261.5	5443.2	73.778	95 $\frac{3}{4}$	300.8	7200.5	84.856
83 $\frac{1}{2}$	262.3	5476.0	73.999	96	301.5	7238.2	85.077
83 $\frac{3}{4}$	263.1	5508.8	74.221	96 $\frac{1}{4}$	302.2	7275.9	85.299
84	263.8	5541.7	74.443	96 $\frac{1}{2}$	303.1	7313.8	85.520
84 $\frac{1}{4}$	264.6	5574.8	74.664	96 $\frac{3}{4}$	303.9	7351.7	85.742
84 $\frac{1}{2}$	265.4	5607.9	74.886	97	304.7	7389.8	85.964
84 $\frac{3}{4}$	266.2	5641.1	75.107	97 $\frac{1}{4}$	305.5	7427.9	86.185
85	267.	5674.5	75.329	97 $\frac{1}{2}$	306.3	7466.2	86.407
85 $\frac{1}{4}$	267.8	5707.9	75.550	97 $\frac{3}{4}$	307.	7504.5	86.628
85 $\frac{1}{2}$	268.6	5741.4	75.772	98	307.8	7542.9	86.850
85 $\frac{3}{4}$	269.3	5775.0	75.993	98 $\frac{1}{4}$	308.6	7581.5	87.071

Diam.	Circum.	Area.	Side of equal square.	Diam.	Circum.	Area.	Side of equal square.
98½	309.4	7620.1	87.293	99½	312.5	7775.6	88.179
98¾	310.2	7658.8	87.514	99¾	313.3	7814.7	88.401
99	311.	7697.7	87.736	100	314.1	7853.9	88.622
99½	311.8	7736.6	87.958				

To find, by means of the Table, the square or circle that will contain the area of a board or surface of given length and width.

RULE.—Find the area of the board or surface by multiplying its width by its length, and in the columns opposite the area thus found, headed respectively "*Diam.*," "*Circum.*," and "*Side of Equal Square*," will be found the dimensions of the circle and square that contains the area.

EXAMPLE.—What is the side of a square, and what the diameter and circumference of a circle, that will contain the same area as a board that is 22 inches wide by 12 feet long?

SOLUTION.—22 inches, width, \times 144 inches, length, = 3168 square inches, area of board: Then, in the table, opposite the area of 3166.9 (the nearest number to 3168) under the columns headed respectively "*Diam.*," "*Circum.*," and "*Side of Equal Square*," will be found 63½ inches, diameter, 109.4 inches, circumference, and 56.275 inches, side of square. *Ans.*

SQUARES, CUBES, AND ROOTS.

TABLE of Squares, Cubes, and Square and Cube Roots, of all numbers from 1 to 1000.

No.	Square.	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
1	1	1	1.	1.	54	2916	157464	7.348469	3.779763
2	4	8	1.414213	1.259921	55	3025	166375	7.416198	3.802953
3	9	27	1.732050	1.442250	56	3136	175616	7.483314	3.825862
4	16	64	2.	1.687401	57	3249	185193	7.549334	3.848501
5	25	125	2.236068	1.709975	58	3364	195112	7.615773	3.870877
6	36	216	2.449489	1.817121	59	3481	205379	7.681145	3.892996
7	49	343	2.645751	1.914933	60	3600	216000	7.745966	3.914567
8	64	512	2.828427	2.	61	3721	226951	7.810219	3.936497
9	81	729	3.	2.090084	62	3844	238228	7.874407	3.957592
10	100	1000	3.162277	2.154435	63	3969	250047	7.937253	3.97957
11	121	1331	3.316624	2.238900	64	4096	262448	8.	4.
12	144	1728	3.464101	2.269428	65	4225	274625	8.052257	4.020285
13	169	2197	3.605551	2.351335	66	4356	287496	8.121030	4.041240
14	196	2744	3.741657	2.440142	67	4489	300633	8.185352	4.061548
15	225	3375	3.872983	2.446212	68	4624	314000	8.216211	4.081656
16	256	4096	4.	2.519842	69	4761	328509	8.306823	4.101556
17	289	4913	4.123105	2.671222	70	4900	343000	8.366600	4.121285
18	324	5832	4.242640	2.620741	71	5041	357911	8.426149	4.140918
19	361	6859	4.358938	2.698402	72	5184	373248	8.485381	4.160168
20	400	8000	4.472135	2.711418	73	5329	389017	8.544003	4.179339
21	441	9261	4.682575	2.750023	74	5476	405224	8.602325	4.198336
22	484	10648	4.690415	2.802039	75	5625	421875	8.660254	4.217163
23	529	12167	4.795431	2.843067	76	5776	438976	8.717797	4.235824
24	576	13824	4.898979	2.884499	77	5929	456533	8.774964	4.254321
25	625	15625	5.	2.921018	78	6084	474552	8.831760	4.272659
26	676	17576	5.099019	2.962495	79	6241	493039	8.888194	4.290841
27	729	19638	5.196152	3.	80	6400	512000	8.944271	4.308870
28	784	21952	5.271502	3.036589	81	6561	531441	9.	4.326749
29	841	24399	5.385164	3.073317	82	6724	551368	9.055395	4.344441
30	900	27000	5.477226	3.107232	83	6889	571787	9.110433	4.362071
31	961	29791	5.567764	3.141381	84	7056	592704	9.165151	4.379519
32	1024	32768	5.656864	3.174802	85	7225	614125	9.219544	4.396890
33	1089	35937	5.744662	3.207534	86	7396	636056	9.273618	4.414005
34	1156	39304	5.830951	3.239612	87	7569	658503	9.327379	4.431047
35	1225	42875	5.916079	3.271006	88	7744	681472	9.380931	4.447960
36	1296	46656	6.	3.301927	89	7921	704969	9.433981	4.464745
37	1369	50653	6.092762	3.332222	90	8100	729000	9.486833	4.481401
38	1444	54872	6.164414	3.361975	91	8281	753571	9.539392	4.497942
39	1521	59319	6.241998	3.391211	92	8464	778698	9.591663	4.514357
40	1600	64000	6.324555	3.419952	93	8649	804357	9.643650	4.530655
41	1681	68921	6.403124	3.448217	94	8836	830584	9.695359	4.546836
42	1764	74088	6.480740	3.476127	95	9025	857375	9.746794	4.562903
43	1849	79407	6.557438	3.503398	96	9216	884736	9.797959	4.577857
44	1936	85144	6.633249	3.530348	97	9409	912673	9.848857	4.594701
45	2025	91125	6.708203	3.556843	98	9604	941192	9.899494	4.610438
46	2116	97336	6.782330	3.583048	99	9801	970299	9.949874	4.626055
47	2209	103824	6.855654	3.608824	100	10000	1000000	10.	4.641589
48	2304	110592	6.928208	3.634241	101	10201	1030300	10.049875	4.657010
49	2401	117649	7.	3.659309	102	10404	1061208	10.099504	4.672330
50	2500	125000	7.071067	3.684031	103	10609	1092727	10.148891	4.687548
51	2601	132651	7.141428	3.708430	104	10816	1124864	10.198039	4.702689
52	2704	140608	7.211162	3.732511	105	11025	1157625	10.246956	4.717694
53	2809	148877	7.280109	3.756266	106	11236	1191016	10.295630	4.732624

No.	Square.	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
107	11449	1225043	10.344080	4.747459	172	29584	6088448	13.114877	5.561198
108	11664	1259712	10.392304	4.762203	173	29929	5177777	13.152946	5.571034
109	11881	1295290	10.400600	3.778856	174	30276	6268024	13.189006	5.582770
110	12100	1333000	10.488080	4.791400	175	30625	5359375	13.228758	5.593445
111	12321	1367631	10.535653	4.805896	176	30976	5451716	13.264999	5.604079
112	12544	1404928	10.583050	4.820254	177	31329	5545233	13.301134	5.614673
113	12769	1444287	10.630145	4.834584	178	31684	5639751	13.337664	5.625226
114	12996	1485544	10.677077	4.848803	179	32041	5735330	13.374058	5.635741
115	13225	1529875	10.723805	4.862944	180	32400	5832000	13.410401	5.646216
116	13456	1567896	10.770329	4.876999	181	32761	5929741	13.445684	5.656652
117	13689	1601613	10.816638	4.890973	182	33124	6028568	13.480737	5.667051
118	13924	1643032	10.862780	4.904863	183	33489	6128487	13.515749	5.677411
119	14161	1685159	10.908712	4.918685	184	33856	6229504	13.550660	5.687734
120	14400	1728000	10.954451	4.932424	185	34225	6331625	13.601470	5.698019
121	14641	1771561	11.001361	4.946087	186	34596	6434856	13.633181	5.708267
122	14884	1815844	11.048536	4.959675	187	34969	6539203	13.674794	5.718479
123	15129	1860867	11.095936	4.973190	188	35344	6644667	13.711100	5.728654
124	15376	1906624	11.143528	4.986631	189	35721	6751269	13.747127	5.738791
125	15625	1953125	11.185339	5.000000	190	36100	6859000	13.781018	5.748897
126	15876	2000376	11.229427	5.013298	191	36481	6967871	13.815075	5.758965
127	16129	2048383	11.274927	5.026526	192	36864	7077888	13.850406	5.768998
128	16384	2097152	11.321708	5.039684	193	37249	7189057	13.882414	5.778996
129	16641	2146689	11.368716	5.052774	194	37636	7301384	13.923388	5.788960
130	16900	2197000	11.417552	5.065797	195	38025	7414875	13.964340	5.798894
131	17161	2248091	11.467222	5.078753	196	38416	7529536	14.005271	5.808798
132	17424	2299968	11.489125	5.091643	197	38809	7645373	14.046188	5.818668
133	17689	2352637	11.532562	5.104469	198	39204	7762392	14.071247	5.828516
134	17956	2406104	11.577836	5.117220	199	39601	7880599	14.100730	5.838342
135	18225	2460375	11.623950	5.129998	200	40000	8000000	14.124235	5.848053
136	18496	2515456	11.671903	5.142563	201	40401	8120601	14.174446	5.857765
137	18769	2571353	11.720699	5.155137	202	40804	8242404	14.212670	5.867440
138	19044	2628072	11.770344	5.167649	203	41209	8365427	14.249786	5.877130
139	19321	2685619	11.789820	5.180101	204	41616	8489664	14.282565	5.886855
140	19600	2744000	11.832159	5.192494	205	42025	8615125	14.317821	5.896568
141	19881	2803221	11.874342	5.204828	206	42436	8741816	14.352790	5.906941
142	20164	2863288	11.916350	5.217103	207	42849	8869743	14.387494	5.915481
143	20449	2924107	11.958220	5.229321	208	43264	8998911	14.422220	5.924493
144	20736	2985694	12.000000	5.241482	209	43681	9129329	14.456988	5.933471
145	21025	3048625	12.041594	5.253588	210	44100	9261000	14.491370	5.943911
146	21316	3112916	12.083040	5.265637	211	44521	9393931	14.525630	5.953841
147	21609	3178563	12.124356	5.277632	212	44944	9528124	14.560209	5.963731
148	21904	3244572	12.165525	5.289572	213	45369	9663597	14.594519	5.973591
149	22201	3307949	12.206555	5.301459	214	45796	9800344	14.628738	5.983426
150	22500	3375000	12.247445	5.313293	215	46225	9938375	14.662378	5.993227
151	22801	3442861	12.288205	5.325074	216	46656	10077696	14.696930	6.003004
152	23104	3511508	12.328828	5.336803	217	47089	10218313	14.730919	6.012744
153	23409	3581577	12.369316	5.348481	218	47524	10360232	14.764826	6.022450
154	23716	3652264	12.409673	5.360109	219	47961	10503459	14.798645	6.032125
155	24025	3723575	12.449899	5.371685	220	48400	10648000	14.832329	6.041761
156	24336	3795416	12.489996	5.383211	221	48841	10793861	14.865988	6.051364
157	24649	3867889	12.529964	5.394690	222	49284	10941044	14.899663	6.060940
158	24964	3940943	12.569800	5.406110	223	49729	11089567	14.933184	6.070486
159	25281	4013679	12.609520	5.417501	224	50176	11239424	14.966629	6.079977
160	25600	4086000	12.649100	5.428885	225	50625	11390625	15.000000	6.089401
161	25921	4158921	12.688577	5.440122	226	51076	11543176	15.033296	6.098769
162	26244	4232452	12.727922	5.451362	227	51529	11697089	15.066599	6.108071
163	26569	4306593	12.767145	5.462556	228	51984	11852352	15.099866	6.117305
164	26896	4401044	12.806348	5.473703	229	52441	12008989	15.133174	6.126469
165	27225	4495825	12.845532	5.484806	230	52900	12167000	15.166550	6.135572
166	27556	4590936	12.884698	5.495865	231	53361	12326331	15.199868	6.144634
167	27889	4686383	12.922848	5.506879	232	53824	12487168	15.233154	6.153664
168	28224	4782168	12.961488	5.517848	233	54289	12649337	15.266437	6.162639
169	28561	4878289	13.000000	5.528775	234	54756	12812904	15.299705	6.171605
170	28900	4974800	13.038404	5.539658	235	55225	12977875	15.332970	6.180544
171	29241	5000211	13.076696	5.550499	236	55696	13144256	15.366229	6.189477

No.	Square	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
237	56169	13312054	15.394804	6.188463	303	91204	27538208	17.378147	6.709172
238	56644	13481272	15.427248	6.197154	304	91904	27181217	17.406895	6.716549
239	57121	13651916	15.459824	6.205821	305	92416	26944464	17.435596	6.723950
240	57600	13824000	15.491933	6.214464	306	93025	26716245	17.464249	6.731316
241	58080	13997520	15.524174	6.223083	307	93636	26496616	17.492857	6.738665
242	58561	14172484	15.556349	6.231678	308	94249	26283444	17.521416	6.745997
243	59049	14349900	15.588457	6.240251	309	94864	26076816	17.549925	6.753313
244	59536	14528784	15.620499	6.248800	310	95481	25876826	17.578395	6.760614
245	60025	14709125	15.652476	6.257324	311	96100	25683480	17.606816	6.767899
246	60516	14890936	15.684387	6.265825	312	96721	25496791	17.635192	6.775168
247	61009	15074225	15.716233	6.274304	313	97344	25316760	17.663521	6.782422
248	61504	15259000	15.748015	6.282760	314	97969	25143399	17.691806	6.789661
249	62001	15445281	15.779738	6.291194	315	98596	24976714	17.720056	6.796884
250	62500	15633060	15.811388	6.299601	316	99225	24816711	17.748269	6.804091
251	63001	15822349	15.842976	6.307992	317	99856	24663384	17.776358	6.811284
252	63504	16013136	15.874507	6.316359	318	100489	24516729	17.804493	6.818461
253	64009	16205473	15.905973	6.324704	319	101124	24376743	17.832554	6.825624
254	64516	16399364	15.937377	6.333025	320	101761	24243424	17.860579	6.832771
255	65025	16594809	15.968719	6.341325	321	102400	24116780	17.888563	6.839903
256	65536	16791808	16.000000	6.349602	322	103041	24000000	17.916472	6.847021
257	66049	16990361	16.031219	6.357859	323	103684	23883384	17.944258	6.854124
258	66564	17190476	16.062378	6.366095	324	104329	23767899	17.972000	6.861211
259	67081	17392151	16.093476	6.374310	325	104976	23653552	18.000000	6.868284
260	67600	17595376	16.124515	6.382504	326	105625	23540341	18.027756	6.875343
261	68121	17799151	16.155544	6.390676	327	106276	23428264	18.055470	6.882388
262	68644	17994476	16.186464	6.398827	328	106929	23317321	18.083144	6.889419
263	69169	18191351	16.217274	6.406958	329	107584	23207512	18.110770	6.896435
264	69696	18389776	16.247976	6.415068	330	108241	23100000	18.138357	6.903436
265	70225	18589851	16.278572	6.423157	331	108900	22993681	18.165902	6.910423
266	70756	18791576	16.309050	6.431226	332	109561	22889552	18.193405	6.917396
267	71289	18994951	16.339413	6.439275	333	110224	22787611	18.220877	6.924345
268	71824	19200000	16.370000	6.447304	334	110889	22687856	18.248287	6.931300
269	72361	19406721	16.400429	6.455313	335	111556	22590299	18.275664	6.938236
270	72900	19615136	16.431676	6.463304	336	112225	22494944	18.303000	6.945149
271	73441	19825251	16.462707	6.471274	337	112896	22401781	18.330282	6.952033
272	73984	19937076	16.493624	6.479224	338	113569	22310800	18.357519	6.958893
273	74529	20050601	16.524429	6.487153	339	114244	22222001	18.384716	6.965729
274	75076	20165826	16.555216	6.495064	340	114921	22135384	18.411865	6.972542
275	75625	20282751	16.586000	6.502956	341	115600	22050945	18.438968	6.979332
276	76176	20401376	16.616784	6.510829	342	116281	21968681	18.466035	6.986109
277	76729	20521701	16.647567	6.518684	343	116964	21888592	18.493068	6.992874
278	77284	20643726	16.678333	6.526519	344	117649	21810681	18.520068	6.999629
279	77841	20767451	16.709093	6.534335	345	118336	21734944	18.547035	7.006376
280	78400	20892880	16.739840	6.542132	346	119025	21661381	18.573968	7.013105
281	78961	21019911	16.770576	6.549911	347	119716	21589992	18.600868	7.019819
282	79524	21148544	16.792555	6.557672	348	120409	21520781	18.627736	7.026516
283	80089	21278779	16.822203	6.565416	349	121104	21453744	18.654575	7.033196
284	80656	21410616	16.852229	6.573139	350	121801	21388881	18.681384	7.039859
285	81225	21544155	16.882193	6.580844	351	122500	21326100	18.708165	7.046508
286	81796	21679396	16.912134	6.588531	352	123201	21265501	18.734919	7.053143
287	82369	21816349	16.941074	6.596202	353	123904	21207084	18.761653	7.059766
288	82944	21955016	16.970012	6.603854	354	124609	21150849	18.788369	7.066369
289	83521	22095399	17.000000	6.611488	355	125316	21096784	18.815068	7.072959
290	84100	22237496	17.029886	6.619100	356	126025	21044881	18.841744	7.079526
291	84681	22381311	17.059872	6.626690	357	126736	21000000	18.868397	7.086079
292	85264	22526844	17.089857	6.634257	358	127449	20957241	18.895028	7.092619
293	85849	22674091	17.119842	6.641801	359	128164	20916604	18.921637	7.100158
294	86436	22823056	17.149828	6.649329	360	128881	20878081	18.948225	7.107693
295	87025	22973735	17.179815	6.656840	361	129600	20840680	18.974792	7.115216
296	87616	23126126	17.209800	6.664334	362	130321	20804391	19.001338	7.122727
297	88209	23280229	17.239785	6.671814	363	131044	20769212	19.027859	7.129226
298	88804	23436056	17.269870	6.679279	364	131769	20735251	19.054359	7.135713
299	89401	23593601	17.299955	6.686730	365	132496	20702504	19.080838	7.142188
300	90000	23752860	17.320000	6.694166	366	133225	20670881	19.107297	7.148641
301	90601	23913831	17.349361	6.701588	367	133956	20640484	19.133736	7.155081

No.	Square.	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
367	134689	4943086	19.157244	7.159599	432	186624	80621568	20.78469	7.559625
368	135424	4983698	19.183521	7.166095	433	187489	8112737	20.80865	7.56533
369	136161	5024309	19.20937	7.172480	434	188356	8174650	20.83366	7.571173
370	136900	5065300	19.235384	7.179064	435	189225	8236755	20.85863	7.576984
371	137641	5106481	19.261384	7.185516	436	190096	8298186	20.88361	7.582786
372	138384	5147884	19.287501	7.191964	437	190969	8359843	20.90861	7.588579
373	139129	5189511	19.313207	7.198405	438	191844	8421727	20.92444	7.594363
374	139876	52313524	19.339073	7.204832	439	192721	8483849	20.94031	7.600138
375	140625	52734375	19.364916	7.211247	440	193600	8546199	20.95622	7.605955
376	141376	53157376	19.390719	7.217652	441	194481	8608821	20.97217	7.611762
377	142129	53582633	19.416487	7.224045	442	195364	8671725	20.98816	7.617551
378	142884	54010152	19.44222	7.230427	443	196249	8734900	20.99419	7.623321
379	143641	54439939	19.46792	7.236797	444	197136	8798347	21.01026	7.629083
380	144400	54872000	19.493588	7.243156	445	198025	8862066	21.01631	7.634836
381	145161	5530631	19.519221	7.249504	446	198916	8926057	21.02239	7.640581
382	145924	55742968	19.544840	7.255841	447	199809	8990320	21.02850	7.646317
383	146689	56181887	19.570385	7.262167	448	200704	9054865	21.03464	7.652044
384	147456	56623104	19.595917	7.268482	449	201601	9119692	21.04080	7.657762
385	148225	57066625	19.62141	7.274786	450	202500	9184800	21.04697	7.663471
386	148996	57512456	19.646882	7.281079	451	203401	9250189	21.05316	7.669171
387	149769	57960603	19.67231	7.287362	452	204304	9315850	21.05937	7.674862
388	150544	58411072	19.697716	7.293633	453	205209	9381793	21.06560	7.680544
389	151321	5886386	19.723082	7.299893	454	206116	9448018	21.07184	7.686217
390	152100	5931900	19.748417	7.306143	455	207025	9514525	21.07810	7.691881
391	152881	59776471	19.773711	7.312383	456	207936	9581314	21.08437	7.697536
392	153664	60236224	19.798989	7.318611	457	208849	9648385	21.09066	7.703181
393	154449	60698457	19.824222	7.324839	458	209764	9715738	21.09697	7.708817
394	155236	61162984	19.849432	7.331067	459	210681	9783373	21.10330	7.714444
395	156025	61629913	19.874640	7.337284	460	211600	9851290	21.10964	7.720061
396	156816	62099340	19.899748	7.343490	461	212521	9919489	21.11599	7.725669
397	157609	62570773	19.924855	7.349696	462	213444	9987970	21.12236	7.731267
398	158404	63044192	19.949967	7.355892	463	214369	10056733	21.12874	7.736856
399	159201	63521199	19.974984	7.362087	464	215296	10125978	21.13513	7.742436
400	160000	64000000	20.000000	7.368281	465	216225	10195505	21.14154	7.748007
401	160801	64481201	20.024984	7.374496	466	217156	10265314	21.14797	7.753569
402	161604	64964804	20.049937	7.380722	467	218089	10335405	21.15442	7.759122
403	162409	65450827	20.074855	7.386947	468	219024	10405778	21.16089	7.764666
404	163216	65939264	20.099753	7.393162	469	219961	10476433	21.16737	7.770201
405	164025	66430115	20.12461	7.399366	470	220900	10547370	21.17387	7.775727
406	164836	66923416	20.149441	7.405560	471	221841	10618589	21.18038	7.781244
407	165649	67419141	20.174241	7.411754	472	222784	10689990	21.18690	7.786751
408	166464	6791731	20.199000	7.417947	473	223729	10761683	21.19343	7.792249
409	167281	68417929	20.223748	7.424140	474	224676	10833568	21.20000	7.797737
410	168100	68921000	20.248500	7.430332	475	225625	10905645	21.20658	7.803216
411	168921	69426531	20.273134	7.436523	476	226576	10977914	21.21318	7.808685
412	169744	69934525	20.297788	7.442713	477	227529	11050375	21.21980	7.814144
413	170569	70444987	20.322400	7.448903	478	228484	11123028	21.22643	7.819593
414	171396	70957944	20.346955	7.455093	479	229441	11195883	21.23308	7.825032
415	172225	71473375	20.371544	7.461283	480	230400	11268940	21.23974	7.830461
416	173056	71991296	20.396077	7.467472	481	231361	11342200	21.24641	7.835881
417	173889	72511711	20.420577	7.473661	482	232324	11415663	21.25309	7.841291
418	174724	73034624	20.445046	7.479850	483	233289	11489328	21.25978	7.846691
419	175561	73559056	20.469489	7.486039	484	234256	11563195	21.26648	7.852081
420	176400	74085000	20.493903	7.492228	485	235225	11637264	21.27319	7.857461
421	177241	74612461	20.518284	7.498417	486	236196	11711535	21.27991	7.862831
422	178084	75141444	20.542633	7.504606	487	237169	11785908	21.28664	7.868191
423	178929	75671967	20.566968	7.510795	488	238144	11860483	21.29338	7.873541
424	179776	76204024	20.591260	7.516984	489	239121	11935260	21.30013	7.878881
425	180625	76737625	20.615528	7.523173	490	240100	12010239	21.30689	7.884211
426	181476	77272776	20.639776	7.529362	491	241081	12085420	21.31366	7.889531
427	182329	77809481	20.663975	7.535551	492	242064	12160803	21.32044	7.894841
428	183184	78347744	20.688160	7.541740	493	243049	12236388	21.32723	7.900141
429	184041	78887569	20.712311	7.547929	494	244036	12312175	21.33403	7.905431
430	184900	79428900	20.736441	7.554118	495	245025	12388164	21.34084	7.910711
431	185761	80000000	20.760539	7.560307	496	246016	12464355	21.34766	7.915981

No.	Square.	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
497	247009	12763473	2.246346	7.921100	562	31684	177 0138	3.706539	8 232371
498	248004	12350592	22.315913	7.9264 8	563	316 84	178453547	23.72762	8 237 63
499	249011	124251499	22.38307	7.931710	564	318 96	179403144	23.74 684	8 262149
500	250020	125010000	22.36079	7.937005	565	319226	18036215	23.769128	8 271729
501	251031	125755501	2.383029	7.94 243	566	3203 8	181321496	23.79 74	8 277903
502	252044	12650308	2.40355	7.947573	567	321486	18228423	23.811761	8 276772
503	253059	127263527	2.42601	7.952847	568	322624	183250432	23.82750	8 281635
504	254076	128027044	22.44944	7.958114	569	323761	184220009	23.8532	8 286493
505	255095	128797625	22.47205	7.963374	570	324900	185193000	23.87462	8 291344
506	256116	129574216	22.49444	7.96867	571	32604	186169411	23.89506	8 296190
507	257139	130353339	22.516660	7.973873	572	327184	18714948	23.916521	8 301030
508	258164	131135512	22.53885	7.979112	573	328324	188132517	23.937418	8 30585
509	259191	131920729	22.561028	7.984344	574	329464	189119224	23.958 97	8 310694
510	260220	132709100	22.583179	7.989569	575	33062	190109375	23.979157	8 315517
511	261251	133500551	22.605309	7.994788	576	331774	191102576	24.	8 320335
512	262284	134297128	22.627417	8.	577	332924	192100033	24.020824	8 325147
513	263319	135095697	22.64953	8.005265	578	334084	193101630	24.041630	8 329954
514	264356	135906744	22.671588	8.010403	579	335241	194104539	24.062118	8 334755
515	265395	136720375	22.69 611	8.015595	580	336400	195112000	24.083189	8 339561
516	266436	137537596	22.715635	8.020779	581	337561	196122941	24.103941	8 344341
517	267479	138358413	22.737634	8.025957	582	338724	197137468	24.12466	8 349125
518	268524	139183832	22.759648	8.031129	583	339888	198156287	24.145 92	8 353904
519	269571	140013859	22.781571	8.036293	584	34104	199176704	24.166091	8 358678
520	270620	140848500	22.803 05	8.041451	585	34222	200120125	24.186773	8 363446
521	271671	141687741	22.824421	8.046633	586	34339	201120866	24.207436	8 368209
522	272724	142531584	22.847319	8.051748	587	34456	202122038	24.223082	8 372966
523	273779	143380039	22.86919	8.056886	588	34574	203124742	24.241 71	8 377713
524	274836	144233204	22.891046	8.062015	589	34692	204128649	24.269 22	8 382465
525	275895	145091125	22.912878	8.067143	590	34810	205133000	24.289195	8 387206
526	276956	145953576	22.934689	8.072262	591	34928	206137601	24.309191	8 391942
527	277919	146820583	22.95648	8.077374	592	35046	207142468	24.330100	8 396673
528	278984	147692152	22.978253	8.082481	593	35164	208147697	24.351591	8 401398
529	279951	148568389	22.99998	8.087579	594	35283	209153284	24.372115	8 406118
530	281020	149449200	23.021724	8.092672	595	35402	210159233	24.392621	8 410832
531	282091	150334681	23.043437	8.097759	596	35521	211165444	24.413111	8 415541
532	283164	151224832	23.065126	8.102839	597	35640	212171917	24.433583	8 420245
533	284239	152119649	23.086792	8.107912	598	35760	213178652	24.454038	8 424914
534	285316	153019136	23.10844	8.112990	599	35880	214185659	24.474476	8 429538
535	286395	153923305	23.130088	8.118011	600	36000	215192900	24.494897	8 434327
536	287476	154832156	23.151673	8.123093	601	36120	216199301	24.515301	8 439099
537	288559	155745689	23.173260	8.128144	602	36240	217205868	24.535688	8 443687
538	289644	156663904	23.194827	8.133186	603	36360	218212609	24.556008	8 448360
539	290731	157586839	23.216373	8.138223	604	36481	219219534	24.576411	8 453027
540	291820	158514400	23.237900	8.143255	605	36602	220226645	24.596747	8 457689
541	292911	159446601	23.25941	8.148277	606	36723	221233936	24.617067	8 462347
542	293904	160383456	23.280893	8.153293	607	36844	222241407	24.637370	8 466999
543	294999	161324969	23.302350	8.158300	608	36965	223249052	24.657655	8 471647
544	296096	162271144	23.323787	8.163300	609	37086	224256879	24.677925	8 476299
545	297195	163221985	23.34521	8.168303	610	37207	225264890	24.698178	8 480926
546	298296	164177496	23.36664	8.173302	611	37328	226273091	24.718414	8 485557
547	299399	165137683	23.388063	8.178299	612	37449	227281488	24.738633	8 490184
548	300504	166102552	23.40949	8.183293	613	37570	228290089	24.758836	8 494806
549	301611	167072109	23.430949	8.188284	614	37691	229298896	24.779023	8 499423
550	302720	168046360	23.452412	8.193272	615	37812	230307907	24.799193	8 504034
551	303831	169025311	23.473889	8.198257	616	37933	231317128	24.819347	8 508641
552	304944	170009968	23.495360	8.203231	617	38054	232326569	24.839484	8 513243
553	306059	171000237	23.516835	8.208202	618	38175	233336230	24.859603	8 517840
554	307176	172006204	23.538304	8.213170	619	38296	234346101	24.879710	8 522432
555	308295	173017875	23.559768	8.218135	620	38417	235356184	24.899799	8 527038
556	309416	174034248	23.581227	8.223098	621	38538	236366489	24.919871	8 531600
557	310539	175056339	23.602684	8.228059	622	38659	237376916	24.939927	8 536177
558	311664	176084152	23.624143	8.233018	623	38780	238387467	24.959967	8 540749
559	312791	177117709	23.645598	8.237971	624	38901	239398144	24.979992	8 545317
560	313920	178157000	23.667053	8.242920	625	39022	240409055	5.	8 549879
561	315051	179202031	23.688508	8.247874	626	39143	241420196	5.019992	8 554437

No.	Square.	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
627	393129	246491883	25.39968	8.58900	692	478864	33133888	26.20589	8.84055
628	394384	247673162	25.05998	8.56337	693	480224	332812557	25.321893	8.849344
629	395641	248861819	25.07987	8.56008	694	481604	334255854	25.343979	8.852598
630	396900	250170000	25.09980	8.572018	695	483000	335762800	25.367282	8.855849
631	398161	251509591	25.11971	8.577152	696	484404	337338400	25.391811	8.86005
632	399424	252880968	25.13961	8.581000	697	485804	338988700	25.417567	8.863337
633	400689	254284637	25.15949	8.586004	698	487204	340718800	25.444689	8.867075
634	401956	254840104	25.17937	8.590723	699	488600	342533000	25.473160	8.871409
635	403225	255447875	25.19920	8.595388	700	490000	344436000	25.503000	8.876010
636	404496	256109456	25.21904	8.599977	701	491404	346431000	25.534264	8.881266
637	405769	256825853	25.23885	8.604252	702	492804	348519000	25.566978	8.887468
638	407044	257609402	25.25866	8.608752	703	494200	349702000	25.601147	8.894706
639	408321	258461119	25.27844	8.613248	704	495600	350991000	25.636299	8.899520
640	409600	259281400	25.29822	8.617738	705	497000	352388000	25.672483	8.904010
641	410881	260179771	25.31797	8.622242	706	498404	353894000	25.709706	8.909336
642	412164	261106988	25.33771	8.626706	707	499804	355419000	25.747971	8.915106
643	413449	262064707	25.35744	8.631183	708	501200	356964000	25.787286	8.921236
644	414736	263053984	25.37715	8.635655	709	502600	358529000	25.827653	8.926931
645	416025	264075865	25.39685	8.640122	710	504000	359114000	25.869175	8.933121
646	417316	265130386	25.41653	8.644588	711	505521	359720000	25.911841	8.939507
647	418609	266218509	25.43619	8.649043	712	506944	360348000	25.955632	8.946190
648	419904	267341192	25.45584	8.653497	713	508369	361000000	25.999940	8.953268
649	421201	268499449	25.47547	8.657946	714	509800	361677000	26.045699	8.960813
650	422500	269693200	25.49500	8.662381	715	511225	362379000	26.092813	8.968783
651	423801	270923451	25.51447	8.666811	716	512656	363106000	26.141383	8.977100
652	425104	272190104	25.53400	8.671266	717	514084	363859000	26.191506	8.985868
653	426409	273493267	25.55364	8.675697	718	515521	364639000	26.243191	8.995106
654	427716	274832944	25.57328	8.680131	719	516960	365446000	26.296447	8.999868
655	429025	276210135	25.59292	8.684545	720	518400	366280000	26.351281	9.005100
656	430336	277624846	25.61249	8.688963	721	519841	367141000	26.407694	9.010813
657	431649	279077069	25.63198	8.693377	722	521284	368029000	26.465706	9.017000
658	432964	280567804	25.65140	8.697784	723	522729	368944000	26.525326	9.023644
659	434281	282107151	25.67095	8.702188	724	524176	369887000	26.586559	9.030740
660	435600	283695100	25.69046	8.706587	725	525625	370859000	26.649408	9.038286
661	436921	285332651	25.70992	8.710982	726	527076	371860000	26.713871	9.046283
662	438244	287019804	25.72936	8.715373	727	528529	372890000	26.780000	9.054732
663	439569	288756559	25.74878	8.719759	728	529984	373949000	26.847817	9.063637
664	440896	290542916	25.76819	8.724141	729	531441	375037000	26.917249	9.073000
665	442225	292379885	25.78759	8.728511	730	532900	376155000	27.015512	9.082813
666	443556	294267456	25.80697	8.732897	731	534361	377294000	27.103701	9.093121
667	444889	296205629	25.82634	8.737200	732	535824	378464000	27.192844	9.103929
668	446224	298194404	25.84569	8.741521	733	537289	379666000	27.283972	9.115240
669	447561	299243781	25.86503	8.745864	734	538756	380899000	27.377100	9.127000
670	448900	300353760	25.88435	8.750130	735	540225	382163000	27.472244	9.139200
671	450241	301524341	25.90367	8.754401	736	541696	383458000	27.569399	9.151913
672	451584	302755524	25.92296	8.758698	737	543169	384784000	27.668566	9.165100
673	452929	304037309	25.94223	8.763030	738	544644	386141000	27.769744	9.178786
674	454276	305369686	25.96140	8.767319	739	546121	387529000	27.872944	9.192900
675	455625	306752665	25.98057	8.771603	740	547600	388949000	27.978166	9.207500
676	456976	308186246	25.99972	8.775892	741	549081	390391000	28.085400	9.222600
677	458329	309670429	26.01923	8.780187	742	550564	391855000	28.194644	9.238100
678	459684	311205214	26.03843	8.785029	743	552049	393340000	28.305888	9.254000
679	461041	312790601	26.05762	8.789846	744	553536	394846000	28.419132	9.269900
680	462400	314427590	26.07680	8.794659	745	555025	396373000	28.534376	9.286200
681	463761	316116181	26.09596	8.799467	746	556516	397921000	28.651620	9.302900
682	465124	317856374	26.11529	8.804227	747	558009	399490000	28.770864	9.319900
683	466489	319649169	26.13468	8.808952	748	559504	401080000	28.892108	9.337200
684	467856	321494566	26.15393	8.813665	749	561001	402691000	29.015352	9.354700
685	469225	323392565	26.17304	8.818359	750	562500	404323000	29.140596	9.372400
686	470596	325343166	26.19161	8.823044	751	564001	405976000	29.267840	9.390300
687	471969	327347369	26.21018	8.827729	752	565504	407650000	29.397084	9.408400
688	473344	329405174	26.22875	8.832409	753	567009	409345000	29.528328	9.426700
689	474721	331517581	26.24730	8.837088	754	568516	411061000	29.661572	9.445200
690	476100	333684590	26.26585	8.841762	755	570025	412798000	29.796816	9.463900
691	477481	335906201	26.28438	8.846432	756	571536	414556000	29.934060	9.482700

No.	Square.	Cube.	Sq. Root.	Cu. Root.	No.	Square.	Cube.	Sq. Root.	Cu. Root.
757	573049	433795093	27.513632	9.113781	82.	675684	56541248	28.67054	9.367505
758	574764	435195112	27.531799	9.117793	82.	677329	567441767	28.68797	9.371302
759	576081	437215719	27.549954	9.121801	82.	678974	559176224	28.705400	9.375096
760	577600	438970000	27.568097	9.125805	85	680825	561515626	28.72281	9.378887
761	579121	440711051	27.586228	9.129808	86	682776	563555975	28.740215	9.372675
762	580644	442450728	27.604347	9.133803	87	684727	565609283	28.757500	9.384460
763	582169	444194917	27.622454	9.137797	88	686678	56766652	28.774882	9.390241
764	583696	44593744	27.640549	9.141788	89	688629	569727289	28.792360	9.396020
765	585225	447679125	27.658633	9.145774	90	689056	571790000	28.809700	9.397796
766	586756	449455096	27.676706	9.149757	91	690581	573856191	28.827170	9.401569
767	588289	451217683	27.694764	9.153737	92	692106	575930368	28.844410	9.405338
768	589824	452981852	27.712812	9.157713	93	693631	578003537	28.861730	9.409105
769	591361	454756660	27.730849	9.161686	94	695156	580093704	28.879056	9.412869
770	592900	456533060	27.748873	9.165656	95	696681	582183875	28.896366	9.416630
771	594441	458314111	27.766886	9.169622	96	698206	584277056	28.913664	9.420387
772	595984	460099648	27.784848	9.173588	97	699731	586373237	28.930961	9.424141
773	597529	461889717	27.802777	9.177544	98	701256	588480422	28.948229	9.427893
774	599076	463684824	27.820686	9.181500	99	702781	590597611	28.965499	9.431642
775	600625	465484975	27.838571	9.185452	100	704306	592720000	28.982753	9.435388
776	602176	467285176	27.856437	9.189401	81	705831	594842321	29.000000	9.439130
777	603729	469085433	27.874279	9.193347	82	707356	596964768	29.017266	9.442870
778	605284	470885732	27.892101	9.197289	83	708881	599087211	29.034561	9.446607
779	606841	472686083	27.910001	9.201228	84	710406	601211584	29.051868	9.450341
780	608400	474486480	27.927886	9.205164	85	711931	603335125	29.069181	9.454071
781	609961	476286931	27.945737	9.209096	86	713456	605459536	29.086499	9.457799
782	611524	478087432	27.963562	9.213025	87	714981	607583947	29.103861	9.461524
783	613089	480087937	27.981377	9.216950	88	716506	609708358	29.121199	9.465249
784	614656	481888448	28.000000	9.220872	89	718031	611832769	29.138544	9.468966
785	616225	483688959	28.017851	9.224791	90	719556	613957180	29.155899	9.472682
786	617796	485489470	28.035691	9.228706	91	721081	616081591	29.173200	9.476395
787	619369	487289981	28.053500	9.232618	92	722606	618206002	29.190500	9.480106
788	620944	489090492	28.071337	9.236527	93	724131	620330413	29.207801	9.483813
789	622521	490891003	28.089143	9.240435	94	725656	622454824	29.225100	9.487518
790	624100	492691514	28.106938	9.244338	95	727181	624579235	29.242401	9.491219
791	625681	494492025	28.124722	9.248241	96	728706	626703646	29.259700	9.494918
792	627264	496292536	28.142494	9.252130	97	730231	628828057	29.277000	9.498614
793	628849	498093047	28.160266	9.256022	98	731756	630952468	29.294300	9.502307
794	630436	500093558	28.178005	9.259911	99	733281	633076879	29.311600	9.505998
795	632025	502094069	28.195744	9.263797	100	734806	635201290	29.328900	9.509685
796	633616	504094580	28.213472	9.267679	81	736331	637325701	29.346200	9.513366
797	635209	506095091	28.231188	9.271559	82	737856	639450112	29.363500	9.517051
798	636804	508095602	28.248893	9.275435	83	739381	641574523	29.380800	9.520730
799	638401	510096113	28.266588	9.279308	84	740906	643698934	29.398100	9.524405
800	640000	512096624	28.284271	9.283177	85	742431	645823345	29.415400	9.528079
801	641601	514097135	28.301913	9.287044	86	743956	647947756	29.432700	9.531749
802	643204	516097646	28.319604	9.290907	87	745481	650072167	29.450000	9.535417
803	644809	518098157	28.337254	9.294767	88	747006	652196578	29.467300	9.539081
804	646416	519898668	28.354893	9.298623	89	748531	654320989	29.484600	9.542743
805	648025	521899179	28.372521	9.302477	90	750056	656445400	29.501900	9.546405
806	649636	523899690	28.390139	9.306327	91	751581	658569811	29.519200	9.550068
807	651249	525899701	28.407745	9.310176	92	753106	660694222	29.536500	9.553732
808	652864	527899712	28.425340	9.314019	93	754631	662818633	29.553800	9.557393
809	654481	529899723	28.442925	9.317859	94	756156	664943044	29.571100	9.561051
810	656100	531899734	28.460498	9.321697	95	757681	667067455	29.588400	9.564705
811	657721	533899745	28.478061	9.325532	96	759206	669191866	29.605700	9.568357
812	659344	535899756	28.495613	9.329363	97	760731	671316277	29.623000	9.571997
813	660969	537899767	28.513164	9.333191	98	762256	673440688	29.640300	9.575634
814	662596	539899778	28.530685	9.337016	99	763781	675565099	29.657600	9.579268
815	664225	541899789	28.548204	9.340833	100	765306	677689510	29.674900	9.582899
816	665856	543899800	28.565713	9.344647	81	766831	679813921	29.692200	9.586524
817	667489	545899811	28.583211	9.348473	82	768356	681938332	29.709500	9.590149
818	669124	547899822	28.600699	9.352295	83	769881	684062743	29.726800	9.593771
819	670761	549899833	28.618176	9.356095	84	771406	686187154	29.744100	9.597393
820	672400	551899844	28.635642	9.359901	85	772931	688311565	29.761400	9.600994
821	674041	553899855	28.653097	9.363704	86	774456	690435976	29.778700	9.604595

No.	Square	Cube.	Sq. Root.	Cu. Root.	No.	Square	Cube.	Sq. Root.	Cu. Root.
887	786766	697864103	9.72545	9.608181	914	891138	81113 3 4	20.724583	9.807736
888	78851	700227072	9.79932	9.611791	915	8930.6	84390.6 5	0.748 2	9.81318
889	7902.5	70259369	29.81610	9.6153.7	916	89491	84650.8 36	30.757113	9.81659
890	792100	7.4969001	29.8387	9.619001	917	89680	849278.1 3	0.71836	9.820117
891	79388	70347971	29.849623	9.622603	918	89870	85197182	0.7896	9.823572
892	79566	7 973228	23.866369	9.626201	919	900601	854670340	30.8684	9.827025
893	79744	712 21957	29.88310	9.629797	920	902500	85747100	30.87070	9.830475
894	79923	714516984	29.899832	9.63330.0	921	904401	860085351	0.83528	9.833923
895	80101	716717375	29.9165 0	9.636981	922	906300	862801409	30.84497	9.837369
896	80281	719321136	29.933259	9.640 69	923	908209	865521177	30.87069	9.840812
897	80460	721731273	29.949988	9.644154	924	910111	86820664	30.88789	9.844253
898	80640	724150792	9.96864	9.647736	925	91202	870933875	30.90 074	9.847692
899	80820	726572699	29.983328	9.651316	926	913936	873721816	30.919 49	9.851128
900	810000	7.90030 0	0.	9.654893	927	915849	8764 7491	30.93416	9.854561
901	81181	731432701	30.016662	9.65848	928	917768	879217912	30.95 57	9.857992
902	81364	733870808	30.033134	9.66 049	929	91968	8819740 0	30.96771	9.861421
903	81548	736314327	30.049955	9.665007	930	921600	8847310 0	30.983800	9.864848
904	81731	738763264	30.06659	9.669176	931	923524	887502681	31.	9.868 72
905	81925	741217765	30.083217	9.672740	932	925444	890273281	31.016124	9.871694
906	82083	743677416	30.099733	9.676301	933	927369	893058347	31.037241	9.875113
907	82264	746142613	30.116440	9.679800	934	929299	89584144	31.058389	9.878720
908	82446	748613312	0.133038	9.683416	935	931225	898632125	31.0 44.0	9.881915
909	82628	751089429	30.1 9 26	9.686970	936	933151	901423016	31.081640	9.885357
910	82810	753571000	30.16 200	9.690521	937	935081	904131063	31.096622	9.888767
911	82992	756058021	0.15270	9.694049	938	937002	907039332	1.11.69	9.892174
912	83174	758550508	30.1993 7	9.697615	939	938901	909955209	31.128704	9.895550
913	833 6	761048497	30.215889	9.701158	940	940800	912673 00	31.1482	9.898983
914	83 39	763551944	0.23242	9.704798	941	942724	915496611	31.169772	9.902383
915	83722	76 06075	30.248966	9.707280	942	944648	9183 048	31.189314	9.905781
916	83905	7685 5 96	30.265491	9.711777	943	946573	9211 7317	31.19947	9.909177
917	84 888	771095213	30.28207	9.716305	944	948506	9241 10424	31. 0877	9.91571
918	84272	773620634	0.298514	9.7 8835	945	950432	926859375	31.22 9 1	9.919662
919	84456	776151669	30.3101	9.72363	946	952357	929714783	1.240.96	9.919351
920	84640	778688001	30.31511	9.72788	947	954281	932574816	31.256999	9.922738
921	84824	781229961	30.34798	9.72941	948	956204	935441762	31.2799	9.9261 2
922	850 8	783777448	30.36412	9.7329 0	949	958128	938313739	31.28877	9.929504
923	8519.5	7863 0467	30.380915	9.736448	950	960040	941192040	31.30499	9.932883
924	8537 6	78888904	30.39738	9.739963	951	961962	944076141	31.3 0911	9.93 261
925	85562	79145312	0.41311	9.743475	952	963884	94696618	31.3 887	9.936636
926	85747	794022176	31.4 0245	9.746985	953	965806	94986 067	31.35250	9.94309
927	85932	796597983	0.44674	9.750493	954	967728	952763904	31.368774	9.947379
928	86118	799178752	30.47309	9.753998	955	969649	955671623	1.384708	9.949747
929	86304	801765789	30.47950	9.757500	956	971570	9585852 6	31.400639	9.9531 3
930	86490	804357000	0.49891	9.761010	957	973491	961404803	31.416 56	9.956477
931	86676	8 6954491	30.512 92	9.764497	958	975412	964420272	31.43247	9.959839
932	86862	809557568	30.528675	9.767992	959	977323	967361669	31.448350	9.9631 8
933	87048	812160237	30.54508	9.771484	960	979244	970299000	31.464265	9.966554
934	87235	814780504	30.56141	9.774974	961	981165	973242.71	31.480155	9.969909
935	87422	817400375	30.577769	9.778461	962	983086	976191488	31.496061	9.97262
936	87609	820025566	31.594 17	9.782946	963	985007	979146657	31.51 902	9.976612
937	87796	822656954	0.6104 5	9.786438	964	986928	982107784	31.532765	9.979595
938	87984	825297672	30.6 6785	9.789905	965	988849	985074875	1.5436 7	9.983104
939	88171	82793 019	0.643106	9.793386	966	990770	988047938	31.559467	9.986648
940	88360	830584000	0.659419	9.79581	967	992691	991026973	31.575 06	9.989570
941	88548	833217621	30.67572	9.799333	968	994612	994011992	31.591138	9.993328
942	88736	835826888	0.692018	9.802813	969	996533	997 02993	31.606961	9.996663
943	88924	838561807	30.70306	9.805271	970	10000000	1000000000	31.622776	10.

THE SOIL.

The soil is made up of decomposed rocks and decayed or decaying organic matter. The proportion of organic matter is small—not averaging in fertile soils more than five per cent. All of the rest of the soil is of a mineral origin, and has at some period formed a part of the rocky crust of the earth.

By the action of air, and heat, and frost, and the friction of running and falling water, and the movement of rocks and stones in moving water, these substances have been sufficiently pulverized to form the foundation material of our present soil.

During uncounted ages these processes have been going on, and they are still active; and, in addition to these, the chemical changes which result from the exposure of pulverized mineral matter to the action of air and moisture, and the successive growth and decay of plants, have operated, and are still operating, to ripen the soil to our uses.

In the early ages, when perhaps the composition of the atmosphere was different from what it is now (and when the soil was surely very different), only plants of a low order, such as are now extinct, could grow at all. These absorbed certain matters from the atmosphere, and, on their decay, gave them to the soil,—thus helping to fit it for the growth

of a higher order of plants, which were in time succeeded by others, and those by others, until, finally, the changes effected in the soil by the action of the chemical forces, and by the deposit of vegetable matter, have enabled it to produce the vegetation required for the uses of man.

Classification of soils.

Some soils were formed mainly of the rocks on which they now lie—as those of the granite region of New England—and these take their names from these rocks, as *granitic* soil, *limestone* soil, *sandstone* soil, &c.

Others have been formed by the deposit, by means of great floods, or the gradual silting of rivers. The latter of these (as the flat lands of the Mississippi Valley) are called *alluvial* soils; and the former (comprising those soils of varied composition in which occur clay, gravel, boulders, &c.) are called *diluvial* soils.

Another classification, which is much more definite, is the following :—

1. PURE CLAY consists of about 60 per cent. of silica and 40 per cent. of alumina and oxide of iron, usually chemically combined.
2. STRONGEST CLAY SOIL consists of pure clay, mixed with 5 to 15 per cent. of silicious sand.
3. CLAY LOAM consists of pure clay, mixed with 15 to 30 per cent. of fine sand.
4. LOAMY SOIL deposits from 30 to 60 per cent. of sand.

5. SANDY LOAM deposits from 60 to 90 per cent. of sand.
6. SANDY SOIL contains no more than 10 per cent. of pure clay.

To analyze the above soils with a view to classifying them.

RULE.—Weigh a portion of the soil and spread it thinly on writing paper, and dry it for an hour or two in an oven, the heat of which is not great enough to discolor the paper—the loss of weight is the quantity of *water* it contained.

Weigh and then boil another equal portion, and when thoroughly incorporated with the water, pour it into a vessel, and allow the sandy parts to deposit until the fine clay is also beginning to settle; then pour off the water, collect the sand, dry as before, and again weigh, which will give the per cent. of *sand* it contained.

The above classification and analysis of soils have reference only to the water, clay, and sand which they contain, while lime is also an important constituent, of which they are rarely entirely destitute. This gives rise to a further classification.

7. MARLY SOIL is one in which the proportion of lime is more than 5, and not over 20 per cent. of the whole weight.

8. CALCAREOUS SOIL, in which the lime exceeds 20 per cent.

To analyze marly and calcareous soils, with a view to their classification as above.

RULE.—Mix 100 grains of the dry soil with half a pint

of water, and add half a wine-glassful of muriatic acid; stir it thoroughly during the day, and let it stand and settle over night. Pour off the clear liquid in the morning, and again fill the vessel with water and stir thoroughly, and when clear again pour it off; dry the soil and weigh it. The loss is the quantity of lime the soil contained. If it exceeds 5 grs., class as a *marly soil*; if more than 20 grs., class as a *calcareous soil*.

9. VEGETABLE MOULDS, which are of various kinds, containing from 15 to 60 or 70 per cent. of organic matter.

To analyze vegetable moulds, with a view to their classification as above.

RULE.—Dry the soil well in an oven, and weigh it; then heat it to a dull redness, over a lamp or bright fire, until the combustible matter is burned away and evaporated. Again weigh it, and the loss is the quantity of organic matter it contained.

Besides the foregoing ingredients, every soil must contain more or less of all the elements which enter into the composition of vegetation. They must hold, in a form adapted to its growth and support, *silex, alumina, carbonate of lime, sulphate of lime, potash, soda, magnesia, sulphur, phosphorus, oxide of iron, manganese, chlorine*, and, probably, *iodine*. They are called the “inorganic or earthy parts of soil,” and constitute from one-half of one per cent. to over ten per cent. of all vegetables. Their analysis is too diffi-

cult and complicated to be attempted by any but a practical agricultural chemist.

The value of soil analysis, even when made by the most careful and skilful chemists, is practically very little. The quantity of matter which is capable of affording food to plants is so very small, in proportion to the whole bulk of the soil, even in those of the most fertile character, that it is questionable whether a sample to be analyzed could be so carefully prepared as to represent the average character of the whole field. Then, again, if we were to procure a correct analysis of a very fertile soil, and then were to crop it for a series of years without manure until it refused to produce paying crops, and were to have it analyzed again, it is not likely that the chemist would detect any change in its composition. In like manner, if we were to add to it 500 lbs. to the acre of bone dust,—enough to make it produce abundantly,—analysis would fail to detect the small quantity of phosphate of lime that we had added in the bones.

Another argument against the value of the analysis of the soil, and a very strong one, is found in the fact that the fertility of the soil depends less on the *quantity* of plant food that it contains than on its *condition*. The roots of plants cannot feed on the *inside* of a pebble; they can only apply their pumps to its surface and take in so much of what is there exposed as can be dissolved in the moisture which goes to form their sap. Neither can roots travel about in the soil; they grow into certain places, and there they remain.

If an inch away from them there is a mass of rich food, they cannot make use of it—save by sending out new shoots to embrace it—but must remain content with the poorer tract in which they lie. Consequently, the uniform *distribution* of the plant food, its *solubility*, and its *exposure* on the surfaces of the particles of the soil are quite as important as its quantity.

Chemical analysis teaches us none of those things—at least it does not teach them so definitely as we would need to know them to be able to make any practical use of its assistance.

In addition to these, fertile soils must also contain *carbon*, *oxygen*, *nitrogen*, and *hydrogen*, which are called the organic parts of soils, from their great preponderance in vegetables and animals. of which they constitute from 90 to over 99 per cent.

General results of analytical examinations of soils.

1. A due admixture of organic matter is favorable to the fertility of a soil.
2. This organic matter is the more valuable in proportion to the quantity of nitrogen it holds in combination.
3. The mineral part of the soil must contain all those substances which are met with in the ash of the plant, and in such a state of chemical combination that the roots of plants can readily take them up in the requisite proportions.

TABLE, showing the composition, in 1000 parts, of different kinds of soil.

CONSTITUENTS.	Fertile without manure.	Fertile with manure.	Very Barren.
Organic matter,	97.	50.	40.
Silica,	648.	833.	758.
Alumina,	57.	51.	101.
Lime,	59.	18.	4.
Magnesia,	8.	8.	1.
Oxide of Iron,	61.	30.	91.
“ of Manganese,	1.	3.	trace
Potash,	2.	trace
Soda,	4.
Chlorine,	2.
Sulphuric Acid,	2.	1.
Phosphoric “	4.	2.
Carbonic “	40.	4.
Loss,	15.	5.

NOTE.—The soil designated “fertile without manure” has been cultivated sixty years without manuring, yielding abundant crops. The soil designated “fertile with manure” has been cultivated over forty years, yielding good crops with ordinary manuring; while that designated “very barren” could scarcely be made to yield anything by the greatest manuring and most careful cultivation.

The following is an analysis of three specimens of very fertile soils, made by Sprengel:—

	Soil near Osterbruch.	From the banks of the Weser, near Hoya. near Weserbe.	
Silica, Quartz, Sand, and Silicates.....	84.510	71.849	83.318
Alumina	6.435	9.350	3.085
Oxides of Iron.....	2.395	5.410	5.840
Oxides of Manganese.....	0.450	0.925	0.620
Lime.....	0.740	0.987	0.720
Magnesia	0.525	0.245	0.120
Potash and Soda extracted by water.....	0.009	0.007	0.005
Phosphoric Acid.....	0.120	0.131	0.065
Sulphuric Acid.....	9.046	0.174	0.025
Chlorine in common Salt.....	0.006	0.002	0.006
Humic Acid	0.780	1.270	0.800
Insoluble Humus.....	2.995	.550	4.126
Organic matters containing Nitrogen.....	0.960	2.000	1.220
Water.....	0.029	0.100	0.150

The above had remained a long time in pasture, and the second was remarkable for the fattening qualities of its grass when fed to cattle.

The following are arable lands of great fertility:—

	Soil from Moravia.	From Ohio.		Soil from Belgium.
		Soil.	Subsoil.	
Silica and fine Sand.....	77.209	87.143	94.261	64.517
Alumina.....	8.514	5.666	1.876	4.810
Oxides of Iron.....	6.592	2.220	2.336	8.316
Oxide of Manganese.....	1.520	0.360	1.200	0.800
Lime.....	0.927	0.564	0.243	Carb. of Lime. 9.403
Magnesia.....	1.160	0.312	0.310	Carb. of Mag. 10.361
Potash, chiefly combined with Silica.....	0.140	0.120	0.240	{ 0.100 0.013
Soda, ditto.....	0.640	0.025		
Phosphoric Acid, combined with Lime and Ox. of Iron..	0.651	0.060	trace	1.221
Sulphuric Acid and Gypsum..	0.011	0.027	0.034	0.009
Chlorine in common Salt.....	0.010	0.036	trace	0.003
Carbonic Acid united to the Lime.....	0.080
Humic Acid.....	0.978	1.304	0.447
Insoluble Humus.....	0.540	1.072
Organic Substances containing Nitrogen.....	1.108	1.011

“Of these soils, the first had been cropped for 160 years successively, without either manure or naked fallow. The second was a virgin soil, and celebrated for its fertility. The third had been unmanured for twelve years, during the last nine of which it had been cropped with beans, barley, potatoes, winter barley and red clover, clover, winter barley, wheat, oats, naked fallow.”—*Johnston*.

Depth of soil—its importance.

If 50 be assumed as the value of a given soil when it is six inches deep, its value when of different depths will be as follows:—

If 3 inches deep, it is worth	38	If 8 inches deep, it is worth	58
4 " " " "	42	9 " " " "	62
5 " " " "	46	10 " " " "	66
6 " " " "	50	11 " " " "	70
7 " " " "	54	12 " " " "	74

Hence each farmer may make an estimate for himself, with regard to every variety of his soil, whether the cost of increasing its depth will equal or exceed its value after the task is performed.

This, of course, supposes that the soil is of the same quality throughout its whole depth, and it refers only to its chemical composition. There are other considerations which make the depth of the soil more important even than the above table will indicate. If a soil is equally rich throughout its whole depth, it would be of more than double value if of double depth; for its ability to withstand drought, and its great capacity to absorb the water of heavy rains (without being made too wet) would make it better, irrespective of the elements of fertility that it might contain. Then again, some soils which are of apparently no value may be made quite fertile by being ploughed a little deeper than has been done.

TABLE, *showing the weight per cubic foot of the different kinds of earth.*

Loose earth or sand.....	95 lbs.	Clay.....	135 lbs.
Common soil.....	124 "	Clay and stones.....	160 "
Strong soil.....	127 "	Brick.....	125 "
Chalk.....	174 "		

NOTE.—23 cubic feet of sand, 18 cubic feet of earth, or 17 cubic feet of clay, make a ton. Eighteen cubic feet of

gravel or earth, before digging, make 27 cubic feet when dug.

As a rough estimate, it may be stated that an acre of ordinary soil weighs 100 tons for every inch of its depth.

EXHAUSTION OF SOILS.

Each crop taken from a field exhausts the soil to the extent of the inorganic or earthy substances that are found in the totality of the crop removed. Unless, therefore, these elements are returned to the soil in some shape it gradually loses its fertility, and finally refuses to produce altogether. Hence the necessity for manuring, irrigating, or *resting* the soil, that it may again, by accumulating these elements, recover its fertility. By returning a crop *in toto* to the soil, by ploughing it in or leaving it to decay and mingle again with it, it accumulates in mass and grows in fertility, not by the substances thus returned to it, but by fertilizing elements gathered in or combined from the atmosphere, by rains and dews descending on it, and by capillary attraction from beneath.

By knowing the composition of the subtracted crops and the added manures, the farmer can keep a debit and credit account with his fields, which will be sufficiently accurate to enable him always to keep his land improving. To enable him to ascertain approximately what his various crops remove from the soil, we introduce the following tables, &c. To

ascertain what will replace this subtraction, let him consult the section on manures.

TABLE, showing the organic substances removed from the soil in 1000 lbs. each of the following crops when perfectly dry.

	Carbon. lbs.	Hydrogen. lbs.	Oxygen. lbs.	Nitrogen. lbs.	Ash. lbs.
Hay.....	458	50	387	15	90
Red Clover Hay.....	474	50	378	21	77
Potatoes.....	440	58	447	15	40
Wheat.....	461	58	434	23	23
Wheat straw.....	484	53	389½	3½	70
Oats.....	507	64	367	22	40
Oat-straw.....	501	54	390	4	51

Johnston.

NOTE.—Of all the vegetable productions which are gathered as food for man or beast in their *dry state*—

Carbon forms nearly *one-half* by weight.

Oxygen rather more than *one-third*.

Hydrogen little more than *five per cent*.

Nitrogen from 1½ to 4 *per cent*.

Earthy matter from 1 to 20 *per cent*.

TABLE, showing the quantity of inorganic matter removed from the soil in 1000 lbs. each of the following crops in their ordinary state of dryness.

	lbs.		lbs.
Wheat.....	about 20	Beans.....	about 30
Wheat-straw.....	" 50	Peas.....	" 30
Barley.....	" 30	Pea-straw.....	" 50
Barley-straw.....	" 50	Meadow Hay.....	" 50 to 100
Oats.....	" 40	Clover Hay.....	" 90
Oat-straw.....	" 60	Rye-grass Hay.....	" 95
Rye.....	" 20	Potatoes.....	" 8 to 15
Rye-straw.....	" 40	Turnips.....	" 5 to 8
Indian Corn.....	" 15	Carrots.....	" 15 to 20
Indian Corn stalk, &c.....	" 50		

Johnston.

TABLE, showing the quantity and kinds of inorganic matter removed from the soil in 1000 lbs. each of the following crops.

	Potash.	Soda.	Lime.	Magnesia.	Alumina.	Silica.	Sulphuric Acid	Phosphoric Acid.	Chlorine.	Oxide of Iron.	Oxide of Manganese.	Total in every 1000 lbs.
Wheat—Grain.....	2.25	2.40	0.96	0.90	0.28	4.00	0.50	0.40	0.10	traces		11.77
" Straw.....	0.20	0.29	2.40	0.32	0.90	28.70	0.37	1.70	0.30			35.18
Barley—Grain.....	2.78	2.90	1.06	1.80	0.25	11.82	.59	2.10	0.19	traces		23.49
" Straw.....	1.80	0.48	5.54	0.76	1.46	33.56	1.18	1.60	0.70	0.14	0.20	52.42
Oats—Grain.....	1.50	1.32	0.86	0.67	0.14	19.78	0.35	0.70	0.10	0.40		25.80
" Straw.....	8.70	0.02	1.52	0.22	0.04	45.88	0.79	0.12	0.05	0.02	0.02	57.40
Rye—Grain.....	5.32	*	1.22	0.44	0.24	1.64	0.23	0.46	0.09	0.41	0.34	10.40
" Straw.....	0.52	0.11	1.78	0.12	3.25	22.97	1.70	0.51	0.17			27.93
Field } Bean.....	4.15	8.10	1.65	1.58	0.34	1.26	0.89	2.92	0.41			21.36
Bean } Straw.....	16.56	0.50	6.24	2.09	0.10	2.20	0.34	2.26	0.80	0.07	0.05	31.21
Field } Pea.....	8.10	7.39	0.58	1.36	0.20	4.10	0.53	1.80	0.38	0.10		24.64
Pea } Straw.....	2.35		27.30	3.42	0.60	9.96	3.37	2.40	0.04	0.27	0.07	49.71
Potatoes } Roots.....	4.028	2.334	.331	.324	.056	.034	.540	.401	.160	.032		8.284
" Tops.....	8.19	.09	12.97	1.70	.04	4.94	.42	1.97	.50	.02		30.84
" Roots.....	2.386	1.048	.752	.254	.036	.388	.801	.367	.239	.032		6.303
Turnips, } Leaves.....	3.23	2.22	6.20	.59	.03	1.28	2.52	.98	.87	.17		18.09
Carrots.....	3.533	.922	.657	.394	.039	.137	.270	.514	.070	.033	.060	6.619
Parsnips.....	2.079	.702	.468	.270	.024	.162	.192	.100	.178	.005	?	4.180
Rye Grass.....	8.81	3.94	7.34	0.90	0.81	27.72	3.53	0.25	0.06			52.86
Red Clover.....	19.95	5.29	27.80	3.33	0.14	3.61	4.47	6.57	3.62			74.78
White Clover.....	31.05	5.79	33.48	3.05	1.90	14.73	3.53	5.05	2.11	0.63		91.32
Lucern.....	13.40	6.15	49.3	3.48	0.30	3.30	4.04	13.07	3.15	0.50		95.52
Sainfoin.....	20.57	4.37	21.95	2.88	0.66	5.00	3.41	9.16	1.57			69.57

Sprengel.

NOTE.—In the foregoing table, the grain, beans, peas, straw, and hay, are estimated after they have been dried in the air; the roots as they have been taken from the field. The potato loses in drying 69 per cent. of water; the turnip 91; the carrot 87; the turnip-leaf 86; the carrot-leaf, parsnip, and parsnip-leaf, each 81; and the cabbage 93.

Besides the *organic elements* present in each of the above crops, and which form about 97 per cent. of the entire dried weight of each, it is not only necessary that all the above

* Included in potash.

inorganic substances should exist in the soil, but that they be also found in a form adapted to the wants of the growing crop.



ANALYSIS OF THE ASH OF THE HOP, *showing the elements it removes from the soil.*

In 100 parts there are of

	Vine & Blossom.	Blossom.		Vine & Blossom.	Blossom.
Silica.....	13.24	21.05	Magnesia.....	4.09	5.77
Chloride of Sodium...	7.73		Sulphuric Acid.....	4.63	5.41
" Potassium.	3.77		Phosphoric "	6.34	9.08
Soda.....	0.13		Phosphate of Iron	3.79	7.45
Potash.....	21.49	25.18	Chloride of Potassium.		1.67
Lime.....	34.79	15.98	Alumina.....		a trace

The following tables, extracted from Waring's Elements of Agriculture, will be found convenient for ordinary computations :—

Amount of Inorganic Matter removed from the soil by ten bushels of grain, &c., and by the straw, &c., required in their production—estimated in pounds :

	Wheat.	1200 lbs. Wheat Straw.	Rye.	1620 lbs. Rye Straw.
Potash.....	2.86	8.97	2.51	11.84
Soda.....	1.04	.12	1.33	.20
Lime.....	.34	4.84	.56	5.91
Magnesia.....	1.46	2.76	1.18	1.58
Oxide of Iron.....	.08	.94	.15	.88
Sulphuric Acid.....	.03	4.20	.11	.05
Phosphoric Acid.....	6.01	2.22	5.64	2.49
Chlorine.....		.79		.80
Silica.....	.14	47.16	.05	42.25
Pounds carried off.....	12	72	11½	66

	Corn.	1620 lbs. Corn Stalks.	Oats.	700 lbs. Oat Straw.
Potash.....	2.78	6.84	1.69	12.08
Soda.....		19.83		
Lime.....	.12	6.02	.39	3.39
Magnesia.....	1.52	4.74	.64	1.59
Oxide of Iron.....		.57	.02	.78
Sulphuric Acid.....		.36	.66	1.41
Phosphoric Acid.....	4.52	12.15	2.80	1.07
Chlorine.....		1.33	.02	1.36
Silica.....	.06	19.16	.18	20.32
Pounds carried off.....	9	71	6½	42


	Buck Wheat.	Barley.	660 bbls. Barley Straw.	2000 lbs. Flax.
Potash.....	1.01	1.90	2 57	11.78
Soda.....	2.13	1.18	.23	11.82
Lime.....	.78	.96	3.88	11.85
Magnesia.....	1.20	1.00	1.31	9.38
Oxide of Iron.....	.14	.20	.90	7.32
Sulphuric Acid.....	.25	.01	.66	3.19
Phosphoric Acid.....	5.40	5 35	1.25	13.05
Chlorine.....		.01	.40	2.90
Silica.....	.09	3.90	28.80	25.71
Pounds carried off.....	11	14	40	100

	Beans.	1120 lbs. Bean Straw.	Field Peas.	1966 lbs. Pea Straw.
Potash.....	5.54	36.28	5.90	3.78
Soda.....	1.83	1.09	1.40	
Lime.....	98.98	13.60	.81	43.93
Magnesium.....	.28	4.55	1.30	5.50
Oxide of Iron.....	.10	.20	.15	1.40
Sulphuric Acid.....	.16	.64	.64	5.43
Phosphoric Acid.....	7.80	5.00	5.50	3.86
Chlorine.....	.13	1.74	.23	.08
Silica.....	.18	4.90	.7	16.02
Pounds carried off.....	17	68	16	80

	1 Ton Turnips.	635 lbs. Turnip Tops.	1 Ton Potatoes.	2000 lbs. Red Clover.
Potash.....	7.14	4.34	27.82	31.41
Soda.....	.86	.84	.93	8.34
Lime.....	2.31	3.61	1.03	43.77
Magnesia.....	.91	.48	2.63	5.25
Oxide of Iron.....	.23	.13	.26	.23
Sulphuric Acid.....	2.30	1.81	6.81	7.05
Phosphoric Acid.....	1.29	1.31	6.25	10.28
Chlorine.....	.61	2.35	2.13	5.86
Silica.....	1.36	.13	2.14	5.81
Pounds carried off.....	17	15	50	118

	2000 lbs. Meadow Hay.	2000 lbs. Cabbage. Water 9-10
Potash.....	18.11	5.25
Soda	1.35	9.20
Lime.....	22.95	9.45
Magnesia.....	6.75	2.70
Oxide of Iron.....	1.69	.25
Sulphuric Acid.....	2.70	9.60
Phosphoric Acid.....	5.97	5.60
Chlorine.....	2.59	2.60
Silica.....	37.89	.35
Pounds carried off.....	100	45

MANURES.



In order to restore to the soil the matters which have been taken from it by the removal of its produce, as well as to add to its power to produce—to make it richer, or to keep it from growing poorer—we make use of what are known as *manures*.

This term is a very comprehensive one, and is taken to mean all substances—whatever their character or origin—which will have the effect of causing a larger growth of vegetation.

Manures may be either *mechanical* or *chemical* in their

mode of action, or they may partake of both of these characters. For instance, barn-yard manure is both mechanical and chemical in its effect.

By reason of its bulk and its coarseness it loosens the soil and makes it more porous when mixed with it; when it is used as a top-dressing it shades the ground, and protects it in a measure against the effect of frost and of too great heat; being a very active absorbent of moisture, it modifies the effect of drought; its decomposition produces heat, and raises the temperature of the soil.

All of these are *mechanical* effects.

On the other hand, it affords to the roots of plants substances which enter directly into their structures, as chemical constituents; it also yields various acids, alkalies, and salts which enter into combination with the constituent parts of the soil, and—in one way or another—make them more available as plant food.

These are *chemical* effects.

The use of Manures.

In the use of manures the farmer should be guided not only by the effect that will be produced on the immediate crop—although this is, of course, the first consideration—but quite as much by the condition in which the soil will be left for the production of future crops. Unless he does this he may find that, while he has reaped a temporary benefit, he has inflicted a lasting injury on his fields.

It will be remembered that in our account of the soil it

was shown that the amount of mineral plant food that is actually present in the soil in an available form is extremely limited. In a state of nature, our fields would produce only such crops as could be fed by the small amount of this plant food which is rendered available from year to year, and there would be no diminution of production. On the contrary, the decay of the crop of one year would probably add to the supply available for the next year. *The removal of the crop by man, not the production of a crop which on decay returns its elements to the soil*, is what impoverishes—is what makes the use of manure vitally necessary on all but virgin lands.

The larger the crop—provided it *decays on the land*—the more the fertility of the soil is increased.

The larger the crop—provided it is *removed from the land*—the more the fertility of the soil is diminished.

If the crop is made larger by the use of manure, and is removed from the land, the manure has caused a larger amount of mineral plant food to be taken away. But if the manure itself contains the full equivalent of what enters into the crop, and so makes up for its drain upon the soil, there will be no impoverishment. If, on the other hand, the manure does not contain the equivalent of the ash-constituents of the crop, but has only stimulated it to take an extra supply from the soil, the injury is obvious.

In some cases, a soil that will produce 10 bushels of wheat without manure will produce 25 bushels if dressed with 100

lbs. of sulphate of ammonia. The extra 15 bushels contain about 18 lbs. of mineral matter more, which was supplied by the manure, and this is equal to one and a half year's supply for the natural crop of the land. The effect of this sort of farming is that the soil is made to produce more than it can afford to in one year, and has its supply of mineral plant food exhausted to the detriment of its future productiveness.

Twenty years ago, the wheat lands of Delaware, which had been producing very small crops, were made, by the use of very small doses of Peruvian guano, to double, triple, even quadruple their yield. The farmers were immensely elated. They had found a sort of philosopher's stone, and a few years would make their fortune. Alas for their hopes—a very few years demonstrated the fact that the guano had been a curse rather than a blessing. Their lands were poorer than ever, and even largely increased doses of the specific were powerless to bring them up even to their old standard.

Had the wheat and straw been consumed on the farm, and all of their mineral constituents returned to the soil, the guano would have been a means of great permanent improvement.

Or, had the same increase of production been effected by the use of a manure containing the full equivalent of what the crop was to take from the soil, the impoverishment of the land would have been prevented.

The foregoing is intended to convey the fundamental ideas which we should bear in mind in deciding what manures we are to use, and in what quantity. It is quite impossible to establish any set of rules which shall be an exact guide for all cases, but the following are always a *safe* guide : —

1. *Apply in the manure the full quantity of the different ash ingredients of the crops that will be produced before manure will be applied again.*

2. *Procure from abroad manure containing the full quantity of the different ash ingredients of all produce sold from the farm, and allow none to be wasted at home.*

A close adherence to these two rules, accompanied by good cultivation, and the draining of such land as needs draining, will make any farmer rich who exercises ordinary judgment and prudence in the management of his affairs.

To speak with scientific accuracy, it is not necessary to return quite *all* that the crops take away.

The processes by which soils were originally formed being still in operation, there is a constant fresh development of plant-food in the ground, and this will, in greater or less degree, compensate for the loss by the removal of crops.

Practically, however, it is best to place this development of fresh matter to the account of improvement, and, by making up the full amount of all removals, to make sure that the land is constantly growing better instead of worse.

As want of space forbids a more full discussion of the

established theories concerning the use of manure, the attention of the reader is called to the following:—

Classification and description of manures.

Manures naturally divide themselves into such as are of mineral, of vegetable, and of animal origin.

Mineral manures are such as originate from various mineral substances, such as lime, which is the product of limestone, marble, chalk, or marl, after the carbonic acid has been expelled by an intense heat; marls, which are composed of carbonate of lime, mixed with clay, sand, or loam; shell sand, calcareous sand, green sand marl, gypsum, phosphate of lime, salt, and salts of various kinds, &c.

Vegetable manures are such as are produced from decomposed vegetable matters, which also contain some of the inorganic or mineral substances.

Animal manures consist chiefly of the flesh, blood, bones, horns, and hair of sea and land animals, and of the solid and liquid excrements of land animals and birds, and also contain some of the inorganic or mineral matters.

Analysis of Fish Guano.

Water expelled by 212° heat..	8.06	Sulphate of Magnesia.....	0.71
Sand.....	0.33	" Potash.....	2.05
Oil.....	2.40	" Soda.....	2.42
Organic Matter	50.72	Chloride of Sodium.....	1.12
Super-Phosphate of Lime.....	9.85	Sulphate of Ammonia.....	2.72
Sulphate of Lime, Hydrated...	19.62	<i>Dr. Apjohn.</i>	

Analysis of Peruvian Guano.

In every 100 parts there are of

Organic Matter, containing Nitrogen, including Urates of Ammonia, and capable of affording from 8 to 17 per cent. of Ammonia, by slow change in the soil.....	50.
Water.....	11.
Phosphate of Lime.....	25.
Ammonia, Phosphate of Magnesia, Phosphate of Ammonia and Oxalate of Ammonia, containing from 4 to 9 per cent. of Ammonia....	13.
Silicious matter from the crops of birds.....	1.

Dr. Ure.

Another analysis.

Water.....	13.09
Organic Matter, containing Ammonia.....	53.17
Common Salt and Sulphate of Soda.....	4.63
Carbonate of Lime.....	4.18
Phosphate of Lime and Magnesia.....	23.54
Silicious Matter or Sand.....	1.39

Johnston.

Professor S. W. Johnson publishes the following table:—

Analysis of Peruvian Guano.

	I.		II.		III.		IV.
Water.....	66.32	65.18	12.63	12.70	68.00	68.70	59.46
Organic Matter.....			52.27	51.46			
Ammonia, potential.....	5.82	5.95	16.03	15.98	17.86	18.85	16.32
“ actual.....	8.93	9.08					
Phosphoric Acid soluble in water.....	4.69	3.64	15.19	14.08			
Phosphoric Acid insoluble in water.....	10.05	10.50					
Sand, &c., insoluble in acids	1.69	1.52	2.45	2.66			
Phosphate of Lime, equivalent to total } Av.	21.28		31.69				
Phosphoric Acid. }							

Analysis of Bolivian Guano.

Water.....	6.91
Organic Matter containing Ammonia.....	55.52
Common Salt and Sulphate of Soda.....	6.31
Carbonate of Lime.....	3.87
Phosphate of Lime and Magnesia.....	25.68
Silicious Matter or Sand.....	1.71

Johnston.

NOTE.—The guano of the *Lobos Islands* is from 25 to 33 per cent. less valuable than the above.

How to select a good article of guano.

1. The drier the better—there is less water to pay for and transport.

2. The lighter the color the better—it is the less completely dissolved.

3. If it has not a strong ammoniacal smell, it ought to give off such a smell when a spoonful of it is mixed with a spoonful of slaked lime in a wine-glass.

4. When put into a tumbler with water and stirred well, and the water and fine matter poured off, it ought to leave but little sand or stones.

5. When heated to redness over a fire or bright flame, until the animal matter is burned away, the ash should nearly all dissolve in dilute muriatic acid.

6. In looking at the printed analysis (which almost all dealers furnish), see that the per cent. of water is small; that the organic matter containing ammonia approaches to 50 or 60 per cent.; that the phosphates do not exceed 20 per cent., and the common salt and sulphate of soda do not exceed 5 or 6 per cent.—*Johnston*.

How to Apply Guano.—From 200 to 500 lbs. per acre is a proper dressing, the largest quantity being required for the more sterile soils. Mix it thoroughly for a few days with five times its bulk of vegetable mould or loam and some

charcoal or gypsum, after breaking the lumps and sifting in alternate layers. Avoid the use of ashes or lime, as they tend to expel the ammonia. Keep it under cover, beyond the reach of water or rains, until used. It may then be scattered broadcast upon meadows or grain, or placed near the seeds or young plants in the hill.

Analysis of bone (crushed) manure.

In 100 parts, there are of

Lime.....	55.5	Carbonate of Lime.....	3.75
Phosphate of Magnesia.....	2.	Fluoride of Calcium.....	3.
Soda and Common Salt.....	2.5	Gelatine (the substance of horn)	33.25

TABLE, showing the comparative value of animal manures, with farm-yard manure as the standard.

100 lbs. farm-yard manure is equal to

125 lbs. solid excrements of the cow.	3 lbs. Dry Flesh.
73 " " " " horse.	5 " Pigeon's Dung.
91 " liquid " " cow.	15 " Liquid Blood.
16 " " " " horse.	4 " Dry Blood.
98 " mixed " " cow.	3 " Feathers.
54 " " " " horse.	3 " Cow Hair.
36 " " " " sheep.	3 " Horn Shavings.
64 " " " " pig.	3½ " Dry Woollen Rags.

Johnston.

NOTE.—The most powerful substances in the above table, viz., dry woollen rags, horn shavings, cow hair, feathers, &c., hold little or no water, and contain the fertilizing elements of the others in very compact forms. They show less immediate sensible effect upon the crop than the others, because, being so dry and compact, they are long decomposing, but continue to evolve fertilizing matter long after the softer and more fluid manures have spent their force.

Decomposed vegetables as manure.

The *characteristic distinction* between animal and vegetable manures lies in the fact of the former containing a much larger proportion of nitrogen than the latter.

There are two grounds upon which the relative values of different vegetable substances as manures may be estimated. *First*, from the quantity and kind of *inorganic matter* they contain. *Second*, from the proportion of *nitrogen* present in each.

TABLE, showing the relative values of decomposed vegetables as manures, from the inorganic matter they contain.

							Inorganic Matter.	
							lbs.	lbs.
1 ton	Wheat Straw	made into manure	returns to the soil			70 to 360	
1 "	Oat	"	"	"	"	"	100 to 180	
1 "	Hay	"	"	"	"	"	100 to 200	
1 "	Barley	"	"	"	"	"	100 to 120	
1 "	Pea	"	"	"	"	"	100 to 110	
1 "	Bean	"	"	"	"	"	100 to 130	
1 "	Rye	"	"	"	"	"	50 to 100	
1 "	Dry Potato-tops	"	"	"	"	"	400	
1 "	Dry Turnip-tops	"	"	"	"	"	370	
1 "	Rape Cake	"	"	"	"	"	120	
1 "	Malt Dust	"	"	"	"	"	180	
1 "	Dried Seaweed	"	"	"	"	"	560	

Johnston.

TABLE, showing the relative values of decomposed vegetables as manures, from the nitrogen they contain.

100 lbs. of farm-yard manure is equal to

130 lbs.	Wheat Straw	Manure.	80 lbs	Fresh Seaweed	Manure.
150 "	Oat	"	20 "	Dried	"
180 "	Barley	"	26 "	Bran of Wheat or Corn	"
85 "	B'kwh't	"	13 "	Malt Dust	"
45 "	Pea	"	8 "	Rape Cake	"
50 "	Wheat Chaff	"	250 "	Pine Sawdust	"
80 "	Green Grass	"	180 "	Oak	"
75 "	Potato Tops	"	25 "	Coal Soot	"

Boussingault.

NOTES.—The *immediate* effect of vegetable manures in hastening the growth of plants is dependent, in a great measure, upon the quantity of nitrogen they contain, which is given off chiefly in the form of ammonia during their decay in the soil, and may be nearly exhausted in a single season.

Their *permanent* effect and value is to be estimated by the quantity and quality of inorganic matter they contain, or ash they leave when burned, and may not be exhausted for several years.

Besides inorganic matters and nitrogen, there are other ingredients in vegetable manures which are necessary to the sustenance and growth of plants.

Each of the elements present in decayed or decaying plants is capable either of ministering to, or preparing food for such as are still alive.

All refuse vegetable or animal matter on a farm, such as straw, leaves, vegetable tops, chips, sawdust, ashes, dead animals, bones, horns, hoofs, entrails, &c., &c., should be carefully saved and composted, or otherwise made into manure for the use of the farm.

Analysis of a manure heap in the condition usually applied to the field.

Fresh.		Dried at 212°.	
Water.....	64.96	Carbon.....	37.40
Organic Matter.....	24.71	Hydrogen.....	5.27
Inorganic Salts.....	10.33	Oxygen.....	25.52
		Nitrogen.....	1.76
		Ashes (inorganic matter).....	30.05

Inorganic matters.

Soluble in Muriatic Acid.		Soluble in Water.	
Silica.....	27.01	Potash.....	3.22
Phosphate of Lime.....	7.11	Soda.....	2.73
“ Magnesia.....	2.26	Lime.....	0.34
“ Iron.....	4.68	Magnesia.....	0.26
Carbonate of Lime.....	9.34	Sulphuric Acid.....	3.27
“ Magnesia.....	1.63	Chlorine.....	3.15
Sand.....	30.99	Silica.....	0.04
Carbon.....	.83		
Alkali and loss.....	3.14		13.01
	86.99		86.99
		Richardson.	100.00

Analysis of other specimens of fresh farm-yard manures.

	Farm yard Manure From Kent.	Farm-yard Manure From Surrey.
Per centage of Ash.....	9.2	9.6
Silica.....	70.79	71.32
Potash.....	3.32	5.14
Soda.....	0.92	1.68
Lime.....	6.90	12.32
Magnesia.....	0.56	0.82
Common Salt.....	1.43	1.22
Phosphate of Iron.....	2.04	2.03
“ Alumina.....	1.53	2.54
Sulphuric Acid.....	1.89	1.57
Phosphoric Acid.....	1.58	1.27
Manganese.....	trace
	90.96	99.91

Allen & Greenhill.

Composition of fresh farm-yard manure (composed of horse, pig, and cow dung, about fourteen days old). Analysis made November 3, 1854, by Dr. Augustus Voelcker, Professor of Chemistry in the Royal Agricultural College, Cirencester, England:—

MANURES.

339

Water.....	66.17	
* Soluble organic matter.....	2.48	
Soluble inorganic matter (ash)—		
Soluble silica (silicic acid).....	.237	
Phosphate of lime.....	.299	
Lime066	
Magnesia011	
Potash573	
Chloride of sodium.....	.030	
Carbonic acid and loss.....	.218	
	—	1.54
† Insoluble organic matter.....	25.76	
Insoluble inorganic matter (ash)—		
Soluble silica { silicic acid }967	
Insoluble silica { silicic acid }561	
Oxide of iron, alumina, with phosphates.596	
(Containing phosphoric acid, .178)		
(Equal to bone earth, .386)		
Lime.....	1.120	
Magnesia.....	.143	
Potash099	
Soda.....	.019	
Sulphuric acid.....	.061	
Carbonic acid and loss.....	.484	
	—	4.05
		100.00
* Containing nitrogen.....	.149	
Equal to ammonia.....	—	.181
† Containing nitrogen.....	.494	
Equal to ammonia.....	—	.599
The whole manure contains ammonia in a free state.....		.034
“ “ “ “ in the form of salts088

According to this analysis one ton (2000 lbs.) farm-yard manure contains—

Soluble silica (silicic acid).....	24 lbs.
Ammonia (actual or potential).....	15 $\frac{3}{8}$ "
Phosphate of lime	13 $\frac{7}{10}$ "
Lime.....	23 $\frac{7}{10}$ "
Magnesia	3 $\frac{1}{10}$ "
Potash	13 $\frac{1}{2}$ "
Soda	1 $\frac{2}{5}$ "
Common salt.....	$\frac{6}{10}$ "
Sulphuric acid	21 $\frac{1}{3}$ "
Water.....	1323 $\frac{2}{5}$ "
Woody fibre, &c.....	579 "

Of course no two samples of farm-yard manure are exactly of the same composition. That analyzed by Dr. Voelcker was selected with much care, as representing a fair average.

GREEN SAND MARL (OF NEW JERSEY).

Protoxide of iron	15.5
Alumina.....	6.9
Lime.....	5.3
Magnesia	1.6
Potash	4.8
Soluble silica.....	32.4
Insoluble silica and sand.....	19.8
Sulphuric acid.....	.6
Phosphoric acid.....	1.3
Water.....	8.0
Carbonic acid, &c.....	3.8
	<hr/>
	100.0

This is an average of three analyses copied from Prof. Geo. H. Cook's Report of the Geology of New Jersey. According to this estimate one ton (2000 lbs.) of green sand marl contains—

Lime.....	106 lbs.	Soluble silicic acid.	648 lbs.
Magnesia.....	32 “	Sulphuric acid.....	12 “
Potash.....	96 “	Phosphoric acid*...	26 “

To give a better idea of the formation and composition of stable manure, the following is copied from “Waring's Elements of Agriculture” :—

“DIGESTION AND ITS PRODUCTS.

“Let us suppose that we have a full-grown ox, which is not increasing in any of his parts, but only consumes food to keep up his respiration, and to supply the natural wastes of his body. To this ox we will feed a ton of hay which contains organic matter, with and without nitrogen, and soluble and insoluble earthy substances. Now let us try to follow the food through its changes in the animal, and see what becomes of it. Liebig compares the consumption of food by animals to the imperfect burning of wood in a stove, where a portion of the fuel is resolved into gases and ashes (that is, it is completely burned), and another portion, which is not thoroughly burned, passes off as *soot*. In the animal action in question, the food undergoes changes which are similar to this burning of wood. A part of the food is *digested* and taken up by the blood,

* Equal to phosphate of lime $56\frac{1}{2}$ lbs.

while another portion remains undigested, and passes the bowels as solid dung—corresponding to the soot of combustion. This part of the dung, then, we see is merely so much of the food as passes through the system without being materially changed. Its nature is easily understood. It contains organic and mineral matters in nearly the condition in which they existed in the hay. They have been rendered finer and softer, but their *chemical* character (their composition) is not materially altered. The dung also contains small quantities of nitrogenous matter, which has *leaked out*, as it were, from the stomach and intestines. The digested food, however, undergoes further changes which affect its character, and it escapes from the body in three ways—*i. e.*, through the lungs and skin, through the bladder, and through the bowels. It will be recollected from the first section of this book, p. 20, that the carbon in the blood of animals unites with the oxygen of the air drawn into the lungs, and is thrown off in the breath as carbonic acid. The hydrogen and oxygen unite to form a part of the water which constitutes the moisture of the breath.

“That portion of the atmospheric part of the hay which has been taken up by the blood of the ox, and which does not contain nitrogen, is emitted through the lungs. It consists, as will be recollected, of carbon, hydrogen, and oxygen, and these assume, in respiration, the form of carbonic acid and water.

“The atmospheric matter of the digested hay, in the blood, which does contain nitrogen, goes to the *bladder*, where it assumes the form of urea—a constituent of urine or liquid manure.

“We have now disposed of the imperfectly digested food (the dung), and of the *atmospheric* matter which was taken up by the blood. All that remains to be examined is the earthy matter in the blood, which would have become *ashes* if the hay had been burned. The readily *soluble* part of this earthy matter passes into the bladder, and forms the *earthy parts of urine*. The more *insoluble* part passes the bowels, in connection with the dung.

“If any of the food taken up by the blood is not returned as above stated, it goes to form fat, muscle, hair, bones, or some other part of the animal; and as he is not growing (not increasing in weight), an equivalent amount of the body of the animal goes to the manure to take the place of the part retained.*

“We now have our subject in a form to be readily understood. We learn that when food is given to animals it is not *put out of existence*, but is merely *changed in form*; and that in the impurities of the breath we have a large portion of those parts of the food which plants obtain from air and from water; while the solid and liquid excrements

* This account of digestion is not, perhaps, strictly accurate in a physiological point of view, but it is sufficiently so to give an elementary understanding of the character of excrement as manure.

contain all that was taken by the plants from the soil and from manures.

“The SOLID DUNG contains the undigested parts of the food, the more *insoluble* parts of the ash, and the nitrogenous matters which have *escaped* from the digestive organs.

“The LIQUID MANURE contains the nitrogenous parts of the digested food, and the *soluble* parts of the ash.

“The BREATH contains those parts of the fully digested food which contain carbon, hydrogen, and oxygen, but *no nitrogen*, or at least a very inconsiderable quantity of it.”

LIQUID MANURE.

We believe there is no system of enriching the land for small gardens, with a view to perfection of crops, so truly economical and so easily available as that of using liquid manure. We occasionally hear of a gardener, or an amateur grower of some special plant or crop, that has practised enriching with liquids, but it is only occasionally; yet the result of every record is in its favor, and a searching inquiry into any extra production of fruit, flower, or plant almost invariably gives watering with liquid manure as the cause. There is in almost every family a waste of liquids, which usually go into the sewer or drain, or possibly upon the road, where they are of no avail, but if saved by being conducted to a tank, would enrich the entire garden spot of vege-

tables, small fruits, furnish stimulus to the rose and other flower borders, and keep the grass-plot green and fresh even in the hottest and driest weather of midsummer. The use of a little plaster (gypsum) occasionally, thrown in and around the tank, would always keep it sweet and clean. By the use and practice of liquid manuring no delay need ever occur in planting-time because of the manure not being on hand, or not being in a sufficiently rotted condition; but planting could proceed, and the application of manure be made at leisure.—*Horticulturist*.

Value of liquid manures.

The urine voided from a cow during one year contains 900 lbs. solid matter, and compared with Peruvian guano at \$50 per ton is worth \$20. It will manure $1\frac{1}{4}$ acres of land, and is more valuable than its dung, in the ratio, by bulk, of 7 to 6, and in intrinsic value as 2 to 1.—*Dana*.

The Urine of the Cow contains of water.....	92.6 per cent.
“ “ Horse “ “	94. “
“ “ Sheep “ “	96. “
“ “ Hog “ “	92.6 “
“ “ Human “ “	93.3 “

The remainder is composed of salts and rich food for vegetables.—*Sprengel*.

Poudrette and Urate.

Poudrette is the name given to the human excrement after being mixed with charcoal dust or charred peat, to disinfect it of its effluvia, and when dried becomes convenient for use or transportation.

Urate is the name given to urine after mixing with it $\frac{1}{8}$

or $\frac{1}{4}$ of its weight of ground gypsum, and allowing it to stand several days. The urine combines with a portion of the ammonia, after which the liquid is poured off and the remainder dried.—*Allen*.

Analysis of night soil.

The excrement of a healthy man yielded in 1000 parts—

Water.....	733.	Mucilage, fat, and animal matter.	167.
Albumen	9.	Saline matters.....	12.
Bile.....	9.	Undecomposed food.....	70.

Man's urine yielded in 1000 parts—

Water	933.	Phosphate of ammonia.....	1.6
Urea.....	30.1	“ soda	2.9
Uric acid.....	1.	Sal ammoniac.....	1.5
Free acetic acid, lactate of } ammonia, and animal mat- } ter	17.1	Common salt.....	4.5
Mucus of the bladder.....	3.3	Phosphate of lime and magne- } sia, with a trace of silica } and fluoride of calcium.... }	1.1
Sulphate of potash.....	3.7		
“ soda	3.2		
			1000.

Berzelius.

Urea is a solid product of urine, and gives in 100 parts—

Carbon.....	19.99	Hydrogen.....	6.65
Oxygen.....	26.63	Nitrogen	46.65

Prout.

THE DRY EARTH SYSTEM.

It has long been a difficult problem to decide in what way to dispose of human excrement so as to make use of its invaluable ingredients as manure, and, at the same time, to avoid the offensiveness which attends its management in China and Japan, and in all countries where it is habitually applied to the soil.

This problem has at last found a satisfactory solution in the invention of the Rev. Henry Moule, Vicar of Fordington, Dorsetshire, England.

This invention is based on the power of common soil, when dried and sifted, to absorb, not only the moisture of human excrement, but its odor as well.

This power of absorbing odors is due to both the clay and the decomposed organic matter in the soil. It was first discovered, or at least first satisfactorily explained, by Prof. Way, chemist to the Royal Agricultural Society of England, whose interesting experiments on the subject are detailed in the Society's Journal.

It is odd that this easy means of arresting the offensive exhalations of human excrement was not long ago generally adopted. We have a practical illustration of this use of earth in the case of animals of the feline race, whose dejections are extremely offensive. They turn and carefully cover these with earth. In the adhesion of the world to many of the tenets of the Mosaic law, it is strange that we have overlooked the sound advice given in the 12th and

13th verses of the xxiii. chap. of Deut., where we read, "Thou shalt have a place also without the camp whither thou shalt go forth abroad; and thou shalt have a paddle upon thy weapon; and it shall be when thou shalt ease thyself abroad, thou shalt dig therewith and shalt turn back and cover that which cometh from thee."

Mr. Moule's invention is susceptible of many modifications. The apparatus which he has devised, and which is coming into quite general use in England, especially in detached country houses and cottages, where there is no supply of water for water-closets, consists of a hopper-shaped reservoir behind and above the ordinary water-closet seat or holding the supply of dry earth,—this forms a back; a water-tight vessel or vault under the seat; and a mechanical arrangement for measuring out the proper quantity of earth (about a pint and a half) and throwing it forward upon the evacuation, which it entirely covers while it absorbs all the moisture.

This apparatus is simple, inexpensive, not liable to get out of order, and cannot be obstructed by frost.

A modification of the same, still more simple, cheap, and equally effective, though much less convenient, consists of a tub or box (filled with dry earth) at the side of the seat, and a common tin scoop with which to throw the earth upon the deposit. This plan is being generally adopted in the prisons and workhouses of England and the British colonies.

In fact, any vessel containing two inches or more of sifted, dry earth, and a second vessel containing a supply

of earth and a scoop or cup with which to handle it, will answer a good purpose on emergency, and will enable the poorest person not merely to mitigate but to absolutely overcome the most offensive accompaniment of sickness.*

While this invention offers relief from untold misery and annoyance to all who cannot conveniently establish water-closets in their houses, its agricultural importance makes it especially interesting to farmers.

It is a fact too well known to need discussion in our limited space, that of all manures none are at once so powerful and so well adapted to the growth of all crops as "night-soil," or human excrement, though its highly offensive character has generally prevented its use, and has associated with it an idea of degradation. In most parts of the country farm-hands would leave their places rather than to have anything to do with the stuff; and where it is commonly used, it is made a nuisance to wide neighborhoods.

By the aid of the dry earth system every real and fancied objection to its use is done away with. The mixed earth and "soil," when dried and pulverized, are absolutely without other smell than that of freshly turned earth; and, although every atom of fertilizing matter has been retained in a most available form, there is nothing by which, from either appearance or odor, its character could be suspected.

The most remarkable part of the whole matter is, that

* For more particular information on this subject, the reader is referred to a pamphlet entitled "Earth Closets, how to make and how to use them," published by the N. Y. Tribune Association.

when the ordure is once decomposed and (by sifting) intimately mixed with the earth, it has the same quality as any other decomposed organic matter, *i. e.*, it acts as a deodorizer. Consequently, the same earth (by drying and sifting) may be used over and over again, always (at least up to the eighth or tenth time of using) being inodorous and as good a disinfectant as fresh earth; therefore the quantity of earth which it is necessary to prepare and store need not be very large, and it may be made so rich as to be equal to Peruvian guano in its effect on vegetation.

In short, in the opinion of the writer, who has had personal experience in the use of the apparatus, in "sickness and in health," the adoption of the dry earth system is "the coming reform."

TABLE, *showing the comparative increase of corn by different fertilizers.*

No. experiment	QUANTITY OF FERTILIZER.	Yield per acre in bushels.	Increase in bush.	Cost of manure.	Increase of corn for each dollar's worth of manure.
1	No Manure.....	28		\$	bus qrts.
2	500 lbs Superphosphate of Lime.....	46	18	12 50	1 14
3	690 " Guano.....	50 $\frac{1}{2}$	22 $\frac{1}{2}$	19 00	1 8
4	300 " Superphosphate Lime & 640 lbs. Guano	58	30	25 10	1 6 $\frac{1}{2}$
5	320 " Guano and 640 lbs. dissolved Bones ..	51	23	18 40	1 8
6	1040 " Guano & 400 lbs. Superphosphate Lime	74 $\frac{3}{4}$	46 $\frac{3}{4}$	38 60	1 6 $\frac{1}{4}$
7	16 loads Stable Manure.....	35 $\frac{1}{2}$	7 $\frac{1}{2}$	16 00	15
8	32 " "	42	14 $\frac{1}{2}$	32 00	14 $\frac{1}{2}$
9	16 " " & 200 bus. leached Ashes...	44	8	12 00 ^c	22 $\frac{1}{2}$
10	16 " " & 640 lbs. Super P. Lime..	49 $\frac{1}{2}$	14 $\frac{1}{2}$	17 80 ^c	18
11	32 " " & 320 lbs Guano & 1320 lbs Superphosphate Lime }	60	17 $\frac{1}{2}$	16 80 ^c	1 4
12	Hog manure from 108 bus. corn.....	43	15	16 20	30

* Only the increase over the experiments 7 and 8 with stable manure alone.

All tables showing the comparative effect of different manures are of very problematical value. There are so many circumstances and conditions of soil, climate, exposure, moisture, previous treatment of the land, &c., &c.—all of which affect, more or less strongly, the amount of the crop—that it is never possible (in the light of our still imperfect knowledge concerning the growth of plants, and their relations to the soil) to decide how far any increase or decrease may be due to the manure used, and how far to other causes.

TABLE, showing the effect produced upon the quantity of the crop by equal quantities of different manures applied to the same soil, sown with an equal quantity of the same seed.

Manure applied.	Return in bushels from each bushel of seed.			
	Wheat.	Barley.	Oats.	Rye.
Blood	14	16	12½	14
Night-soil.....	..	13	14½	13½
Sheep-dung.....	12	16	14	13
Horse-dung	10	13	14	11
Pigeon-dung	10	12	9
Cow-dung	7	11	16	9
Vegetable manure.....	3	7	13	6
Without manure.....	..	4	5	4

Moisture absorbed by different manures.

1000 parts horse-dung, dried in a temperature of 100°

Fahrenheit, absorbed by exposure to the air at a

temperature of 62° Fahrenheit, moisture, parts 145

1000 parts cow-dung, under same circumstances, “ 130

1000 parts	pig-dung, under the same circumstances,	parts 120
"	sheep-dung, " " "	81
"	pigeon-dung, " " "	50
"	rich alluvial soil, " " "	14
"	fresh tanners' bark, " " "	115
"	putrified " " "	145
"	refuse marine salt, " " "	49½
"	soot, " " "	36
"	burnt clay, " " "	29
"	coal ashes, " " "	14
"	lime, " " "	11
"	sediment from salt-pans, " " "	10
"	crushed rock-salt, " " "	10
"	gypsum, " " "	9
"	chalk, " " "	4

TABLE, showing the number of loads of manure and the number of heaps to each load required to each acre, the heaps at given distances apart.

Distance of heaps apart, in yards.	NUMBER OF HEAPS IN A LOAD.									
	1	2	3	4	5	6	7	8	9	10
3	538	269	179	134	108	89½	77	67	60	54
3½	395	168	132	99	79	66	56½	49½	44	39½
4	203	151	101	75½	60½	50½	43½	37½	33½	30½
4½	239	120	79½	60	47½	39½	34½	30	26½	24
5	194	97	64½	48½	38½	32½	27½	24½	21½	19½
5½	160	80	53½	40	32	26½	22½	20	17½	16
6	131	67	44½	33½	27	22½	19½	16½	15	13½
6½	115	57½	38½	28½	23	19	16½	14½	12½	11½
7	99	49½	33	24½	19½	16½	14	12½	11	10
7½	86	43	28½	21½	17½	14½	12½	10½	9½	8½
8	75½	37½	25½	19	15½	12½	10½	9½	8½	7½
8½	67	33½	22½	16½	13½	11½	9½	8½	7½	6½
9	60	30	20	15	12	10	8½	7½	6½	6
9½	53½	26½	18	13½	10½	9	7½	6½	6	5½
10	48½	24½	16½	12	9½	8	7	6	5½	4½

EXPLANATION.—In the left hand column are placed the distances of the rows and the heaps in each row (*i. e.*, the distances between the heaps in each direction), and at the top of the columns will be noticed the number of heaps intended to be made of each load; the point where the two meet gives the number of loads per acre which will be required for that purpose.

EXAMPLE 1.—Required the number of loads necessary to manure an acre, dividing each load into six heaps, and placing them $4\frac{1}{2}$ yards apart?

SOLUTION.—In the left hand column find $4\frac{1}{2}$ (the distance of the heaps apart), and opposite it to the right, under 6 (the number of heaps in each load), will be found $39\frac{3}{4}$.
Ans.

EXAMPLE 2.—A farmer has a field containing $5\frac{1}{2}$ acres, over which he wishes to spread 82 loads of manure. Now, 82 divided by $5\frac{1}{2}$ gives 15 loads per acre, and by referring to the table it will be seen that the desired object can be attained by making 4 heaps of each load, and placing them 9 yards apart, or by 9 heaps at 6 yards apart, as may be thought most advisable.

NOTES.—A cubic foot of half-rotten stable manure will weigh 56 lbs.; if coarse or dry, 48 lbs.

A load of manure is about 36 cubic feet, and of the first quality will weigh 2016 lbs.; of the second, 1728 lbs.

Eight loads of the first kind spread over an acre will give

108 lbs. to each square rod, and about $3\frac{1}{2}$ lbs. to each square yard.

Five loads will give 63 lbs. to each square rod.

To find the number of loads of manure required to the acre, for a given number of lbs. per square foot.

RULE.—Multiply 43560 (the number of square feet in an acre) by the number of lbs. you wish to spread on each square foot, and divide the product by 2016, and the quotient will be the number of loads required.

EXAMPLE.—Required, the number of loads of manure to cover a 2-acre field, giving 2 lbs. of manure to each square foot?

SOLUTION.— $43560 \times 2 \times 2 = 174240 \div 2016 = 86.4$ loads.

Ans.

ARTIFICIAL MANURES.

It is a self-evident truth that if we sell, we must buy, or we must be content to see our stock on hand reduced.

This principle applies nowhere else with more force than to the stock of mineral plant-food in the soil. This is, after all, our “stock in trade”—ammonia, carbonic acid, and water;—the sources of nearly ninety-nine-hundredths of our crops we can draw from the floating capital of the world, and, except in the case of ammonia, we need give ourselves but little trouble about them. With the mineral matters, however, the case is very different. Some of them, it is true,

are so abundant and so universally distributed that they do not demand much attention ; but some others, on the other hand, have been distributed by nature with so sparing a hand, that our constant care should be given to keeping our supply of them undiminished. They exist only in the soil ; the winds cannot waft them to us, nor do they come, as ammonia does, in every summer shower. They are the hard currency of our banking system, and our business will always be limited by the amount we have in our vaults, and by the promptness with which we make good their loss when we have put them in circulation.

This fact has created a demand for *artificial manures*, the theory of whose production is, that the phosphate of lime which has found its way into the bones of animals, and has thus become, for the moment, unavailable to the farmer, shall be returned by some process which shall convert refuse bones into manure, or that it shall be replaced from some other source, as from the phosphatic guanos from which superphosphate of lime is largely made ; and that potash, lime, &c., shall be collected, in the form of ashes, &c., &c., and returned to the soil.

If all the artificial manures that have been put into the market had been honestly made, the demand for them would have been much greater even than it now is.

But the fact that their composition can be ascertained only by careful chemical analysis, which farmers are incompetent to make, has led to no end of fraud, and one never

knows, in purchasing a ton of superphosphate, poudrette, guano, &c., whether he is or is not paying for half a ton of coal-ashes or other worthless dirt. The consequence of this has been that many farmers have bought a little superphosphate as an experiment, have found no beneficial result from its use, and so have given it up as a bad job and pronounced the whole system of artificial manuring a swindle. The example of each man has had its effect on his neighbors, and there is, consequently, a wide-spread belief that all artificial manures are humbugs.

At the same time, there are so many who do fully understand the value of these fertilizers, and whose land absolutely needs their aid, that the manufacture and sale of such as are of established good quality has reached enormous proportions.

On farms where large stocks of cattle are fed, and for lands which are enriched by the raising of clover as a green crop, the necessity for the use of foreign manures is much less than where the crops are mainly sold off, and no recuperative process (such as the use of green crops) is adopted.

There is, in all fertile lands, a large reserve stock of mineral plant-food which is not yet in a proper condition to be taken up by roots, and if the cropping is not too severe—the produce being mainly consumed at home, and the manure economically used, or the frequent use of green crop manuring being resorted to—the gradual development, in an available form, of these mineral matters will maintain

the land in a fair state of fertility for a very long time, and here the use of mineral manures is less obvious than in other cases.

It is a fallacy, however, to suppose that these lands do not *need* mineral manuring. By the system pursued, we are simply drawing on the capital stock, and, sooner or later, we shall touch bottom. It all looks fair enough now, but at some future day we or our successors must pay the penalty of our improvidence by finding that the land will no longer produce good crops without the use of more purchased manure than can profitably be applied to them.

The only safe rule (the only *honest* course, when we consider the fact that we are only life-tenants of our farms, and are in duty bound to leave them, unimpaired if not improved, to those who are to come after us) is to bring back on to the farm, every year, as much of the more valuable elements of vegetable ashes as we have sold off from it, whether in meat milk, grain, or hay. In this way only can we be sure that our land and our crops will each year improve.

The great deficiency of our older soils is in the items of phosphoric acid and potash. (Lime is more often needed as an agent for the development of matters already contained in the soil than as a direct food for plants.)

While ammonia has been classed among the non-essential elements of manure, its action as a stimulant is so remarkable that it is, commercially considered, the most valuable of all.

Professor S. W. Johnson of the Sheffield Scientific School, Yale College—the highest authority in America—gives the following as the analysis of the best *Superphosphate of Lime* that ever came under his examination:—

*Analysis of Mapes' Improved Superphosphate of Lime.
Manufacture of 1852.*

Water.....	4.54
Organic and volatile matter.....	22.96
Sand and matters insoluble in acids.....	1.48
Soluble phosphoric acid.....	10.65
Insoluble " "	10.17
Ammonia.....	2.78
Phosphate of lime equivalent to phosphoric acid....	45.11

The following is also from Johnson:—

Analysis of Coe's Superphosphate. Manufacture of 1856.

Water, organic and volatile matters.....	38.02
Sand and matters insoluble in acids.....	3.37
Soluble phosphoric acid.....	3.84
Insoluble " "	17.84
Ammonia.....	3.04
Phosphate of lime, equivalent to phosphoric acid...	46.47

Johnson also gives the following analysis of *bone-ash*, or the residue of burnt bones:—

Analysis of Deburg's Bone Meal.

Water.....	3.04
Organic and volatile matters, mostly charcoal.....	2.07
Sand and insoluble matters.....	11.19

ARTIFICIAL MANURES.

359

Lime	42.17
Phosphoric Acid.....	35.42
Carbonic "	1.23
Magnesia and sulphuric acid, with undetermined matters.....	4.88
	<hr/> 100.00

Also the following:—

Analysis of Bone Dust.

Water	8.75
Organic matter.....	27.25
Sand	5.37
Earthy phosphates.....	45.32
Carbonate of lime and loss.....	13.31
	<hr/> 100.00
Ammonia.....	2.98

Also the following:—

Analysis of Fish Guano, or the refuse of Fish Oil Works.

Water	9.67
Organic (animal) matter.....	67.78
Sand	2.05
Lime.....	3.76
Soluble phosphoric acid.....	3.38
Insoluble " "81
Ammonia yielded by animal matter.....	8.36

Purchasers of manures will find the following table—taken

from Judd's Agricultural Annual for 1868—of great value, as affording a good *general* guide in determining the value of manure by the use of an analysis:—

PRICES OF STANDARD FERTILIZERS, AND A STANDARD FOR PRICES.

The prices of some of the standard fertilizers offered in the New York market simply as such, in December, 1867, are as follows:—

Peruvian Guano, in quantities of 50 tons, per long ton, (gold).....	\$60.00
do do in smaller quantities the price varies with the premium on gold; with gold at 35 per cent. prem., per 2000 lbs.....	85.00
Baker's or Jarvis' Island Guano—a phosphatic Guano from the S. Pacific Ocean, which should contain equivalent to 60 to 70 per cent. of bone phosphate of lime; per 2000 lbs.....	
Superphosphate of lime, per 2000 lbs.....	55.00
Bone, fine ground, in 250 lb. bbls., per 2000 lbs.....	45.00
Flour of bone, per 2000 lbs.....	60.00
Fine floated bone, per 2000 lbs.....	65.00
Fish manure, dry and finely ground, per 2000 lbs.....	45.00
do unground, per 2000 lbs.....	30.00
Gypsum or plaster, sold in quantities of 7 bbls., per bbl. (250 lbs)...	1.75
Shell lime, in bulk, per bushel.....	.10
do per bbl.....	1.50
Sulphuric acid of 66 degrees, (oil of vitriol) per lb.....	2½c.
do do of 60 degrees, (pan acid).....	2¼c.

Carboys containing about 150 lbs. of this acid cost \$3 each, and may be returned when empty.

The following table was prepared by John B. Laws, of Rothampstead, England. The money values of the manure resulting from feeding the several substances are based on

the English (gold) prices of manure; they would be considerably higher here, but this does not affect their *relative* value.

Average Composition, per cent. and per ton, of various kinds of Agricultural Produce, &c.

	PER CENT.					LBS. PER (LONG) TON.					Value of manure in \$ & cts. from 1 ton (2000 lbs.) of food.
	Total dry matter.	Total mineral matter (ash).	Phosphoric acid reckoned as phosphate of lime.	Potash.	Nitrogen.	Total dry matter.	Total mineral matter (ash).	Phosphoric acid reckoned as phosphate of lime.	Potash.	Nitrogen.	
1. Linseed cake.....	88.0	7.00	4.92	1.65	4.75	1,971	156.8	110.2	27.0	106.4	19.72
2. Cotton-seed cake.....	89.0	8.00	7.00	3.12	6.50	1,994	179.2	156.8	70.0	145.6	27.86
3. Rape cake.....	89.0	8.00	5.75	1.76	5.00	1,994	179.2	128.8	39.4	112.0	21.01
4. Linseed.....	90.0	4.00	3.38	1.37	3.80	2,016	89.6	75.7	20.7	85.1	15.65
5. Beans.....	84.0	3.00	2.20	1.27	4.00	1,832	67.2	49.3	23.4	89.6	15.75
6. Peas.....	84.5	3.40	1.54	0.96	3.40	1,803	53.8	41.2	21.5	76.2	13.38
7. Tares.....	84.0	3.00	1.03	0.60	4.20	1,852	44.8	36.5	14.8	94.1	16.75
8. Lentils.....	88.0	3.00	1.89	0.96	4.50	1,971	67.2	42.3	21.5	96.3	16.51
9. Malt dust.....	94.0	8.50	5.23	2.12	4.20	2,106	190.4	117.1	47.5	94.1	18.21
10. Locust beans.....	85.0	1.75	1.25	1,904	39.2	28.0	4.81
11. Indian meal.....	88.0	1.20	1.13	0.35	1.80	1,971	29.1	25.3	7.8	40.3	6.65
12. Wheat.....	85.0	1.70	1.87	0.50	1.80	1,904	38.1	42.0	11.2	40.5	7.08
13. Barley.....	84.0	2.20	1.35	0.55	1.65	1,882	49.3	20.2	12.3	37.0	6.12
14. Malt.....	95.0	2.60	1.60	0.65	1.70	2,128	58.2	35.8	14.6	28.1	6.65
15. Oats.....	86.0	2.85	1.17	0.50	2.00	1,926	63.8	26.2	11.2	44.8	7.70
16. Fine pollard*.....	86.0	5.60	6.44	1.46	2.60	1,926	125.4	144.2	32.7	58.2	13.53
17. Coarse pollard†.....	86.0	6.20	7.52	1.49	2.58	1,926	128.9	168.4	33.4	57.8	14.26
18. Wheat bran.....	86.0	6.60	7.95	1.45	2.55	1,926	147.8	178.1	32.5	57.1	14.59
19. Clover hay.....	84.0	7.50	1.25	1.30	2.50	1,882	168.0	28.0	29.1	56.0	9.64
20. Meadow hay.....	84.0	6.00	0.88	1.50	1.50	1,882	134.4	19.7	38.6	33.6	6.43
21. Bean straw.....	82.5	5.55	0.90	1.11	0.90	1,848	124.3	20.2	24.9	20.2	3.87
22. Pea straw.....	82.0	5.95	0.55	0.89	1,837	133.3	19.0	19.9	20.2	3.74
23. Wheat straw.....	84.0	5.00	0.55	0.65	0.60	1,882	112.0	12.3	14.6	13.4	2.68
24. Barley straw.....	85.0	4.50	0.37	0.63	0.50	1,904	100.8	8.3	14.1	11.2	2.25
25. Oat straw.....	83.0	5.50	0.48	0.93	0.60	1,850	123.2	10.7	20.8	13.4	2.90
26. Mangel wurzel.....	12.5	1.00	0.09	0.25	0.35	280	22.4	2.0	5.6	5.6	1.07
27. Swedish turnips.....	11.0	0.68	0.13	0.18	0.22	246	13.4	2.9	4.0	4.6	0.91
28. Common turnips.....	8.0	0.68	0.11	0.29	0.18	179	15.2	2.5	6.5	4.0	0.86
29. Potatoes.....	24.0	1.00	0.22	0.43	0.35	537	22.4	7.2	9.6	7.8	1.50
30. Carrots.....	13.5	0.70	0.13	0.23	0.20	302	15.7	2.9	5.1	4.5	0.80
31. Parsnips.....	15.0	1.00	0.42	0.56	0.22	336	22.4	9.4	8.1	4.9	1.14

* Middlings, Cannelle.

† Shipstuf.

TILE DRAINING.

I have preferred to head this article as I have, rather than to say simply "draining" or "under-draining," because I believe in the use of tiles under all circumstances when it is possible to procure them, and because the making of stone drains is understood by every farmer who lives in a region that is blessed with wet land and stone.

At the same time, I would not be thought to undervalue the usefulness of stone drains. Neither the stone nor the tile has any influence, in itself, on the fertility of the soil. Any material by the use of which we can make a passage-way through the soil will make a perfectly good drain, *as long as it keeps the passage open.*

The question is to be decided simply by the consideration of cost and durability; and here the tiles have an immense advantage.

In the first place, they are very much *cheaper* than stone; and in the second, the drain which they make is very much more likely to be permanent.

It will, I am aware, strike many farmers whose land is encumbered with stones, as a singular proposition that it is cheaper to pay twenty-five or thirty dollars per acre for tiles, when there are stones on the place that it would be an advantage to get rid of. But it is a fact, nevertheless. The

cost of collecting the stones, of breaking (or selecting them) to a proper size, of laying them in the drain, and of protecting them from the rattling down of loose dirt among them, and from the burrowing down of field-mice, is very great, and in addition to this we have to calculate the cost of digging the very much wider ditch that is required for their use.

To drain land in the best manner there are required about sixty rods of drain four feet deep, and fifty cents a rod for the above items (which is the utmost that tile should cost) would not pay one-half of the actual cost of stones, if we calculate the labor of teams and men at anything approaching their full value.

As to durability. A tile drain, when properly laid, is packed closely in the most compact subsoil within our reach, has its joints (which are very close) encased in an earthen collar, is closed at its upper end by a flat stone against the tile, and its outlet secured by a grating. No dirt can get in to stop it up, and no vermin can use it for a camping ground. The only thing (except in rare instances the roots of trees) that can enter it at all is the water that it is intended to carry away.

Of course I speak of a tile drain that is made of good materials and is made in a proper manner. It is very easy to make a drain that will not be worth the cost of the tiles, not worth anything; and many such drains are made by careless or ignorant people, who, seeing their uselessness,

are loud in the praise of stone drains, and never want to see another draining tile so long as they live.

A *good* tile-drain, made of good clay and well burnt, properly laid on a uniform descent, *and having a good outlet*, is practically as permanent as the earth in which it is imbedded.

And now, *how to make such a drain*. It would take much more than the few pages that can be here devoted to the subject to tell. All that my space will allow me to do is to give a few general rules and directions, which will suffice to enable a farmer to understandingly decide for himself whether he will make his drains of stones or of tiles; and a few arguments which may convince him that he *cannot afford* to let his wet land go undrained.

The draining tile is made in several forms, known as the "round," the "sole," and the "horse-shoe." The last mentioned represents the first step that was taken in advance of the use of stones, and it has long been condemned as an inferior article by all who have had experience in the use of the other kinds. The sole-tile, which has an egg-shaped orifice, and has a flat side to lie upon, is theoretically very good, and is really very good, only not the best. The flat side is a delusion, for the reason that it generally is *not flat*, being very liable to be warped out of shape in the burning, while the uneven drying of the clay before it is burnt, or the friction of the die through which it is moulded, is very apt to so distort its shape as to make it difficult to make a good joint.

The round tile, if well made, is much better, is practically perfect. A tile does not need a flat side to lie upon, for in nine cases out of ten the bottom of the ditch is not flat, and as soon as each piece is put in its place, and while it is held there by the tile-layer, a second man covers it sufficiently to hold it firmly. The smaller sizes have collars or rings to fit them, and these keep the joints "in line" and prevent loose dirt from rattling into the wider openings. Another great advantage of the round tiles is that, if they don't fit each other as they are first laid, they can be turned over until the slight inequalities of the two ends will correspond.

All of the larger tile makers now make the round tiles, and most of them make them very well. A machine invented by Mr. Tiffany (of the Crosmann Clay and Manufacturing Company, Woodbridge, New Jersey) moulds the tiles more smoothly, and presses them harder, than any other yet brought into use. Mr. C. W. Boynton, of Woodbridge, however, seems to have brought more real talent to the manufacture of tiles than any one else who has undertaken the business, and his pipes are probably the best now made, inasmuch as they are two feet long—twice the usual length—and are supplied with connecting pieces for admitting lateral drains into the main trunk lines. Heretofore it has been the custom to pick a hole in the side of the tile of the main drain, and to bring the end of the lateral against it, closing the irregular openings by covering them with bits of broken tile or small stones; and it was nice work to

avoid breaking the pipe, and at the same time to make the joint so accurately as to neither retard the flow nor to admit earth from the filling.

Boynton's pipes, which are shown in the accompanying cuts, have a branch piece nicely fitted to the side of the pipe that is to form a part of the main, the branch forming a part of the lateral. On the end of this branch a collar may be placed to receive the end of the lateral, making as good a joint at the junction as at any other part of the drain.

Before this improvement was made, it was often necessary, where a tile came into the main, to make a silt-basin to catch any silt that might be deposited by the more sluggish flow of the water at that point. By its aid these silt-basins may be, in nearly all cases, dispensed with, as the lateral enters in an oblique direction, and the velocity of its flow will be imparted to that of the main.



FIG. 1.

FIG. 2.



FIG. 3.

Fig. 1 shows the round tile; Fig. 2, the collar; Fig. 3, the manner of laying these; Fig. 4, the connecting joint of the

main with a branch to receive the lateral; and Fig. 5 the




FIG. 4.

manner of laying the tiles at the junction of a lateral drain with the main.




FIG. 5.

Rules to be observed in making Tile Drains:—

1. Every drain (unless there is some special reason to the contrary) should run directly down the steepest descent of the land—not obliquely, but straight down the hill.
2. Wherever possible, the drains should be four feet deep, *especially* when the subsoil is a stiff clay hard-pan.
3. When the drains are four feet deep, they should be forty feet apart. If only three feet deep, they should be only twenty feet apart; and if more than four feet, they may safely be placed at greater distances than forty feet.

4. The rate of fall or inclination of a drain should not decrease as it approaches the outlet. It may be increased as much as is convenient. The rule is, to keep the water running faster and faster, rather than slower and slower, as it gets on in the drain.

5. The outlet should always be clear and free—never, if it can possibly be avoided, so arranged as to be obstructed by mud or dead water.

6. The tiles should have no porous material of any kind over them, but should be imbedded (and firmly packed) in the closest clay that is accessible.

7. In digging the ditch, always commence at the lower end and work toward the top; in laying the tiles, commence at the upper end, and continue toward the outlet.

8. Never have tiles laid by the piece (or rod), but always by the day, and by the most faithful and careful man that can be found; if possible, do it yourself, and remember that the golden rule of draining is that, *as the weakest link of a chain is the measure of its strength, so is the worst laid tile of a drain the measure of its goodness.**

If the drains are laid at distances of forty feet it will take just about one thousand feet of tiles to drain an acre.

As to the sizes of tiles required, it will make a difference whether the fall is rapid or slight; but under all ordinary circumstances, where there are no springs to be disposed of, only the natural drainage of the land itself (its accumulated

* Talpa, or the Chronicles of a Clay Farm.

rain-fall), the first 1500 feet in length, whether it be a single drain or several laterals, may be made of the smallest sized tiles ($1\frac{1}{2}$ inch). Beyond this amount and up to 5000 feet, 2-inch tiles will suffice. From 5000 to 10,000 feet use 3-inch, and from 10,000 to 20,000 feet use 4-inch.

These sizes would not suffice for the immediate removal of all the water of a very heavy rain-fall, but it is to be remembered that before the water can get to the tiles it must filter slowly through four feet of soil, and could reach the drain but slowly, were it ever so large. Then again, it is not important that the water of a heavy rain be removed within an hour of its falling; it does no harm to have it settle slowly away, so long as it really does settle away, and does not stand to be evaporated from the surface, nor to flow off over it; and it is desirable that the drains should occasionally run "more than full," so that a strong flow of water may wash out any obstructions that may have accumulated in them.

The question should not be so much how large a tile is necessary to carry the water, as how large a tile will the water (after heavy rains) be able to flush and keep clean.

In the foregoing, I have simply stated rules and principles which have been proven by long experience to be correct. The evidences of their truth and reliability, and the arguments on which they are founded, could not be set forth in the limited space which has been allowed for the subject in this book. The object here is to set forth rules and to give




FIG. 6.

Tools used in laying drain tile.

directions. Those who are desirous of investigating reasons will find them stated in other works which are devoted to the fuller discussion of the various topics here touched upon.

The ditches are usually dug, in this country, with the ordinary pick, spade, and shovel, with the single addition of a narrow scoop to work in the narrow bottoms of the drains. Such a scoop may be made by cutting a common, round-pointed, long-handled shovel down to a width of four or five inches.

In Europe, where much more extensive operations of drainage are carried on than are known in this country, sets of tools especially adapted for all the different operations are used. One set of these is shown in Fig. 6.




FIG. 7.

The position of the workman in cutting a narrow ditch

for a tile, or rather in finishing the bottom of the ditch with the scoop, is shown in Fig. 7.




FIG. 8.

The manner of securing the outlet so as to keep out ver-




FIG. 9.

min, and, at the same time, to prevent the earth from caving in about the end of the drain, is shown in Fig. 8.

The manner in which draining tiles are moulded from moist clay may be learned from Fig. 9, which represents a strong wooden box filled with clay, which, by the pressure of a lever, is forced out through holes which have the shape of the outside of the tile. A plug stands in the middle of each hole (supported from within, so that the clay can entirely surround it as it comes out), which makes the bore of the tile.

WHY SHOULD LAND BE DRAINED?

There is one condition of soil that is the most favorable for the growth of nearly all agricultural plants—that is a condition of porousness, moisture, warmth, and aëration. The roots of plants need to be in a *dark* place, to be *surrounded by moisture* (this is very different from being soaked in water), and to be sufficiently supplied with air.

There are other conditions of fertility, such as richness in plant-food, &c., which, although of the utmost importance, are apart from our present subject. What we have now to do with is the *mechanical* state of the soil, as distinguished from its chemical composition and action—that is to say, with its moisture, its temperature, the ease with which roots can penetrate it in search of nutriment, and the opportunity for the admission of atmospheric air to their vicinity.

The effects of drainage on the chemical constitution of the soil, and on the chemical action of its ingredients as

affecting vegetation, is very great ; but it is not necessary to the strength of the argument that they should be detailed here, and their sufficient discussion would require too much space.

Moisture.

By the moisture of the soil we mean a condition resembling that of a sponge which has been dipped in water and then lifted out and allowed to drain. While in the water it was saturated—that is, all of its pores were *filled* with water—but on being removed the water all runs out from its pores, except the small amount that adheres (by capillary attraction) to its substance.

In like manner the undrained soil, after a heavy rain, is saturated. All of the spaces between its particles are filled with water. After draining, this water all passes away, except the small amount which adheres to the surfaces of the particles, and that which fills the more minute pores of these particles. There is enough water in the soil in this condition to supply the demands of plants ; but there is not—as there was before draining—so much as to interfere with their healthy growth.

Not the least beneficial effect of draining is that which is the result of the admission of air to its lower and cooler parts, causing a deposit of moisture in dry weather, which is sufficient to supply the needs of vegetation, and to greatly mitigate, if it does not even entirely overcome, the effects of drought.

That land should be made damper by being made more dry, that under-draining should be one of the best preventives of the ill effects of drought—this is the apparently anomalous proposition on which one of the strongest arguments in favor of draining is based.

When we see a field baked to the consistence of a brick, gaping open in wide cracks, and covered with a stunted growth of parched and thirsty plants, it seems hard to believe that the simple laying of hollow tiles, four feet deep, in the dried-up mass, would do anything at all toward the improvement of its condition; for the present season it would not, but for the next it would, and for every season thereafter, and in increasing degree, so long as the tiles continued to act as effective drainage.

The baking and cracking, and the unfertile condition of the soil are the result of a previous condition of entire saturation. Clay cannot be moulded into bricks, nor can it be dried into lumps unless it is first made soaking wet. Dry, or only damp clay, once made fine, can never again be made lumpy, unless it is first made thoroughly wet, and is pressed together while in its wet condition. Neither can a considerable heap of pulverized clay, kept covered from the rain, but exposed to the sun and air, ever become even apparently dry, except within a few inches of its surface. After under-draining has had time to bring the soil, to a depth of two or three feet, to a thoroughly drained condition, it will equally prevent it from being baked into lumps, or

from becoming, for any considerable depth below the surface, too dry for the purposes of vegetation. In the first place, the water of heavy spring rains, instead of lying soaking in the soil until the rapid drying of summer bakes it into coherent lumps, settles away and leaves the clay, within a few hours after the rain ceases, and before rapid evaporation commences, too much dried to crack into lumps.

The other direct effect of under-draining is to remove from below, water which, if not so removed, would be evaporated from the surface.

The formation of a crust on the surface of the ground is in direct proportion to the quantity of water that is removed by evaporation, and the crust constitutes a barrier against the admission of air. Consequently the larger the quantity of water that is removed by the drains, the smaller is the obstacle offered to the entrance of air. The more constantly the lower parts of the soil are relieved from excess of water and supplied with air, the more deeply will roots descend; and the more frequently will the air in the lower soil be changed, the easier its communication with the atmosphere.

On these two principles depends the immunity from drought which under-draining helps us to secure. In dry weather the soil gets its moisture from the deposit of dew, on the surface during the night, and on the surfaces of the particles of the lower soil constantly, day and night.

Temperature.

The temperature of the soil is a matter of the utmost consequence. Seeds cannot germinate, and plants cannot grow without there being a certain amount of heat in the soil, and there is no means by which this heat is so much and so constantly reduced as by the evaporation of water from its surface. In proportion as we remove by the means of under-draining the water which would, if not so removed, remain to be evaporated, we allow the soil to attain a higher temperature, and so to become more productive.

The penetration of roots.

In a soil that is usually *too wet*, the roots of plants confine their operations to the few inches of dry soil at the surface, as they will not push into a cold, compact, wet subsoil. Draining removes the water from the subsoil, allows it to become sweet and warm and loose, and fit for the entrance of roots, which are thereby enabled to seek farther for a greater quantity and a greater variety of food.

The circulation of air.

Atmospheric air, if not absolutely necessary to the life and action of the roots of plants, greatly favors their growth and their absorption of food. Aside from its direct supply of carbonic acid to the feeding parts of the roots, it brings moisture to the soil by which they are surrounded, and aids in preparing its nutrient constituents for assimilation.

ROTATION OF CROPS.

The experience of practical farmers very early demonstrated the necessity for adopting a system of changes in the crops grown on the same soil. Thus, we find in the writings of Columella, Varro, Theophrastus, and others who in ancient times wrote on the subject of agriculture, distinct rules laid down as to the course of cultivation to be pursued in order to prevent the exhaustion of the soil, or, rather, to prevent it from failing to produce a particular crop so long as it was fertile for anything, and to enable it to make full use of whatever manures were applied to it.

In more modern times, the *reasons why* rotations are necessary have been, in a measure, explained by the aid of chemistry, but we have not materially improved on the *practice* of those who cultivated the soil 2000 years ago.

The various crops appropriate different elements from the soil, or the same elements in different proportions. Of course, by raising the same crop year after year from the same field, its quantity and quality not only yearly deteriorate, but the soil becomes exhausted of the special ingredients which go to support the growth of that particular product, while it accumulates the elements especially adapted to some other crop.

The principle on which rotations are based may be readily understood from the following illustration :—

What are known as the root crops contain, in their ashes, a very large proportion of potash. The average amount of this substance contained in the ash of potatoes, turnips, beets, and carrots, is fully *fifty per cent.* of the whole ; that is, they contain as much of this single ingredient as of all the other mineral ingredients combined. Wheat, rye, oats, and barley, on the other hand, contain an average of only *twenty-five per cent.*, or only one-half as much of this as of all the other ingredients.

If we examine their content of phosphoric acid, however, we shall find the case quite different. For instance, the four root crops above named contain an average of only about *thirteen per cent.* of this element, while the four grain crops contain an average of about *thirty-seven per cent.*

Again, lime forms but about *three per cent.* of the ash of most root crops, while it exists in clover and most of the fodder plants to the extent of about *thirty-five per cent.* of their ash.

If we were to follow through the whole range of the mineral constituents of our crops, we should find similar variations in the amounts appropriated by the different plants which are commonly grown on our fields.

Now, suppose that on a field of average quality we find that wheat or some other grain grows to advantage. Stimu-

lated by the profits of the cultivation of this grain, we continue to grow it year after year, without intermission. The result is that—sooner or later, often within two or three years—we find the yield steadily diminishing. One reason for this is that we have been constantly robbing the soil of undue amounts of phosphoric acid, and (without rendering it unfertile for some other crops, such as potatoes) we have seriously impaired its capacity for the production of wheat. If, instead of raising wheat the second year, we had raised potatoes, or clover, or some plant of an entirely different character from wheat, we should have drawn more evenly on *all* of the resources of the land, and should have postponed the exhaustion of its stock of available phosphoric acid.

Here then comes in play, also, another element which it is necessary for the farmer to consider, namely :—there are constantly going on in the soil (which may be considered a natural chemical laboratory) certain chemical and mechanical processes, whose effect is to continually set free from other combinations and prepare for the use of plants the various minerals which constitute their ashes. Therefore, if we bring a grain crop into the rotation only once in four, five, or six years, the simple action of these processes will, in the intervening time, set free enough phosphoric acid for a second crop. Soils differ, not only in their composition, but in the rapidity with which their elements are set free; consequently we find some soils on which the same crop may

safely be tried every second or third year, and others on which we must allow a much longer interval.

The same rule that applies to the soil holds good also with regard to manures. These almost always contain various matters which go to feed plants, and we must study to so arrange our crops as to make profitable use of all that they can yield; and, if they are of a sort to need time and the action of the chemical and mechanical influence of the atmosphere and of the soil for the complete development of all of their constituents, we must adjust our crops, so far as possible, to take up these constituents as they are prepared for use.

The foregoing is the basis of the chemical theory of rotations.

In addition to this, we must consider the influence exerted on the soil by the roots which are left in the ground when the crop is removed. This element of the influence which plants exert on plants which are to follow them in the same soil is especially important in the case of clover, which is so active in its fertilizing effect, that it may be assumed that we have overcome our great difficulty in bringing up a poor soil when we have enabled it to grow a good crop of clover. One especial virtue of this plant is that it sends its roots far into the subsoil, and thus appropriates, by means of its vigorous feeding powers, useful materials which were out of the reach of the roots of plants of other species. These materials are deposited in the substance of the plant, and (on its decay

when ploughed in, or on the decay of its roots when these alone are left in the soil) they are presented to the new crop in a most acceptable form. The raising of other green crops to be ploughed in for manure, is advantageous for the same reasons.

Two most valuable accessions to the rotation of crops will be found in the root crops, and in green forage crops to be either cured for winter use or fed to animals kept on the "soiling" system. To these crops the richest animal manures may be profitably applied, and, while they will make a most luxuriant growth, they will "draw the fire" of the manure, and leave the land in the best condition for the growth of grain crops.

Copeland says:* "When it was discovered that roots of all kinds were not only good food, but the best food for cattle, those farmers who believed in the discovery cultivated roots, and found, not only that their value as food was inestimable, but that, with a given expenditure in manure and labor, roots gave a larger return in value than any other crop. This was the turning point, the rising tide-wave of improving agriculture. The new crop was an improvement in every respect. It restored fertility better than the fallow, gave an immense amount of fodder, and insured a corresponding increase in manure, from the greater number of cattle which could be fed from the farm.

"Under the old system—the same pursued in New Eng-

* Country Life, page 435.

land at the present day—there was a large and a small white crop, one large yield of hay, then smaller and smaller, then good pasture, then poor. This rotation gave a change from better to worse. The new practice demonstrated that there need be no “worse.” It showed that a root crop should follow the sod and should be followed by grain; that again by grain or grass and clover; that by pasture and roots. At first it was made a point that a white crop should never be taken two years in succession, and after going through roots and grass it was found, on returning to the white crop, that the ground was so much richer than before, that a number of bushels was taken previously unheard of in the neighborhood.”

Liebig says:* “The succession of crops in rotation is always made dependent upon the cereals; the preceding crops are selected of such a kind that their cultivation will not injure, but rather improve the succeeding corn crop. The selection of the particular kind, however, is always governed by the condition of the soil. In a field abounding in stalk and leaf constituents, it is often found useful to have wheat preceded by tobacco or rape, rye by turnips or potatoes, since these plants, by drawing from the soil a large amount of leaf and stalk constituents, serve to restore a more suitable proportion between the straw and corn constituents for the future cereal crop, and, at the same time, to diminish in the arable soil those conditions which favor the growth of weeds.

* The Natural Laws of Husbandry, page 227.

Prof. James F. W. Johnston says :* “Two practical rules are suggested by the fact that different plants require different substances to abound in a soil in which they shall be capable of flourishing.

“1. To grow alternately as many different *classes* or *families* of plants as possible, repeating each class at the greatest convenient distance of time. In this country (England) we grow, chiefly, root crops—corn plants refined for seed—leguminous plants, sometimes for seed (peas and beans), and sometimes for hay or fodder (clover and tares), and grasses ; and these in alternate years.

“Every four, five, or six years, therefore, the same class of plants comes round again, and a demand is made upon the soil for the same kinds of food in the same proportion.
* * * * * A perfect rotation would include all those classes of plants which the soil, climate, and other circumstances allow to be cultivated with a profit.

“2. A second rule is, to repeat the same *species* of plants at the greatest convenient distance of time. * * * * *

“Instead, therefore, of a constant repetition of the turnip every four years, theory says, make the carrot or the potato take its place now and then, and instead of perpetual clover, let tares, or peas, or beans occasionally succeed to your crops of corn.†

* Agricultural Chemistry, page 493.

† “*Corn*, in English agricultural writing, is a general term corresponding to our *grain*.”

"The land loves a change of crop because it is better prepared with that food which the new crop will relish than with such as the plant it has long fed before continues to require.

"It is for this reason that new species of crop or new varieties, when first introduced, succeed remarkably for a time, and give great and encouraging returns. * * * *

"It is constant variety of crops which, with rich manuring, makes our market gardens so productive, and it is the possibility of growing in the fields many different crops in succession that gives the fertility of a garden to parts of Italy, Flanders, and China."

The rotation to be adopted may be best selected by each farmer for himself—keeping in mind the foregoing principles—with reference to his soil, his market, his climate, the price and supply of labor in his neighborhood, and the extent to which he can accumulate manure.

The rotation which the writer has adopted for his own farm is the following:—

First year :—Indian corn, on sod land, manured the previous autumn with the entire accumulation of manure in the barn cellar, then ploughed and left in the rough furrow for the fullest exposure to frost, harrowing thoroughly before planting time.

After the crop is taken off in the fall, the land to be ploughed and again left in the rough furrow to winter.

Second year :—Roots, the ground being properly divided between carrots, mangel wurzel, turnips, and parsnips.

For this crop the land is cross-ploughed in the spring, dressed with one-half of the winter's accumulation of manure in the cellar, and from 100 to 250 lbs. of superphosphate of lime, both sowed broadcast on the furrow and thoroughly harrowed in.

Third year :—Green forage crops for "soiling" cattle—mainly oats and Indian corn in successive sowings.

These crops receive the balance of the winter's manure, and a good-portion of the land is cleared off in time for winter rye to be sown.

Fourth year :—The winter rye is cut green, very early in the season, for "soiling" the cattle, and on the land not occupied with it a crop of green fodder is grown that can be got off by August 1st.

In the early autumn the land to be sown to wheat, and seeded down with timothy and clover.

Fifth year :—The grain harvested and the growth of grass and clover left on the land.

Sixth year :—Two cuttings of hay to be taken off, and the land to be manured and ploughed in the fall for the succeeding crop of corn, with which the rotation recommences.

PROPERTIES AND COMPOSITION OF MILK, BUTTER, &c.

Composition of Milk in 1000 parts.

Water.....	840
Casein.....	40
Milk-sugar.....	45
Butter, or oil.....	40
Phosphate of lime.....	17
Phosphate of magnesia.....	4
Chloride of potassium.....	9
Common salt.....	2
Free soda.....	3
	1000

NOTE.—Milk is heavier than water in the proportion of 103 to 100.

The rapidity with which cream rises to the surface depends upon the temperature to which it is exposed.

New milk, set aside, will cream in

36 hrs. if the temperature of the air is 50° Fahrenheit.					
24	"	"	"	55°	"
18 to 20	"	"	"	68°	"
10 to 12	"	"	"	77°	"

At a temperature of 34° to 37°, it may be kept two to

three weeks without throwing up any noticeable amount of cream.

Cream contains the greater part of the fatty matter of the milk, a small portion of the curd, and considerable water.

Good cream, when skilfully churned, will yield about one-fourth of its weight of butter.

The temperature at which *milk* can be churned most economically is 65° Fahrenheit.

The temperature at which *cream* can be churned most economically is at 58° Fahrenheit.

Butter contains more or less of all the ingredients of the milk. Essentially it consists of the fat of milk mixed with about one-eighth of its weight of water, a small quantity of casein or curd (cheesy matter), and of saline matter. The casein seldom exceeds two per cent. of the whole weight.

The *fat of butter*, when solidified by pressing out the oil, is identical with the solid fat of the human body.

The *oil of butter* is a peculiar kind of fat not hitherto detected in any other substance.

These two ingredients vary considerably with different samples ; hence the different degrees of *hardness* which different samples present. The solid fat abounds more in winter ; the liquid fat more in summer. They are in about the following proportions in 100 parts :—

	Summer.	Winter.
Solid fat.....	40	65
Oil of butter.....	60	35

The main cause of butter becoming rancid is the chemical decomposition which the casein or curd it contains undergoes by exposure to the air. This chemical change in the cheesy matter may be prevented—

1st, By thoroughly washing and salting before the cheesy matter has had time to become altered by exposure to the air ;

2d, By taking care that any water that may remain in or around the butter be kept perfectly saturated with salt ;

3d, By carefully excluding the air from the vessel in which the butter is packed.

About half a pound of the best Ashton salt is used to 10 pounds of butter.

Milk contains a peculiar kind of sugar called milk-sugar, which, being highly soluble in water, passes off in the whey and goes to fatten pigs. In some countries it is extracted and made an article of commerce.

The main cause of milk becoming sour is the chemical change which this sugar undergoes, without fermentation and therefore without loss, into an acid called lactic acid.

This lactic acid is the cause of the *curdling* of the milk, which may be hastened by hastening the change of the milk-sugar into lactic acid by the addition of any other acid, such as vinegar or rennet.

Pure casein is nearly insoluble in pure water, either by boiling or otherwise. By adding, however, a little soda to the water, it dissolves and returns to its milky condition ;

when, by adding some more milk-sugar (or lactic acid), it again curdles.

The milk of nearly all animals contains the same ingredients. The best known varieties consist nearly of—

	Woman.	Cow.	Ass.	Goat.	Ewe.
Casein.....	1.5	4.5	1.8	4.1	4.5
Butter.....	3.6	3.1	0.1	3.3	4.2
Milk-sugar....	6.5	4.8	6.1	5.3	5.0
Saline matter..	0.5	0.6	0.3	0.6	0.7
Water.....	87.9	87.0	91.7	86.7	85.6
	<hr/> 100.	<hr/> 100.	<hr/> 100.	<hr/> 100.	<hr/> 100.

The butter and cheese producing quality of milk is shown by the following

TABLE.

100 lbs. milk contains about	3	lbs. pure butter.
100 lbs. “ “ “	7.8	lbs. “ cheese.
100 lbs. “ averages “	3.5	lbs. common butter.
100 lbs. “ “ “	11.7	lbs. “ cheese
100 lbs. skim-milk yields “	13.5	lbs. skim-milk “
1 qt. wine measure weighs	35	oz.
1 qt. milk “ “	41	oz.

The milk of different cows varies much in richness. We have known one from 65 lbs. of whose milk were made 64 oz. of butter. A full milk cheese contains about 33 per cent. of water, and a skim-milk cheese about 60 per cent.

Butter at 50 cents per pound will yield about as much profit as cheese at 15 cents, making no allowance for the value of skim-milk over whey.

BUTTER AND CHEESE-MAKING.

The Butter Dairy.—The quality of butter doubtlessly depends more upon the manufacture than upon all other causes combined, yet it is true that the cows, the grass or food, and the water, have much to do with the delicacy of its flavor and richness of its color. It is a notorious fact that eight-tenths of the butter that is sold in the market brings from *five to fifteen* cents per pound less than it would have done had it been properly manufactured. Factory cheese for the same reason brings from *three to eight* cents per pound more than dairy. It costs no more to make a good article than an inferior one, and when this fact is fully appreciated, thousands of dollars will be saved annually to the dairyman farmer.

Milk-room.—The best milk-room is one through which a stream of pure spring water flows, and a reservoir under the “pan rack” is very desirable. When this cannot be had, select a room or building on the north side of the house, through which fresh air can freely circulate. If a cellar is chosen, it should be dry and thoroughly ventilated by large latticed windows and doors. No decaying vegetables should be allowed to remain in it, as the milk and cream easily become tainted. Close and damp cellars are

entirely unfitted for a milk-room, and should not be used. The temperature of the milk-room should be as uniform as possible, ranging from 55° to 65° . When the weather is cold, a fire should be kept in a stove on which a basin of pure water is placed, to prevent the air from becoming so dry as to form a crust on the cream. When too warm the temperature can be reduced by hanging wet linen sheets near the doors and windows, the lower edges of which dip into a vessel of water.

Cleanliness.—In every department of butter-making the utmost cleanliness should be observed. Milk and cream rapidly absorb noxious gases, and are especially affected by the acids and gases which arise from the decomposition of sour milk or cream. Every utensil used in connection with the dairy should be scalded every time used in boiling water, in which, occasionally, a small piece of bicarbonate of soda has been dissolved. All traces of milk or cream accidentally spilled on the floor should be carefully removed.

Setting the Milk.—As soon as the milk is drawn from the cow it should be strained into the setting pans, to a depth of not over two inches. The complete raising of the cream, especially in warm weather, is thus greatly facilitated. In summer the temperature of the milk should be reduced as soon as possible to about 62° . Powdered ice put into the pail before straining is best; setting the pail in cold spring or well-water for a few minutes will answer. A small piece of crystallized soda about the size of a common acorn,

dissolved in a little water, put into each pail of milk before straining, to correct the acidity as it is formed, will increase the quantity of cream, and improve the quality of the butter. Milk, if kept at the proper temperature, need not stand over thirty-six hours. If the cream does not rise in that time, the quality of the butter will be impaired by the formation of a bitter acid, which gives to the butter a disagreeable flavor. In winter the quantity of cream will be increased, and its quality improved, by bringing the milk to a temperature of about 120° before setting.

Cream.—As soon as the cream is taken from the milk it should be placed in stone jars or tin pails and set in a cool place. Sprinkle a small handful of fine salt over the top of the cream, and let it stand until churned. Should there be any milk at the bottom of the jar it should be separated from the cream, for the cheesy particles of the sour milk become mixed with the butter during the process of churning, and give it the white cheesy appearance which is sometimes observed when the butter “comes white.” The cheese decomposes upon exposure to the air, and renders the butter rancid. Such butter should never be packed with the good, for it will surely spoil the whole; “a little leaven will leaven the whole lump.”

Churning.—The proper temperature at which to churn cream is from 55° to 60°, and care should be taken that the cream be “washed down” so that all will granulate at the same time. When the butter “has come” to the size of

peas, draw or pour off the buttermilk, and pour into the churn a pail of cool water, and thoroughly "gather" by the aid of the "dasher" the butter into a compact mass; after which remove it to the butter-bowl. It should be again washed until the water is free from the least trace of milkiness, and then salted. Use the best Ashton salt, and if free from water one-half pound of salt is sufficient for 10 pounds of butter. Common salt should never be used, for it contains impurities which injure the butter. The cheapest salt in this case is certainly not the most economical. While the salt is being worked in, if too soft, let it stand in a cool place not over three or four hours, then work again and pack. While working, absorb all the moisture from the butter with a sponge covered by a linen cloth, previously moistened in cold water, and continue to work until all the brine is absorbed. No milky brine * should be allowed to remain in the butter, for it decomposes and injures it. During the process of working the temperature of the butter should not be higher than 55° or 58°. When it becomes warmer than this it loses its waxy, granular appearance, and becomes sticky and greasy. When the salt is not thoroughly worked in, the butter will have a streaked or marbled appearance.

Packing.—Place no undissolved salt in the bottom of the

* We have known those who would not work the brine out of the butter "because," say they, "it will weigh less;" mistaken shrewdness, to gain a penny they lose a pound. That it is necessary to leave brine in the butter to "keep it" is a great mistake.

tub or pail, unless covered with a cloth so the butter cannot come in contact with it. If this caution is not observed when sold, four or five pounds of butter is thus rendered comparatively worthless. Never pack a poor "churning" with the good butter, thinking it will not be found out. The sale of many a good firkin of butter is spoiled by a few pounds of poor butter becoming rancid in the centre or bottom, which taints the whole package. If there is any butter that is even suspicious, put it by itself.

Select neat pails, tubs, or firkins made of white oak, and cleanse them by placing in each about a pound of the common bicarbonate of soda, and then filling with boiling water, letting the water remain for twenty-four hours. Great care should be used in cleansing pails that are to be re-filled,* as they are usually bedaubed to a greater or less extent with rancid butter. A neglect of this precaution will often cause great loss. Butter until the first of June should be packed in pails or tubs and shipped as soon as made. This butter will keep sweet only a short time. As soon as the weather becomes too warm to ship without risk, pack in firkins, being careful to exclude the air as far as possible while packing. When the firkin is filled to within an inch of the top, dissolve two tablespoonfuls of white coffee sugar, and a piece of saltpetre about the size of a common bean, in sufficient strong brine to cover the butter and

* Pails or tubs after being once used, if properly cleansed, are preferable to new ones.

exclude the air. Place it in a cool dry cellar, and do not disturb it until ready to be shipped. In the fall the butter should be packed in pails or tubs and sold as fresh butter. An air-tight butter pail or tub is very desirable for shipping spring and fall butter.

Test of good butter.—Good butter should have a granular, waxy consistency, and a rich yellow color, except in the winter and spring, when the color is of a pale yellow or nearly white. When cut it should not soil the polished blade of the knife, and the cut surfaces should be free from a dewy appearance. The taste and smell should be entirely free from the slightest trace of rancidity, for if not, however good otherwise, when exposed to the air for a few days it will become almost worthless. The flavor of butter is various, generally depending upon the season, the water, the food of the cows, &c. The preference is merely a matter of choice. If butter upon being cut or repacked is covered with small drops of milky brine, it shows that it has not been sufficiently washed and worked, and although sweet it will not remain so if exposed to the air. When opened for use it should be immediately covered with a strong brine. When it is sticky or greasy, it shows that it was too warm while being churned and worked, or has been overheated since. Such butter is rancid, or will become so as soon as opened.

Setting-pan.—To insure a perfect separation of the cream from the milk a setting-pan has been successfully used in

England. It consists of a large tin pan about four inches deep, holding from four to six pails of milk. It may either set on a table or float in a reservoir of running spring water. Where running water is not to be had, the proper temperature may be obtained by the dripping of melting ice. At one end is a tube covered with a fine strainer to prevent the escape of the cream, through which the milk is to be drawn off, leaving the cream in the pan. All the cream may be secured by rinsing the pan in a little warm water.

The Cheese Dairy.—The superiority of factory cheese is entirely due to the great care exercised in its manufacture. But little cheese is now made by private dairies, for it can be better and more economically manufactured at the factory. With proper management it is more profitable for those who do not live near a cheese factory to make butter, unless they provide themselves with all the necessary apparatus.

Rich Cheese.—The richness of cheese varies in proportion to the amount of the butter that remains entangled in the curd. The following brief directions are from a practical cheesemaker:—

“When two milkings are united, strain the evening’s milk and cool by means of pieces of ice dropped into the pails before straining. In the morning take off all the cream, mix it with twice the quantity of new milk. Add warm water enough to raise it to the temperature of 98°. Rub annatto through a silk cloth sufficient to make the curd the color of rich cream. Into this put rennet sufficient to

curd in 35 minutes. Stir the whole into the milk previously raised to the temperature of 85°. The milk should be warmed by means of a pail of hot water set into it, but never by putting it over the fire, for the least burning of the milk will spoil the cheese. While the curd is setting, cover with a cloth to prevent the surface from cooling. The method of cutting, scalding, and pressing depends upon the varieties of cheese to be manufactured. About $\frac{1}{4}$ of a pound of the best Ashton salt is sufficient for 20 lbs. of curd. Care should be taken that the whey be entirely expressed."

The different varieties of cheese come to market under the names of Cheddar, Cheshire, and Gloucester. These are English cheese. The Dunlop cheese is from Scotland. The Dutch cheese is made in the north of Holland. The Parmesan cheese is made in Italy. Factory cheese is the best manufactured in this country, some of it being equal to the English. The private dairy cheese is of every grade and quality, from the richest Cheddar to that made of skim-milk.

Thermometer.—In the butter and cheese dairy the thermometer should be a constant companion. Those who trust to sensations are not aware how easily they may be deceived. Let a person put one hand in cold water, the other into warm, then both into another vessel, and it will *feel* warm to one hand and cold to the other. The only certain guide is the thermometer; its cost is but a trifle, it will save many dollars annually.

Ice-house.—Next in importance to the thermometer is the ice-house. Many farmers say “I can’t afford it.” They should say “I can’t afford to be without it.” It will save three times its cost every year. The method of building the following is so simple, and involves so trifling an expense that no man need have an excuse.

Select a place on the north side of some building; lay a floor twelve feet square on scantlings, one foot from the ground. Set firmly in the ground, near each corner, two posts, from four to six inches square, and about eight or ten feet long. When the weather becomes cold, place on the floor saw-dust, tan-bark, or rye-straw, to the depth of eight or ten inches. On the top, place another floor of the same size, putting a curb *inside* the posts to keep the filling between the floors in its place. Next make a curb ten feet square and six inches deep, and fasten the corners with common gate-hooks. On a cold day place the curb on the centre of the floor, put in two inches of tan-bark, and dash water over the bottom until it forms a coat of ice that will not leak. Fill the curb with water and let it stand until frozen solid. With boiling water thaw the curb loose, raise it to the top of the frozen mass, fill and freeze as before. Continue so doing until the mass is of the desired height. Place boards on the *inside* of the posts, and fill the space with tan-bark or rye-straw; nail boards on the *outside* of the posts and fill the space with rye-straw; cover the top with tan-bark to the depth of ten inches. Over the whole

put a roof, to shield from the sun and rain. Cut and take the ice from the top. Ice can be thus kept the entire season. If a stream of running water can be turned into the curb, the labor of filling will be much lessened.



SOILING CATTLE.

This is a rather unmeaning expression, and its origin is no more clear than is the fitness of its application; still it has come into such general use that it is now too late to change it.

It is applied to the feeding of cattle in yards or in stables, with grass or other green fodder, cut and hauled to them.

This practice is very rapidly growing in favor in all localities where land is very high priced, where manure is largely used, where the finer class of animals are kept, and where for any reason it is desired to keep a large stock on a small place. It is the best foundation of what is called *High Farming*.

It has been found by experiment that if a field bearing luxuriant grass or clover is divided into two equal parts, one half being used as pasture and the crop of the other being cut and fed in the stable as often as it grows to a sufficient height, this latter half will support, for the same time, four times as many animals of equal weight as will the depastured portion; and while the usual allowance of pasture land is at the rate of two acres for each cow, the allowance of land in soiling, where the system is practised in the best manner, is at the rate of only one-half of an acre for each cow.

Of course, this would not hold good on ordinary land which had been in no way prepared for the practice, but after one or two years' preparation by judicious use of the manure made by the animals fed, and by the aid of proper management, any fair land will support, on the system of soiling, four times as much stock as if they grazed upon it constantly and voided upon it all of their manure.

It was for a long time questioned, and very naturally too, whether cattle would remain in good health if they were deprived of the exercise which they necessarily take in getting their own food in the fields; but ample experience has proved that, if they are allowed good yards in which to exercise for a short time, once or twice a day, they keep in better condition and are less liable to disease than when they are exposed to the various changes of the weather in the fields.

It is also sometimes objected that this treatment is an unnatural or an artificial one. To this the reply is that our domestic animals are artificial productions. In nature we see no working oxen, and no cows give during the whole year a tenth part of the quantity of milk that cows have been forced to give in a state of domestication.

With the writer, the soiling of cattle is not a matter of theory. He has adopted the system on his own farm, and has sufficient evidence in his own practice of its substantial advantages.

Perhaps the most practical way to give an idea of the

manner in which stock is managed under the soiling system will be to describe the operations as there carried out.*

The farm† comprises sixty acres, lying in a nearly square body, and *all in one field*. Adjoining the main farm there is a small field in which to pasture calves during their first summer only, but it is not intended that the older animals shall ever feed except in their stalls.

In the centre of the farm there is an enclosure of about four acres, within which are concentrated all of the farm buildings; outside of this there is nothing to interfere with cultivation—no interior fences, rocks, nor trees.

The barn-yards occupy two acres of what was formerly an apple-orchard, and in the middle of this stands the barn (40 ft. \times 100 ft.). This has a cellar under the whole for the accumulation of manure, and (one corner of it) for the storage of roots. The main floor—the whole extent of the building—is occupied by two rows of stalls, the animals facing a central passage-way, through the entire length of which there runs a railway with a car, for distributing the food. The next floor above is used for the storage of hay and grain and of implements, and for the cutting and steaming of food in winter. Each floor and the cellar can be entered by loaded teams.

On the cattle floor there is a system of water-troughs which are constantly supplied from a tank on the floor

* To make this description more complete, a few improvements which are contemplated for the coming year are spoken of as though now in operation.

† Ogden Farm, Newport, R. I.

above, which is filled by a wind-mill, from a running spring. By this means water is always kept within reach every animal.

The floor is divided into four principal parts, separated from each other by bars which run (one on each side of the barn) from the rear of the stalls to the wall; and each of these divisions has its own door, communicating with a yard nearly half an acre in size, surrounded by a four-foot stone wall, and sufficiently shaded by the remains of the former orchard. Each set of animals has its own quarters and its own ample exercising ground, so that all danger from over-crowding is avoided.

They are turned out for exercise in pleasant weather at 8 A.M. and at 2 P.M., and are kept out (by closing the doors) for about two hours each time. If the doors are left open they return to their stalls almost immediately. Being abundantly fed, they show no disposition to move about, and I am satisfied that they give more milk and keep in better condition than if they were allowed the best pasture without shelter, even in the summer time.

Five times a day they are given as much green fodder as they will eat. This is cut in the field, loaded on to a cart, and hauled to the upper floor of the barn, where it is dumped through a trap-door into the car, by which it is carried to the stalls. The manure is dropped through an open slat-floor, and through scuttles, into the cellar, whence it is drawn in wagons directly to the field, having been well

worked over by hogs while in the cellar. Thus it will be seen that the labor of attending to a large stock of cattle is reduced to the lowest possible amount.

ARRANGEMENT OF CROPS FOR SOILING.

The amount of land that it is necessary to appropriate for the supply of fodder for each animal must, of course, depend on the quality of the land and on the degree to which its productiveness is forced.

Under all ordinary circumstances, one-half acre of land, in good heart and in good tilth, should be allowed for each full-grown milch cow of the ordinary breeds (more for short-horns), but, under high cultivation, this will allow a considerable amount of the produce to be cut for winter use.

The regular soiling crops are the following :—

Winter Rye,

Cabbages,

Oats,

Clover,

Grass, and

Indian corn.

Many other crops are available, such as Hungarian grass or millet, wheat, Jerusalem artichoke, sainfoin, &c., but the foregoing are the regular dependence of American farmers, and are the best for common use.

The best essay that has yet been written in this country

on the subject of "soiling" was prepared for the Massachusetts Agricultural Society by the Hon. Josiah Quincy, and was published in the Journal of that Society for 1820.

His recommendation is as follows :—

"1. As early in April as the state of the land will permit, which is usually between the 5th and the 10th, on properly prepared land, sow oats at the rate of four bushels to the acre.

"2. About the 20th of the same month, sow oats or barley, at the same rate per acre, in like quantity and proportions.

"3. Early in May, sow, in like manner, either of the above grains.

"4. Between the 10th and the 15th of May, sow Indian corn (the flat Southern being the best) in drills, three bushels to the acre, in like quantity and proportions.

"5. About the 25th of May sow corn in like quantity and proportions.

"6. About the 5th of June repeat the sowing of corn.

"7. After the last-mentioned sowing, barley should be sown in the above-mentioned quantity and proportions, in succession, on the 15th and 25th of June, and on the 1st of, or early in July; barley being the best qualified to resist the early frosts."

Mr. Quincy depended on the mowing of the best of his grass land to carry his stock through the month of June, or from the earliest pasturing season to the 1st of July,

when he expected his first sowing of oats to be ready for the scythe. After the first killing frost, he depended on the tops of about twelve acres of root crops, for the use of fifteen cows.

The plan which I have adopted is a modification of the above, and is as follows (for twelve cows) :—

1. Early in the autumn sow three acres of winter rye, to be cut from May 15th to June 15th.

2. Early in April, three acres oats, to be cut from June 15th to July 1st.

3. Late in April, two acres oats or barley, to be cut from July 1st to July 15th.

4. Early in May, two acres oats or barley, to be cut from July 15th to August 10th.

5. Middle of May, two acres corn, to be cut from August 10th to September 1st.

6. Middle of June, the three acres from which rye has been cut to be sown with corn, to be cut from September 1st until September 20th.

7. Early in July, the first three acres sown with oats to be resown with barley, to be cut from September 20th until the harvest of roots and cabbages furnishes a stock of green refuse, which will suffice until winter feeding commences.

This is an allowance of twelve acres for twelve cows, and assumes that the latter end of the season will be helped out by root tops, &c. The reason for appropriating so much land

is that the soil is not yet in sufficiently good condition to *insure* an ample supply from a much smaller area. In a season of extraordinary drought the whole of the product may be consumed, but in any ordinary year a very large part of it would be in excess, to be cured and stored for winter use, and to furnish a supply of dry food, with which occasionally to alternate with the fresh fodder, to prevent the too great relaxation of the bowels which a free use of succulent food sometimes causes.

In September three acres of the four comprising Nos. 4 and 5 should be sown with winter rye for the following spring's use, and the rotation should follow in regular order. If all of the manure made in the soiling season were to be used on these twelve acres year after year, I am satisfied that they might be made in time to support, during the whole of the usual pasturing season, thirty milch cows, or five cows for each two acres.

In my own case, as one of my reasons for adopting the system of soiling has been that it is the best help in bringing up a worn-out farm, I shall each year raise my forage on fresh land, so as to give the whole place the benefit of the treatment.

Of course, a rule which will apply in one region may not be the best for another, and each farmer must decide for himself the extent to which he can profitably adopt the system on his farm, and also what crops will best accomplish the desired end in his own case.

Where it is desirable to plough as little as possible, clover and grass may with advantage enter much more largely into the arrangement.

Two general principles, however, may be stated as applicable to all of the more temperate regions of our Northern States—

1. The *earliest* abundant food will be secured by the use of winter rye.
2. The *best* and most abundant food for the later summer and earlier autumn time will be secured by the use of Indian corn.

ARGUMENTS IN FAVOR OF SOILING.

Mr. Quincy states the following as the leading advantages of this system:—

- “ 1st. The saving of land.
- “ 2d. The saving of fencing.
- “ 3d. The economizing of food.
- “ 4th. The better condition and greater comfort of the cattle.
- “ 5th. The greater product of milk.
- “ 6th. The attainment of manure.”

On the subject of the 3d item—the economy of food—he says: “There are six ways by which beasts destroy the article destined for their food—1. By eating; 2. By walking; 3. By dunging; 4. By staling; 5. By lying down; 6. By

breathing on it. Of these six, the first only is useful. All the others are wasteful."

The other points he elucidates with equal force, but at too great length for full quotation here.

The statement made above that a milch cow may be kept during the ordinary pasturing season upon the produce of one-half acre of land, while of land of the same character at least two acres would be necessary on the pasturage system, is sufficient to illustrate the saving of land. Yet this statement, which will be supported by the testimony of all who practise the system on land of good quality, is far below the estimate of many who have had a lifelong experience of soiling in Europe. Some of them place the proportion in favor of soiling as high as 1 to 7. Of course the amount of stock which may be fed from the produce of a single acre depends very much on the manner in which that acre is cultivated, and the question of the cost of labor must determine whether it is or is not profitable to force the production beyond a given extent.

As to fencing, it is only necessary to remind nearly every farmer of his own experience of the first cost of building, and of the yearly cost of repairing the fences of his own farm, and to say that by the soiling system, when completely carried out, *all interior fences may and should be entirely dispensed with.*

Add to the question of expense, the fact that useless headlands and their nurseries of noxious weeds are got rid of,

and that the plough can be driven, if desired, straight through from one side of the farm to the other, and the argument needs no re-enforcement.

Concerning the condition of the cattle, the following is stated by Quincy:—"One writer asserts that he has kept a large herd for several years in this way, and during the whole time 'he never had an animal essentially sick, had never one die, and had never one miscarry.'" The general result of the experience of hundreds of farmers in Europe, and of considerable experience in America, is, that cattle are really better off in every way, under the protection of the soiling barn, with its ample and regularly supplied food, and with the advantage of daily currying and exercise, than when left to shift for themselves exposed to the vicissitudes of the weather.


The quantity of milk may never be so large as it is during the flush weeks of June, when the cows are gorging their maiden appetites on rich grass; but the consumption of food from the first of May to the first of November (and *consequently* the yield of milk) will be much greater.

"Last, but by no means the least," the question of manure asserts its claim to the fullest consideration. Were it not for this item of the calculation the arguments in favor of soiling would lose more than half their force.

The immense superiority, both in quality and evenness of distribution over the soil, of manure which is made and kept under cover, over that which is dropped at random on pas-

ture fields; and the advantage of being able to apply it when we please, where we please, and in such quantities as we please, are too well known to all who have to use manure to produce paying crops, for any argument on the subject to be necessary. There is no way in which so much manure of such excellent quality can be landed on the farm without a far greater outlay of money than is necessary to pay for all the labor required for ploughing, sowing, "tending," cutting, and hauling the food, and for currying and feeding the animals under the most complete soiling management.

Of course the manure argument does not hold (nor is the system of soiling to be recommended) for those districts of the West where the laughing harvest follows the tickling hoe; where straw is burned in the fields, and barns are moved to get away from the accumulated manure. But for the older settled countries of the East and South (and for the future West—the West with its "inexhaustible fertility" exhausted) it does hold, and with such force that as population grows more dense—and farmers more wise—it alone, even if there were no other advantage in the system, must in time compel the rapid increase of the practice of soiling.



APPARATUS FOR STEAMING FOOD, &c.

STEAMING FOOD FOR STOCK.

A more recent improvement than "soiling" in the keeping of cattle, on farms where it is important to make every pound of food tell with the fullest effect in the production of meat, muscle, or milk (and on what farm is this not important?), is the *steaming of food* in winter.

Although this practice has been the subject of much less experiment than soiling, and is, consequently, less generally recognized as worthy of adoption, enough is known of its advantages, both by experience and from theory, to make its brief discussion necessary to the completeness of this book.

During the past year I have investigated the subject with some thoroughness, and have determined to adopt it on my own farm; and I can hardly do better than to give here some account of my investigations, in order that my readers may decide for themselves the soundness of my reasons for the determination.

My serious attention was first called to the matter by an article in the Report of the Department of Agriculture for 1865, written by Mr. E. W. Stewart of North Evans, N. Y. He therein details his own experience of ten years in steaming food for a large stock of cattle and horses, gives a succinct statement of the reasons why steaming is beneficial,

and sustains his own opinion by the concurrent testimony of other practical farmers who have found the practice beneficial.

The following are the results of the operation as stated by Mr. Stewart:—

“1. It renders mouldy hay, straw, and corn-stalks perfectly sweet and palatable. Animals seem to relish straw taken from a stack which has been wet and badly damaged for ordinary use; and even in any condition, except ‘dry rot,’ steaming will restore its sweetness. When keeping a large stock, we have often purchased stacks of straw which would have been worthless for feeding in the ordinary way, and have been able to detect no difference, after steaming, in the smell or the relish with which it was eaten.

“2. It diffuses the odor of the bran, corn-meal, oil-meal, carrots, or whatever is mixed with the feed, through the whole mass; and thus it may cheaply be flavored to suit the animal.

“3. It softens the tough fibre of the dry corn-stalk, rye-straw, and other hard material, rendering it almost like green succulent food, and easily masticated and digested by the animal.

“4. It renders beans and peas agreeable food to horses, as well as other animals, and thus enables the feeder to combine more nitrogenous food in the diet of his animals.

“5. It enables the feeder to turn everything raised into

food for his stock, without lessening the value of his manure. Indeed, the manure made from steamed food decomposes more readily, and is therefore more valuable than when used in a fresh state. Manure made from steamed food is always ready for use, and is regarded by those who have used it as much more valuable, for the same bulk, than that made from uncooked food.

“6. We have found it to cure incipient heaves in horses ; and horses having a cough for several months at pasture, have been cured in two weeks on steamed food. It has a remarkable effect on horses with a sudden cold and in constipation. Horses fed upon it seem much less liable to disease ; in fact, in this respect, it seems to have all the good qualities of grass, the natural food of animals.

“7. It produces a marked difference in the appearance of the animal, at once causing the coat to become smooth and of brighter color—regulates the digestion, makes the animal more contented and satisfied, enables fattening stock to eat their food with less labor (and consequently requires less to keep up the animal heat), gives working animals time to eat all that is necessary for them in the intervals of labor ; and this is of much importance, especially with horses. It also enables the feeder to fatten animals in one-third less time.

“8. It saves at least one-third of the food. We have found two bushels of cut and cooked hay to satisfy cows as well as three bushels of uncooked hay, and the manure in the case of the uncooked hay contained much more fibrous

matter unutilized by the animal. This is more particularly the case with horses."

Other publications on the subject fully confirm Mr. Stewart's estimate, and we commend his essay, which is accessible to all, to the careful attention of every feeder of farm stock.

In January (1868) I visited the farm of Messrs. S. & D. Wells, at Wethersfield, Conn., for the purpose of examining their cow stable and its fixtures.

The leading features of this establishment are a constant water-supply, and apparatus for cutting and steaming food.* The latter was introduced at a cost of about \$500. It comprises a three-horse steam-engine of very simple construction, a tubular boiler of sufficient capacity to run the engine, a strong *power* stalk-cutter, and a chest for steaming food.

There were about thirty cows in the stable. They receive steamed food morning and night, and dry hay at noon. The steamed food consists of hay of poor quality, straw, or corn-stalks, cut to short lengths, sprinkled until thoroughly wet, and then dusted with bran or meal, and steamed for about two hours.

The engine has power enough to cut in a couple of hours

* The water is brought from a living spring and flows through galvanized iron pipes which form the connections between the bottoms of small iron troughs standing at the head of the partitions which divide each pair of stalls. The last trough overflows through a pipe near its top, and the water wells up to the level of this overflow in each trough of the series. By this simple arrangement, a constantly changing supply of water is kept always in front of the cattle.

a supply sufficient for the whole week, and enough is steamed at one charge to last for three or four days. Steam is made only twice in each week (once for cutting and steaming, and once for steaming only), and then only for a short time.

The steaming box is about four feet square and eight feet high. The materials are put into the box from the floor above that of which the cow stable is an extension, and are removed through a door in one of its sides on the feeding floor. Elevated a short distance above the bottom, there is a false bottom perforated with many holes. The steam is let in below this, and is thus allowed to rise evenly through the the whole mass.

The box is made of two thicknesses of one-inch, matched spruce boards (one set running up and down, and the other across). The doors are not made with any very great care to prevent the escape of steam, nor does it seem to be considered necessary to do more than to have the box strong enough to hold its burden of wet fodder.

The Messrs. Wells find that Mr. Stewart's opinion—given above—is, in all essential particulars, sustained by the results of their experience. They think that steaming adds one-half to the feeding value of fodder.

It was what I saw on their farm, more than anything else, which caused me to decide on adopting the system in my own practice. My apparatus is not yet completed, and I cannot, therefore, speak on the subject with the authority of a successful experimenter; but from all that I can learn, I

am satisfied that the advantages of steaming have hardly been overrated.

The theory of the process (in a nutshell) is this: Cattle and horses in a state of nature live the year round on succulent green herbage. When the cold weather begins to cut short the supply in the more northern latitudes, they migrate toward the south. Man steps in and keeps them in the colder climate. He substitutes dried grass for fresh grass. Steaming will, in a great measure, restore hay to the condition of green grass. Also, many constituents of hay, straw, &c., are insoluble and indigestible. By the action of heat and moisture they become soluble, or at least are reduced to a condition in which they are easily available to the digestive organs of animals. Starch-grains, according to the best authorities, are coated with a layer or cuticle which resists—to a great extent—the action of the juices of the stomach, while its interior parts, could they be directly exposed, would readily be assimilated; therefore, as heat causes the interior of the grains to swell and burst their coating, exposing themselves on the surface, as the interior parts of a kernel of corn do in “popping,” the process of steaming (or any cooking) makes the starchy part of food more readily available.

Examinations of the droppings of animals fed on cooked and uncooked food furnish results which confirm the foregoing opinion.

Carefully conducted experiments on animals of equal

weight, and of like condition in all respects, invariably show that those which are fed on cooked food take on fat, and form bone and muscle more rapidly than those which get only raw food. If, after a certain time, the food is changed—the cooked being given to the animal that has been receiving the uncooked, and *vice versa*—the rapidity of growth will change too. The trial has often been made, and the result has been invariably the same.

In fact, in all of the essays and opinions on the subject of cooking food for domestic animals, in this country and in Europe, I have failed to find the first one that is not decidedly favorable.

Steaming, of course, is valuable only because it is a means of *cooking*, and the arguments in its favor bear equally on the subject of *boiling*. *Steaming* is rapidly coming into use because of its greater convenience and economy.

How to make a Steaming Apparatus.—Any device by which steam may be generated under a very slight pressure—barely sufficient to cause it to penetrate the mass to be cooked—and conducted to the vessel in which the steaming is to be done, will accomplish the desired purpose; but, of course, the more convenient the arrangement, and the less the waste of steam (whether by condensation or otherwise), the more economically the process may be performed, as to both time and fuel.

Mr. Stewart suggests a plan which, from its cheapness, will answer a good purpose where the stock to be cooked

for is small, or where it is desired to experiment on a small scale.

It is a box made of well jointed 2-inch pine, seven or eight feet long, and about two and a half feet wide, with a bottom of No. 16 sheet iron, nailed securely on to the lower edge of the sides and ends, and turned up a little outside of them—say half an inch. This box has a false bottom, of wood or iron, placed about three inches above the fast bottom, and perforated with many small holes, and a closely-fitting cover over the top.

It stands on brick walls which do not come quite so far out as the wooden sides of the box. At one end of the chamber enclosed by these walls there is a wood fire-place, and from the other end a chimney rises.

The space between the bottom and the false bottom is partly filled with water, cut hay mixed with meal or bran is put in the box above the false bottom, the cover is closed, and the fire is started. The steam rises through the perforations in the false bottom, and cooks the mass above it.

A much more complete apparatus for steaming, and in large practice a more economical one, comprises a boiler for generating the steam, a box in which to place the food, and a wooden, or well protected steam-pipe to connect the two. The box should have a perforated false bottom, and the steam should be introduced beneath this, so that it may diffuse itself uniformly through the mass.

The boiler may, of course, be of any pattern that will secure the economical generation of steam. A discarded engine-boiler will answer every purpose if it is strong enough to bear a pressure of, say, five or ten pounds to the inch—a slight pressure being necessary to force the steam through the mass of hay.

D. R. Prindle's Agricultural Boiler, which is shown in the accompanying cut, is admirably adapted for this use.



FIG. 1.




FIG. 2.

Prindle's Agricultural Steamer and Cauldron (shown in Figs. 1 and 2) is the invention of Mr. D. R. Prindle, of East Bethany, New York, and is largely manufactured by Messrs. Savery & Co. of Philadelphia.

Its popularity seems to be rapidly increasing, and there is no question that it is the best steaming apparatus for the use

of all farmers who do not employ steam-engines that has yet been invented.

It consists of a cauldron set over a furnace arranged to burn either wood or coal, and furnished with a dome which fits closely over it and is keyed down so as to make a steam-joint. It is provided with a test-cock to show when it needs the addition of water, a safety-valve which is also a *vacuum* valve, a funnel for filling, and one or more pipes to convey the steam to the cooking-boxes.

Aside from its use in steaming fodder for cattle, it may be used to heat water to scald hogs, or for other purposes, to warm buildings, to cook roots or meal for hogs or grain for fowls, and for a variety of other purposes for which hot air, hot water, or steam are useful.

For farm use, especially when constant steam is not required, Prindle's steamer is much better than an engine-boiler, as it works only at a very low pressure, and is consequently quite safe, and is much cheaper when we consider the cost of setting up the larger engine-boiler, and its more expensive transportation.

Full particulars concerning the Prindle steamer may be obtained by application to the inventor.

I have not determined, in my own case, what power to adopt for the cutting of my long fodder. The question is about evenly balanced between a small steam-engine, a wind-mill, and a railway horse-power, for final use; but as the first cost will be less, I shall commence with the horse-power

belonging to a threshing machine, and a Prindle boiler, changing to one, the engine or mill, at a future day, if it seems desirable.

It is hardly prudent to make any positive calculations in advance of actual experiment, but I anticipate—and I base my calculations on a very careful survey of the whole field—a saving of about *forty per cent.* in the cost of feeding my stock, over the present system of feeding only the best hay uncut. A part of the saving will be due to the more digestible condition of the food, and a part to the fact that a much cheaper quality of hay, or straw, or corn-stalks can be largely used. A saving of very much less than this, when from thirty to forty head are to be provided for, will be enough to make a fair profit on the business.

The various uses for which steam can be adapted seems to be but little understood by the masses. Fear of explosions, scalding, &c., as well as want of knowledge of its great advantages, has thus far prevented its general introduction.

The want of a perfectly safe and easily managed low pressure apparatus with which to accomplish all the requirements of domestic use, has also been a great drawback. The great advantages of cooking, heating, boiling, &c., by steam, are obvious when it is remembered that it can be done with much less water and fuel, requiring but little care of the operator, and using wooden vessels (if desired) of any kind, size, or shape (a great desideratum). By its use there

is no re-filling of *kettles* (the ordinary mode) to get a desired quantity; no constant watching or stirring, or removal of the substance while hot, to prevent burning; no cleaning of kettles for every separate job, which can be done by steam. By the use of this powerful agent, large quantities may be boiled or steamed, or several vessels (if need be) treated at the same time; and when desirable, the steam can be conveyed in pipes or logs to some little distance, using proper care in protecting the same from condensation; thus avoiding, many times, danger from fire, and accommodating itself to all the various purposes of domestic economy, as well as in the manufacturing of many articles or compounds, when danger from burning or explosion is so common. By steam the clothes may be boiled at any point in the barrel or tub; the bath-tub may be warmed in an adjoining room; the farm or stock-feeder could easily cook in quantities at a time, or scald his hogs, steam his barrels, &c., &c. We believe that when a cheap, simple, and perfectly safe apparatus is once introduced, that the subject (as it deserves) will receive much more attention, as by steam *all classes* might as easily be benefited.

ADVANTAGES OF COOKED FOOD.


The *American Agriculturist* for January, 1860, says: "Experiments made by C. M. Clay, of Kentucky, showed that one bushel of dry corn made 5 lbs. 10 oz. of pork; of boiled corn, 14 lbs. 7 oz., and boiled meal, 16 to 18 lbs."

Morton's Cyclopædia of Agriculture (than which there is no higher authority in Europe) says: "As to steaming food for cattle, there is abundant experience to recommend it. The process of cooking renders soluble that which would otherwise be imperfectly digested. It removes, in some cases, what would otherwise be unwholesome; and it renders savory what would otherwise be distasteful."

Loudon's Encyclopædia of Agriculture remarks: "Unless food be thoroughly deprived of its vegetative powers before it enters the stomach, the whole nourishment which it is capable of affording cannot be derived from it. The most effectual mode of destroying the living principle is by the *application of heat*, by steaming or boiling."

The *Society of Shakers*, at Lebanon, N. Y., famous for pork-raising, say: "For fattening animals, swine particularly, we consider three of cooked equal to four of raw meal."

GARDENING FOR MARKET.



While market-gardening, as a systematic business, is quite distinct from farming, there is no farmer who lives near a town who may not make the raising of certain crops on a small scale very profitable. Success in this branch of the business of the farmer requires that the land to be devoted to its prosecution be dry, warmly situated, with a good exposure, and *rich* and again *rich*.

The amount of manure which may be *profitably* applied to land intended for the growth of market vegetables has hardly any limit. One hundred cartloads of good horse

manure to an acre, *every year*, will pay more *profit* than will fifty loads; and I am inclined to believe that even two hundred loads would pay better still.

The cultivation of vegetables entails, in any case, a heavy outlay for labor, seed, expenses of marketing, &c., and these are about the same (except in the matter of marketing) for a light as for a heavy crop—it takes a certain amount of produce to pay the cost, and up to this point there is *no profit*. Beyond this point, except the cost of the manure, it is nearly all profit, and the more we can stimulate excessive production the more rapidly will the ratio of profits increase over the expenses.

No farmer can hope to become really successful in raising vegetables for market until he is prepared to expend—including the value of the manure used—at least \$300 annually on every acre of his garden land. With this outlay, if his soil is good and well placed, and his market is a good one, and *if he is the right man for the business*, he ought to make a clear profit of \$500 per acre.

The character of the market should be well understood. If there is a manufacturing town near by, or any town having a population which includes a large proportion of laboring people, the case is a simple one.

It should be well understood that it does not pay (at least so far as gardening is concerned) to feed *the rich*. They are like the black sheep of the flock, that don't eat so much as the white ones—there are not so many of them, and, as

another reason, they do not eat so largely of coarse vegetables. A hearty Irish laborer, with a stout hardworking wife and a table full of healthy children, will use up cabbages and turnips in a way to delight the heart of a gardener; and the atmosphere of a manufacturing town will evaporate a farmer's load of these vegetables as the sun dries up the morning mists.

To any one who is disposed to venture an acre or two in gardening, no better service can be done than to recommend him to read Peter Henderson's "Gardening for Profit," wherein are laid down precise rules for the management of every department of the business.

We have here only space to give a few practical hints which will be chiefly of use to *farmers* who propose to devote a portion of their time to the simpler kind of gardening.

It may be given as a general rule, that the only crops that it will pay the *farmer* to raise, in his market garden, are beets, cabbages (early and late), sweet corn, cucumbers, onions (rare-ripes), parsnips, radishes, spinach, and tomatoes.

The size, arrangement, and equipment of the garden.—We will suppose a farmer to be about to embark in this business, and that he is willing to invest in it a capital of one thousand dollars. Of course the same general rules will apply for a more or less extensive operation. He should select two acres of light dry land (if he has it, and if not he should thoroughly underdrain it), if possible with an exposure to

the east or south. If it is sheltered from the north and west by an orchard or by other trees, so much the better.

The land may be more economically arranged if it lies in about a square body, and should be fenced on the north and west sides with a tight board fence six or eight feet high. A fence of the latter height, made in the best manner, of pine boards, capped with a spruce rail, will cost in the vicinity of New York about \$200 for 600 running feet. This fence should set close to the ground, so that the wind cannot draw under it, and it will have the effect of very materially modifying the climate, and enabling the growing of much earlier vegetables.

Close in the northwest corner he should then set up two parallel rows of hemlock boards, nailed to 2x3 stakes, driven into the ground. The back line of boarding should be 12 inches high, parallel to the fence and three feet distant from it. The other row should be 8 inches high, parallel to and 6 feet and 2 inches distant from the first, outside measurement. Both to be 187 feet long, with boards to close up the ends, and the ground enclosed by them should be spaded and manured. This is the "cold frame," which is to be covered by 50 sashes, each 3 feet 9 inches wide by 6 feet $2\frac{1}{2}$ inches long, having four rows of glass, each containing nine 8x10 lights set lengthwise across the space—the rails being ten inches apart. The sashes to be made of $1\frac{3}{4}$ inch stuff and strengthened by a flat rod of iron (1 inch by $\frac{3}{16}$ inch) let in flush on the under side and screwed fast to the bars

and rails, across the middle of the sash. It is best to make the sashes in the best manner, as they are a very important part of the permanent stock in trade of the garden. They will cost, at an outside price, \$250.

The ground of the garden should be deeply ploughed and subsoiled in July or August, and if the weeds that grow upon it are likely to ripen their seeds, they should be mowed down late in the fall. Before winter sets in, the largest amount of horse manure that can be bought for \$200, delivered, should be spread upon the surface, and left exposed to the rain and melting snow of the winter.

About the middle of September, sow in a well-prepared seed-bed in an old garden, twelve ounces of the seed of Jersey Wakefield cabbage, and four ounces of Fottler's Improved Brunswick. At about the same time sow on three feet of one end of the cold frame, one ounce of black-seeded butter lettuce, and one ounce of early-curved Simpson lettuce, giving to each about nine square feet. These are to remain where they are sown during the winter. The cabbage plants will be large enough to transplant about six weeks from the time of sowing, when they are to be "pricked out" in the cold frame two inches apart each way, which will give about 800 plants to a sash. These plants should be well watered, and sprinkled with a light coating of air-slaked lime.

They will need to be protected by the glass until they are firmly rooted (the sashes being tilted up at the back to give them air whenever the sun is on them), and on frosty nights,

and they should be gradually accustomed to the cold air, so that they may be able to withstand the hard freezing that they will get in the winter; all through the winter they should have air whenever the frost is thawed from the under side of the glass, and on fine days the sashes should be stripped off from them altogether. The end where the lettuce plants are standing should have less air, and should have the protection at night of an old carpet thrown over the sash. Directly in front of the cold frame there should be a second frame made of exactly the same size and character. This should be filled with straw, leaves, or other rubbish which will keep it from freezing, and about the last of February or the first of March its covering should be removed and about three inches of well-rotted manure should be dug into it—not too deeply. The lettuce plants are now to be transplanted to this frame, at distances of six and a-half or seven inches each way (about seventy plants to a sash), and covered by the sashes which may now be taken entirely from the hardened cabbage plants. If light board shutters have been provided to cover the cabbages during severe storms, it will be better, but they will stand any amount of hardship after their winter's training. The lettuce plants should have plenty of air during fine weather (and some air whenever it is not freezing), should be abundantly watered if the season is dry, and should be forced by as much heat as can be given them without depriving them of air. They will be ready for market about the middle of May, when lettuce

usually sells in towns (not in the larger cities) for from 8c. to 12c. per head.

During the latter part of April, plant sixty three-inch pots with half a dozen seeds each of White Spine cucumber, and set them in a warm light room in the house. By the time the lettuce is sold off these will be sturdy plants, and they should be thinned to three in each pot. Now dig holes a foot deep, and a foot in diameter, at intervals of three feet in the lettuce frame, and fill them with very thoroughly rotted and rich compost, covering it with a little soil. On each of these plant the contents of a pot, without disturbing the roots of the plants, and cover closely with the sashes. Give a little air in the middle of the day, but cover close from 4 P.M. until 10 A.M., and during all chilly weather; water copiously, and uncover to all warm rains.

By the latter part of June the picking will commence (at from 5c. to 30c. each), and it may be continued as long as the price is not less than 1c. each. This crop is more uncertain and varying in its results than lettuce, but it usually pays well, and is very inexpensive.

Now let us sum up the probable income of 50 sashes, managed as directed above:—

35,000 cabbage plants, at \$10	\$350
3,500 lettuces, at 8c.	280
Cucumbers (from \$25 to \$100), say.....	50
	<hr/>
	\$680

This is earned with a small investment, and the labor is

mainly done in the fall and winter, when other work is slack; and it has the great advantage of coming in early, when there is a demand for ready money to pay for labor, &c.

Five hundred tomato plants may be started in the kitchen window, or in a small hot-bed, and by the middle of April they may be pricked out in one end of the lettuce frame. As early in May as the danger of frosts has passed, they should be set out at intervals of fifteen inches along the foot of the fence on the north and west sides of the field, to be trained up against it (tacked fast), and kept trimmed to single stems. At a height of six feet they should be pinched off and their growth kept close. They should be planted in a very rich soil, and well watered. They can hardly fail to produce early crops, and ought to sell for \$75 to \$100.

Now we come to the management of the field crops.

If we could only raise cabbages year after year on the same land, our business would be a very simple one. We might take two crops yearly (an early and a late one) of the most profitable and easily raised vegetable on our list.

But, unfortunately, one crop in two years is all we can reasonably hope for, as the "club-foot" will surely attack an immediately succeeding crop on the same ground, and our best plan is to arrange to grow as many cabbages as we safely can—making this point our constant aim—and to occupy the land as profitably as possible the rest of the time.

Therefore, the field should be divided into two equal parts,

one side being prepared for cabbages and the other for such other crops as will not interfere with the growth of cabbages the next year.

The first operation is the preparation of the ground for early cabbages, for which we devote a space of about one acre.

The manure which was spread in the fall should be lightly ploughed in—not deep enough to turn up the old sod—and a thousand pounds of Peruvian guano, two thousand pounds of fish guano, or fifteen hundred pounds of bone-dust, should be evenly sown over the ground, and thoroughly harrowed in. Either of these manures will cost about \$40. As early as it is possible to get the ground into proper condition, as described above, the cabbage plants in the cold frame should be set out, in rows two feet apart, and about 16 inches apart in the rows. It will probably be best to plant three-fourths of the piece with the Jersey Wakefield, and the remainder with the Brunswick, which will begin to be fit for market at about the time when the Wakefield is all sold.

This amount of land will receive about 15,000 plants, leaving about 20,000 plants to be sold from the frame. If the value of cold frame plants is understood in the vicinity, they will be readily taken up at \$10 per thousand.

If there is a good summer market for lettuce, the Early Curled Simpson may be set out between the rows of cabbage, when it will grow to a marketable size before the whole

ground will be required by the main crop. In the neighborhood of small towns this will not be worth while, as there is but little demand for lettuce after June 1st.

As soon as the cabbages are planted—and this may be done even so early as in March, if the weather is fine—the other half of the garden should be manured and prepared in the same manner, and planted with beets, onions, parsnips, spinach, and radishes; the first four in about equal proportions, and in the following manner:—

Beets (of the Bassano and the early turnip-rooted blood variety) should be very thickly planted in rows 18 inches apart—thickly, because the early frosts may cut off a part of the crop—and when they are fairly up, they should be singled out to intervals of about 4 inches in the rows.

The *onions* should be “sets” raised the previous year. These may usually be bought for from \$6 to \$10 per bushel, according to size—the smallest bearing the highest price. They should be set in rows 9 inches apart, and at intervals of 3 inches in the rows, being firmly pressed down in the bottom of the line made by the marker. Every seventh row should be omitted to leave room to walk among the crop, and the sets should be entirely covered by raking the beds evenly over.

Onions raised from the seed are rather a farm than a garden crop, and will not pay to raise on land so expensively manured as that under consideration.

Onions raised from “sets” are called *Rare Ripes*, and

they always meet a ready sale in any market where there is a market for any vegetables. Still, as it is considerable work to tie them, it will be best not to raise more than one-quarter of an acre of them.

Parsnips should be planted early in May on well prepared (deeply loosened) ground, in rows 27 inches apart, the seed being strewn thickly in the rows, and the plants finally thinned to intervals of six inches. The reason for putting the rows so wide asunder is that it enables us to cultivate the crop with the horse-hoe at a time when labor can be ill spared for hand-hoeing.

Spinach.—This crop, the first year, must be planted in the spring; by planting very early, on ground so heavily manured, it will be in market ahead of green peas, and will bring a good price, but after these are plenty it can hardly be sold at any price. The cultivation of this crop is extremely simple. The seeds are sown pretty thickly (say 10 lbs. per acre) in rows about 12 or 14 inches apart, and the land kept clean until it is large enough to cut.

For all subsequent years, spinach should be planted about September 15th, on the ground from which the Brunswick cabbage has been taken, this being first well manured with animal manure. It will require (above the latitude of New York) a light covering of seaweed, leaves, or straw during winter. Coming very early into market, it often brings *four dollars a barrel*.

Radishes are a stolen crop, and, to a limited extent, they

may be very profitably grown. It is best to raise both the long scarlet and the short top turnip-rooted varieties—the former for common trade, and the latter for those who are more choice in their taste, the proportion of each being regulated according to the character of the market.

The seed may be sown, rather thinly, with a seed drill between the rows of beets. No cultivation is needed. The seed is the only cost except the preparation for market, and this need be applied only to so much as there is a sale for; the rest can be simply cut out with a push hoe, before the beets will require the whole ground.

We have now provided for the planting of all the land, and will need to commence promptly to use the hoes, of which at least two should be kept going incessantly until the crops are all firmly established, and are able to hold their own against weeds. In fact, at no time during the growth of the crops, until they are too large to be worked among without injury, should weeds be allowed to grow *at all*. If they once get started so that there must be a fight to get rid of them, we may as well say good-bye to all hope of profit, for they will require more labor than it will be pleasant to pay for, and the crops will be materially injured by them. If, on the other hand, every foot of the land be lightly hoed over (or even raked with a light iron rake until it becomes too hard) once a week, there will be no weeds to kill, and the plants themselves will be sufficiently benefited by the operation to pay the cost.

Harvesting the crops, and preparing them for market.

The first sales will be of *radishes* and *spinach*. Long radishes are pulled, and tied in bunches, and then thrown into water. In a few minutes they are taken out by the tops, laid against a board which stands sloping into the water, and there washed clean with a wisp-broom.

The round radishes grow at the top of the ground, and so little dirt adheres to them that they only require to be soaked for a few minutes and then shaken in the water.

Spinach is simply cut off at the top of the root and packed (dry) in barrels—40 lbs. being a barrel. It is the easiest of all the crops, except cabbages, to prepare for market.

Parsnips are, as every farmer knows, either left in the ground until spring, or taken up in the fall and stored like any other roots.

Beets are pulled when about half grown; the outside leaves torn off so as to leave only enough to hold them by securely, the roots washed clean, and tied in bunches of four or five, according to the varying custom of different markets.

Onions (rare-ripes) are pulled when the bulb has a diameter of three-quarters of an inch or thereabouts—the larger the better—and, after the removal of the dead skin, are tied in bunches of five or ten. For the New York market, they must be washed. For Eastern markets this is not necessary. It is quite an addition to the cost of preparation.

Cabbages (the early sorts) are simply cut off near the

ground, with nearly all their leaves, and, if they are to be shipped, are packed in barrels or crockery-crates. They will stand a good deal of rough treatment.

Prices of Early Vegetables.—On this subject but little can be said that will be a criterion for different localities, except that in nearly all of the smaller towns they sell for from 50 to 100 per cent. above the New York quotations. The cause of this anomalous condition is that these towns are nearly always supplied with *early* vegetables from the larger cities.

Probably the following may be taken as a fair average of prices in towns of from 10,000 to 50,000 inhabitants, during a series of years:—

Cabbages, 8 cents each.

Onions (rare-ripes), 50 cents per dozen bunches of five each.

Beets, 75 cents per dozen bunches of five each.

Radishes, 30 cents per dozen bunches of about ten each.

Spinach, \$1.50 per barrel.

Second Crops.—We have now cleared all of the land except that which is occupied by the parsnips. This produces but one crop during the season, and we have not very much more to expect from the use of the land. Our profit must have come mainly from the early crops. Still, enough may be expected to make a fair return for the labor of cultivation, and for the use of the land and manure, and the land needs to be cultivated for its own sake. The gardeners about the

large cities, having a market for everything green that they can raise during the whole year, and for some crops, such as celery and salsify, which meet with no sale in small places, find their second crops very profitable; but, in our case, the chances are that we must be content with small returns from this source.

We are debarred from raising rutabagas, or French turnips, and late cabbages, for the reason that these cannot follow our crop of cabbages, and if they were made to follow any of the other crops they would injure the land for the growth of early cabbages the next year.

Celery is a good crop for land that is in good condition, but it is hardly worth raising for small markets.

Horseradish, sweet herbs, mangel wurzel, sweet corn, and common turnips are about the only safe reliance. Of these, the first is the most profitable, as it finds a ready sale among the pickle-makers in cities. Concerning its cultivation, the following is copied from an article furnished by Peter Henderson for the Report of the Agricultural Department for 1865:—

“The culture is very simple, and so far very profitable. The plants or *sets* used are the pieces broken off from the main root in its preparation for market. These are cut into lengths of about six inches, and are from one-quarter to one-half inch in diameter. They are planted *between the rows* of cabbage or cauliflower as soon as these crops are planted in the spring, and about the same distance apart between

the plants. The set or root is planted perpendicularly, three inches under the surface. There is no danger in planting the sets thus deep, for horseradish is particularly tenacious of life, and will start and push through the soil even if planted much deeper. The motive in planting it under the surface is to delay its starting, so as not to interfere with the cabbage crop, which may close over it without any injury whatever to the horseradish. It sometimes happens, however, either from planting too near the surface, or by the sets being very strong, that the horseradish grows so strongly as to interfere seriously with the cabbage crop. In such cases it must be cut off by the hoe, and this will not injure it in the slightest degree. We have often had to hoe it off twice before the cabbage crop was ready. It will be borne in mind that it is the root only of this crop that is wanted, and that, being grown mostly in the late summer and fall months, the removal of the leaves in June, or July even, does not in any way affect the crop.

“As soon as the cabbages have been cut off the stumps are dug up, and the ground deeply hoed, so as to encourage the growth of the horseradish crop. This rarely requires to be done more than once, the rapid growth of the leaves smothering all weeds. It attains its full growth of root by the end of October, when it may be dug up; but, being an entirely hardy plant, we usually defer lifting it until all our more tender vegetables are secured, so that the time of digging it up is usually in November and December. It is then placed

in pits adjacent to the vegetable house, so that it can be got at conveniently, and trimmed during leisure time in winter. Its preparation for market is very simple, being merely trimming off the small roots (which are kept for next season's planting), washing, by rinsing them around in a large tub; weighing—for it is all sold by weight—and packing in barrels.

“The average weight per acre is four tons, and for the past five years it has sold for \$200 per ton, or \$800 per acre. During March of last year it sold as high as \$250 per ton. I have always considered it the most safe and profitable crop of our gardens.”

Whether these results could be obtained if the production of horseradish were largely increased, it is impossible to say; but there is no doubt that its cultivation will remain fairly remunerative.

Sweet herbs are a safe crop to raise, even at a distance from market, as they can be dried and stowed away in a loft until the leisure time of winter allows them to be bunched and packed for shipment. Henderson estimates the average yield per acre at \$500.

The varieties usually grown for commercial purposes are thyme, sage, summer savory, and sweet marjoram. The cultivation of all of these is precisely the same.

The plants are raised from seed sown in April in a very fine and rich seed-bed, and they are planted out in the field, at any time after they are large enough up to the last of July,

in rows about 12 inches apart, and at somewhat less than this distance in the row. They should be kept free from weeds until they cover the ground. At this stage each alternate row should be cut out, after which the crop will spread and occupy the whole ground again, and in very favorable seasons it will sometimes close up after alternate rows have been taken out a second time.

Mangel-wurzel (or field beet) is a safe crop for the farmer to raise, inasmuch as it is the best of all the roots for cattle food; and, in rich ground, it produces enormously, while it does not interfere with the growth of cabbages the following year.

For a second crop the plants should be raised from seed planted very early in May, and it should be set out at distances of 30 inches by 15 inches. It is a perfectly safe and easy crop to transplant, if care be only taken not to attempt the operation until the roots are at least as thick as the little finger.

The distances recommended as the best ones at which to set the plants are larger than are usual in this country, but on land so rich as that under consideration, the leaves will cover the whole space, and the roots will grow to an enormous size, giving a larger yield than if more thickly set out.

Sweet corn is a fair crop to raise for market, but its cultivation is so well understood by all that it is only necessary to say here that it should follow the spinach and the onions, which are the first out of the ground in June.

Common turnips are the poorest paying of all the articles recommended for a second crop, but they are also raised with very little trouble, and as the seed may be sown at any time in July, they are often available to follow the last removed of the first crops, except the Brunswick cabbages, and these will not usually be cleared off in time to prepare the ground for anything but spinach for the following spring.

Profits.—This is hardly a safe subject for estimate; so much depends on the land, the situation, the man, and the market, that one will gain where another would lose, and the ratio of profits will vary from zero to an almost fabulous amount. However, under any favorable circumstances, a man tolerably well qualified for the business, *provided he will use manure with what he may think a wasteful hand*, might expect about the results of the following table, for an average of ten years.*

Expenses:—

Rent and taxes, say.....	\$30
Interest on cost of improvements and tools, say on \$800, at 7 per cent.....	56
Wear and tear.....	100
Manure (2 acres).....	160
Labor (equal to two men for the whole year)	1000
Seeds and plants.....	50
Total.....	\$1,396

* The first year, the outlay for manure will be more, and, owing to the crude condition of the soil, the returns will be less.

Receipts :—

From use of 50 sashes, as per previous estimate.....	\$680 00
From 450 tomato vines on the fences (say 25c. each).....	112 50
1 acre, 10,000 cabbages at 8c.....	800 00
3 tons horseradish (2d crop).....	500 00
$\frac{1}{4}$ acre beets, 300 dozen bunches at 75c.	225 00
$\frac{1}{4}$ " onions, 500 dozen bunches at 50c.	250 00
$\frac{1}{4}$ " spinach, 50 barrels, at \$2.....	100 00
$\frac{1}{4}$ " parsnips, 200 bushels at 75c....	150 00
Radishes from among beets and cabbages, say.....	100 00
$\frac{1}{4}$ acre sweet herbs (2d crop).....	100 00
$\frac{1}{4}$ " sweet corn (2d crop).....	25 00
$\frac{1}{4}$ " mangel-wurzel, say 250 bushels at 40c....	100 00
$\frac{1}{4}$ " common turnips.....	25 00

Total..... \$2,487 50

Deduct expenses 1,396 00

Net profits.....\$1,091 50

Of course there are chances that the profits will be much less than the above amount, but there are at least equal chances that they will greatly exceed it.

STEAM CULTIVATION.


For many years it has been a dream of American inventors to devise some means by which a locomotive steam-engine could be made to take the place of the team in ploughing.

Thus far, although some of the devices have been made to work tolerably well, none of them have achieved such success as to commend them to general use.

It has fallen to the lot of England to make the first application of steam to ploughing that has been so decidedly successful as to come into very general use. They have abandoned the idea of making the steam-engine travel at the front of the plough, and place it on one of the headlands, broadside to its work, an "anchor" standing opposite to it on the other side of the field.

Under the engine there is a horizontal windlass, five feet in diameter, and a similar windlass is attached to the anchor. A steel wire rope passes around these two windlasses, its ends being fastened to the carriage to which the ploughs are suspended, and which forms a link in the endless chain.

The windlass under the engine is so arranged that it clasps the rope firmly on those parts where its pulling force is exerted, and lets go as the rope leaves it in its movement toward the anchor.



FOWLER'S PATENT STEAM PLOW IN OPERATION.

The ploughs are set in "gangs" on a tilting frame. One end of the frame carries right hand, and the other end left hand ploughs. The ploughman sits on the end of the frame which is "in work," and guides the carriage by means of a steering wheel. His weight holds the end on which he sits down to its work, and tilts the other end up, so that its ploughs are in the air. If the width of the field is considerable, "rope porters" or guiding wheels keep the rope from running on the ground, and thus save power and prevent wear and tear.

The ploughs being ready to commence their work at the side of the field next to the engine, this is set in motion and the ploughs are drawn toward the anchor; when they arrive at the anchor side of the field, the ploughman changes his seat to the other end of the frame, and the engine is reversed, drawing the ploughs toward it; and in this manner they are moved back and forth until the whole length of the field is ploughed. They are then moved to the ends of the headlands and these are ploughed.

The engine is a locomotive, and advances along the headland so as to be always opposite its work, and the anchor is moved at the pleasure of the operator, by the action of its windlass.

The ploughs are used in all cases where there is a sod or a long stubble to be turned under, but fallow land is cultivated by the substitution of long-toothed grubbers, which work at a greater depth.

The construction of the steam ploughing apparatus, and its mode of operation, are shown in the illustrations which accompany this article.

Among the advantages claimed for it are the following:—

1. Greater rapidity of work, allowing land to be speedily prepared for the crop while in the proper condition, thus greatly lessening the danger that planting will be delayed by rains.
2. Cheapness of work—the cost (in England) being reduced from about \$5 per acre, the cost with horses, to about \$1.25, the cost with the steam apparatus.
3. Improved condition of the land.
4. Better drainage.
5. Greater activity in the performance of all the work of the farm.

Concerning *rapidity* of work, it may be stated that a 14-horse engine set will *plough* from 9 to 12 acres per day, and do the work better (deeper) than it can possibly be done with any ordinary farm team.

At the Annual Show of the Royal Agricultural Society at Bury St. Edmonds, in 1867, Fowler's cultivator smashed up light stubble-land at the rate of 50 acres per day of 10 hours, and did the work at a cost of about 25c. per acre, including all charges for fuel, wear and tear, and attendance.

Anything which places it in the power of the farmer to prepare his land for planting at so rapid a rate as even 8

acres per day, must do much to free him from the annoyance of frequent delays from wet weather at a time when it is important that everything proceed rapidly.

The comparative cost of cultivation, when done by steam instead of horses, is, of course, dependent on circumstances. On small farms, and for use in small fields of irregular shape, the cost of maintaining an expensive set of machinery, and the time lost in moving from one field to another, would more than make up for any saving in the actual cost of the work. On farms having 250 acres of land under the plough, and having few fields of less than 10 acres, the saving in cost of work would be very great.

This saving of cost, however, is of minor consequence as compared with the other advantages of steam cultivation.

The improved condition of the land, including its better drainage, is the great argument in favor of the process.

The movement of the ploughs is nearly twice as rapid as that of the horse-plough, and the furrow, instead of simply being turned over, is thrown from the mould-board so rapidly that it is much more thoroughly pulverized. As the furrows are all laid in one direction, there are no dead furrows left when the work is done. In the ordinary ploughing of an acre of land it receives 350,000 foot-marks per acre, one-half of these being upon the earth at the bottom of the furrows, which in time becomes compacted to an almost water-tight condition. In steam ploughing, the land is not touched by a hoof, and when (as is often the case) all the operations of

harrowing, rolling, and seed-drilling are done by steam, it is left in a condition most favorable to the growth of the crop, and to the rapid subsidence of water of rains—assuming that the land is either naturally or artificially under-drained.

Not the least benefit of steam cultivation (accompanied by the use of the steam-engine for threshing, grinding, fodder-cutting, &c.) is found in the greater activity which is imparted to all the business of the farm. The same difference, but in less marked degree, is to be observed in the use of horses instead of oxen.

The motive power sets the *time* of the whole establishment, and as the use of oxen leads to a slow, drawling, listless habit, so steam gives an activity and bustle to everything which makes wages and board tell with better effect on the year's performances.

In the Journal of the Royal Agricultural Society, for 1867, there is a very elaborate report of the results of the examinations of the committees which had been appointed "to inquire into the results of steam cultivation in use by 135 farmers and stock companies in England."

The following are some of the conclusions at which they arrived :—

"In nearly all the cases reported it will be seen that the expenses of cultivation are very much reduced, and yet that a larger amount of produce is said to have been realized.

"Not only are the operations themselves better done, quicker done, less expensively done, but all kindred and col-

lateral movements have had imparted to them a speed and 'whirr' characteristic of steam; men acquire the habit of doing the day's work in the day, and of not leaving it for the morrow. The day's labor, too, on a steam farm represents more work with less distress to the physical frame of the laborer, and better remuneration. Steam is working a revolution—slightly manifested as yet, so that we can only speak of tendencies in farm practice, and in the character of the rural population. They are being trained for the era of machinery in agriculture.

"In most cases an increase of produce, in some instances as much as 8 bushels per acre (of wheat), has resulted from steam cultivation.

"We may state as our general conclusion that steam tackle, whether of Fowler, Howard, Smith, or other makers, is now so far perfected and settled in form and details, that it may be classed among old-established, standard farm machinery, and no longer among the novelties of the day.

"We find, as the result of experience, that which we already anticipated theoretically, viz., that the increased depth of surface and the absence of pressure greatly increase the absorbing powers of the soil, and consequently assist the action of the drains.

"Mr. Wm. Smith, of Woolston, England, was one of the pioneers of steam cultivation, and is still one of its most zealous advocates. A short time ago he extended an invitation to all who were interested in the subject to visit his

farm and witness the operation of his tackle, and to see its effect. He communicates to a London paper the substance of the statements he made to his visitors, and from this I extract the following, as serving to illustrate the completeness with which the system has been tried and found satisfactory :—

“You must see that these fields are not only heavy clay, but hilly and uneven, and the face of them shows that they are well drained as well as well cultivated.

“This field, No. 3, on which you stand, together with No. 2, through which you have passed, and No. 1 (light land), which I will hereafter show you, contain 32 acres, and were smashed by steam-power on the 31st of August, and the 1st, 2d, and 3d of September, at the following cost :—

Labor.....	£3 14 0
Coal.....	1 12 0
Oil.....	2 6
Interest on money, and wear and tear.....	2 9 6
	<hr/>
	£7 18 00

Or 4s. 9d. per acre (about \$1.15).

* * * * *

“Now I will let you know what the operations and cost of seed-beds have been on these four fields under steam culture for 14 years, taking those on field No. 3 to represent the lot :—

“10 steam-power smashings, 2 ridgings and subsoilings

by steam-power, 2 cultivatings and drillings at one operation, each by steam-power; 1 cross cultivating by steam-power, 1 cross cultivating and seeding at one operation by steam-power, 7 horse cultivatings, 1 horse subsoiling, 1 ridge ploughing by horses. The total cost of these operations has been £6.11.9, or 9s. 5d. per acre as the average cost of a seed-bed, exclusive of planting or drilling, except those planted by steam-power.

"The cropping on No. 3 during that period has been 1 of peas, 2 of barley, 5 of beans, 5 of wheat, and 1 of Swedes.

"The average yearly produce under steam culture has been, on these four fields, quite 14 bushels per acre more than it had been under horse culture.

* * * * *

"Now let us look into the working of the tackle since the 5th of October last. On that day I started it on No. 4 (heavy land), 10 acres, ridging and subsoiling it for beans. It was finished on the 6th at 4.10 P.M. The depth of work is 9 inches; the consumption of coal 1 ton; and the pressure on the engine 60 lbs.

"We then shifted the tackle nearly half a mile to No. 1 (heavy land), 8 acres, and we finished that field at 12.15 P.M. on the 8th. Depth of work 10 inches; consumption of coal 16 cwt.; pressure on the engine 65 lbs.

"We then shifted the tackle a quarter of a mile to No. 3 (light land, part 1), 5 acres, and finished it on the 9th at 12.20 P.M. Depth of work 10 inches; consumption of coal 9 cwt.; pressure on engine 60 lbs.

"We then shifted the tackle a mile to No. 6 (light land), 14 acres, working all day on the 10th (the 11th was Sunday), working again all day on the 12th, and we finished on the 13th at 8.40 A.M. Depth of work 11 inches; consumption of coal 29 cwt.; pressure on engine 65 lbs.

"We then shifted the tackle more than a mile to No. 3 (light land, part 2), 6 acres, and finished it on the 14th at 11.50 A.M. Depth of work 10 inches; consumption of coal 10 cwt.; pressure on engine 60 lbs.

"We then shifted the tackle to where it now stands, on No. 4 (light land), for you to see it at work.

* * * * *

"It is not a set of new-fangled tackle, got up for the purpose of racing, for the common portable engine has done 10 years' hard work. It has done, in addition to my ploughing, a lot of threshing and grinding, yet it is in capital trim. The cost of repairs has been but a mere trifle.

"The windlass has had 10 years' work. In 1858 it did 55 acres for the late Prince Consort, on the Flemish farm, Windsor, and I have worked it on my farm ever since.

* * * * *

"I have worked the rope 7 years. The first year it got out several times. At one or other of the splices it pulls in two; indeed it has not, during these 9 days' work, gone through a day without a break; therefore the men have had the mending of the ropes to do as well as the ridging and subsoiling.

* * * *

"My average quantity of work per day is much greater,

and my average consumption of coal per acre is much less, than it has ever been before. This is mainly due to my land having been deeply worked so many times before.

“The sum total of all this evidence proves plainly that the Woolston system of applying steam-power to the cultivation of the soil gives clean dress and cheap seed-beds, and that fancy tackle is not needed on show days. The boy that you see working the implement is only 14 years old. He has done all my work this year, and well too. The work is before you to speak for itself.”

It is found, for use in neighborhoods where the farms are small, that it is the best plan to form joint-stock companies to own and operate the tackle—hiring it out by the day or by the acre, and giving the precedence to stockholders. This plan would work the best among the smaller farmers of our Eastern States—but at the West, where the proprietorships are larger, it will be most advantageous to have the apparatus, with its engine to do other work, attached to the farm.

It is sometimes objected that much of the land in this country is too rough and too stony for the steam-plough ever to gain a foothold. The same objection was made twenty years ago to the use of the mowing machine in New England, and there is every reason to suppose that when the advantages of the steam-plough are once fully realized, even the hillsides of Vermont will smile under its influence.

HOUSE PAINTING.

The following receipts and directions are condensed from a practical English work on the art of house painting. They are principally designed for the inexperienced and those who, living at a distance from cities, have great difficulty in obtaining first-class workmen.

To make the work satisfactory, it is very necessary for the workman to have *very clean* all the vessels, brushes, and cans he may require in the course of his work, such as the various paints, pots, or vessels in which he mixes or from which he uses his colors. These are sometimes bought at the shops, handsomely made of stout tin, and such are easily kept clean, and save their expense in color, which is readily brushed down their smooth sides. He will also require a marble slab and muller, to grind the finer colors used in painting. Sometimes a small cast-iron mill is useful not only to grind colors, but to pass the tinted color through, so that it may be more thoroughly mixed.

It is presumed the workman will know what brushes he will require, according to the work he has in hand.

In preparing to paint a good dwelling, after having obtained the necessary colors and brushes, see that you have a few pounds of good pumice stone, a quire or two of assorted sand paper, to smooth the inequalities in the work; some

twenty pounds of putty, to stop up after the first coat in every part of the house; a sufficiency of fine slaked lime, and a proper number of large and small vessels, to mix the colors in and use it from; a few pounds of soaked glue, &c.

If the wood-work be new, and no wall work required, you will go over it carefully with a small brush, and some of the glue size, colored with red lead, covering what knots and stains may appear in the wood, after which the priming coat of almost all oil, and good white lead, tinted with Indian red, should be evenly brushed over the work; and, as soon as dry, the putty knife and putty should follow, to stop all the cracks and nail holes. Then should follow the second coat, with a little spirits of turpentine in the oil, and the color slightly tinged with blue black. This is generally thought sufficient for the attic and third stories. But the rest of the house is usually finished with old ground white lead, thinned with spirits of turpentine. The roof, if covered with tin, should be painted once in three years. There are many different methods in use. Some paint with raw oil, dry Spanish brown and a little red lead, to dry it, for fear of a rain; others, with Spanish brown, more red lead, and half whale oil with the linseed oil; others use yellow ochre and black, mixed in the same oils; others use a roof paint, made by boiling paint skins in whale oil, and carefully straining them while warm, reserving the remaining skins, to stop the leaks around chimneys and dormer windows. This last mentioned paint is probably

serviceable from its elasticity. In the country, many paint their roofs and out-buildings in the same way, using sometimes Venetian red from its brightness.

Many complaints are continually made that white lead, and colors composed thereof, do not endure, and are quickly beaten off by exposure to the sun and rain. This difficulty occurs as much from the manner of using the paint as from its quality. As this occurs in outside work, it is to be attributed, first, to the condition of the work to be painted, being generally in such a state as to absorb the oil from the first coat, thereby leaving it in a dusty state, and liable to be washed off by the first rain. This can be guarded against, only by filling the old work, in painting two thin coats over it, one upon the other, as soon as dry; and finishing it with one thicker coat, to protect it and shed the rain. A fourth coat, if the immediate expense is not heeded, will repay its cost in additional service and beauty.

The white lead can be procured of any requisite quality at the color stores. It is thought that the best article is the most economical, as it works out with more ease, and repays the difference of cost in its appearance. Linseed oil is also better for having due age, for the same reasons as the white lead, working with softness and advantage after parting with the water which is generally combined with new oil.

The quality and fineness of the white lead used adds materially to the work, and that which is well ground, and

has such mellowness from age as will cause it to work smoothly under the brush, in connection with good linseed oil, will certainly repay any reasonable additional cost. The first coats should always be mixed with clear linseed oil; the fourth coat may be used with boiled oil and one-quarter part spirits of turpentine.

Putty is best purchased at a good color store, where you can depend upon its being made of good dry whitening and linseed oil. It should be carefully and freely used after the work has had one coat of paint, for the fresh paint holds the putty very firmly.

Harmony of Colors.—Red looks well with blacks, whites, or yellows. Blues harmonize with whites and yellows. Greens, with whites, black or yellow. Gold, with blacks or browns. White appears well with any color. Purple, pink and white, &c., &c.

MIXING PAINTS.

A Beautiful White Paint.—For inside work, which ceases to smell, and dries in a few hours. Add one pound of frankincense to two quarts of spirits of turpentine; dissolve it over a clear fire, strain it, and bottle it for use; then add one pint of this mixture to four pints of bleached linseed oil, shake them well together, grind white lead in spirits of turpentine, and strain it, then add sufficient of the lead to make it proper for painting; if too thick in using,

thin with turpentine, it being suitable for the best internal work on account of its superiority and expense.

For a Pure White Paint.—Nut oil is the best ; if linseed oil is used, add one-third of turpentine.

To Mix Common White Paint.—Mix or grind white lead in linseed oil to the consistency of paste, add turpentine in the proportion of one quart to a gallon of oil ; but these proportions must be varied according to circumstances. Remember to strain your color for the better sorts of work. If the work is exposed to the sun, use more turpentine for the ground color to prevent its blistering.

For Knotting.—Mix white or read lead powder in strong glue size and apply it warm.

Common Flesh Color.—Stain your white lead with red lead, and mix with oil and turps.

Fine Flesh Color.—Is composed of white lead, lake and vermilion.

A Beautiful Color for Carriages, &c.—Mix carmine lake with black japan.

Cream Color.—This is a mixture of chrome yellow, the best English Venetian red, white lead, and red lead, in oil.

Pearl Gray.—White lead, with equal portions of Prussian blue and lampblack, mix with oil and turps.

Fawn Color.—Grind some burnt and raw terra sienna very fine. Two or three pounds of this is sufficient to stain

white lead for a large building. This color is of a superior shade, and very excellent for inside work.

Blue.—Grind Prussian blue in turps; other blue very fine in linseed oil, and mix it with white paint to the tint required.

Buff.—This is a mixture of French yellow, chrome yellow and white lead, tinged with a little Venetian red, oil and turps.

Straw.—A mixture of chrome yellow and white lead, oil and turps.

Drab.—Raw and burnt umber and white lead, with a little Venetian red, linseed oil and turps. *Another.*—Burnt umber and white lead, with a little Venetian red, oil and turps, as before.

Steel.—Mix white lead, Prussian blue, fine lake and verdigris, in such proportions as to produce the required color.

Purple.—White lead, Prussian blue and vermilion, or lake, with oil and turps.

Violet.—Is composed of vermilion, mixed with blue-black, and a little white.

French Gray.—White lead and Prussian blue, tinged with vermilion; and for the last coat substitute carmine for the vermilion. Mix with oil and turps.

Silver.—Use white lead, indigo, and a small portion of blue-black, as the shade may require.

Gold.—Mix Naples yellow or patent yellow with a small quantity of orange chrome and a little Spanish white.

Dark Chestnut.—Mix red ochre and black. Use yellow ochre when you require to lighten the color, in oil.

Salmon.—White lead, tinged with the best English Venetian red, oil and turps.

Peach Blossom.—White lead, tinged with orpiment; mixed with oil and turps.

Drab.—White lead with a little Prussian blue and French yellow, linseed oil and turps. *Another.*—White lead, with a little French yellow and lamp-black, linseed oil and turps. *Another.*—White lead with a little chrome green and blue-black.

Lead.—This is a mixture of lamp-black and white lead, with a little litharge.

Chocolate.—Mix lamp-black and Venetian red with a little red lead, or litharge, to harden the color and give a drying quality. The colors must be ground, and mixed with boiled oil and a little turps.

Dark Red, for Common Purposes.—Mix English Venetian red in boiled oil with a little red lead and litharge, to give a drying quality.

Orange.—Mix red lead and French yellow with linseed oil and turps, or use deep chrome yellow.

Bright Yellow for Floors, &c.—White lead and linseed oil, mixed with some French yellow, and a little chrome

yellow to brighten it ; some red lead, burnt white vitriol and litharge added to give it a very drying quality. This color mixed with equal parts of boiled oil and turpentine, and used very thin.

Dark Yellow.—Mix French yellow in boiled oil, adding to it a little red lead and litharge, to give the paint a drying quality.

Light Yellow.—This is a mixture of French yellow, chrome yellow and white lead, with oil and turps. *Another.*—French yellow, white lead, and red lead. *Another.*—Grind raw terra sienna in turps and linseed oil ; mix with white lead. If the color is required of a warmer cast, add a little burnt terra sienna ground in turps.

Olive Green.—A suitable, cheap, and handsome color for outside work, such as doors, carts, wagons, &c.

Grind separately Prussian blue and French yellow in boiled oil, then mix to the tint required with a little burnt white vitriol to act as a drier. *Another.*—Black and blue mixed with yellow, in such quantities as to obtain the colors or shades required. For distemper, use indigo and yellow pink mixed with whiting or white lead powder. *Another.*—This is a mixture of Prussian blue, French yellow, a small portion of Turkey umber, and a little burnt vitriol. Grind the same way. *Another, in oil.*—Mix Prussian blue and chrome yellow. Grind the same. *Another shade.*—A mixture of Prussian blue and French yellow, with a small

Quantity of white lead and Turkey umber and burnt white vitriol. Grind the same.

Light Green.—White mixed with verdigris. A variety of shades may be obtained by using blue and yellow with white lead.

Grass Green.—Yellow mixed with verdigris. *Another.*—Mix one pound of verdigris with two pounds of white lead. Walnut oil is the best for this purpose.

Invisible Green, for outside work.—Mix lamp-black and French yellow, with burnt white vitriol. These colors mix in boiled oil. Burnt vitriol is the best drier for greens, as it is powerful and colorless, and consequently will not injure the color.

To Paint a Bronze.—Grind good black with chrome yellow and boiled oil; apply it with a brush, and when nearly dry use the bronze powder at certain parts and the edges also; the effect will be a brassy hue.

A Good Imitation of Gold.—Mix white lead, chrome yellow, and burnt sienna, until the proper shade is obtained.

Tar Paint for Fences, Roofs, &c.—Common tar mixed with whiting. Venetian red or French yellow, according to the color required. This should be warmed in a large iron kettle in the open air, and applied with a large painting-brush. It is an excellent preservative of the wood, and looks well for rough work.

Paint Driers.—*Litharge.*—This is a useful drier, and

may be used in all kinds of paints, except greens and very delicate colors. *White Vitriol or Copperas*.—This turns into water, especially when used in black paints; and is almost useless for any color till the water of crystallization is evaporated, and then it becomes a powerful drier, and may be used for every delicate color, as it is perfectly transparent; but when used in its raw state in white paint, has the effect of turning it yellow. *Sugar of Lead*.—This is a very useful and transparent drier, not so powerful as white vitriol, but it may be used with it to advantage.

Milk Paint for In-door Work.—The quantity for one hundred square feet:—One quart of skimmed milk, three ounces of lime, two ounces of linseed or poppy oil, one pound and a half of Spanish white or whiting. Put the lime into a clean bucket, add sufficient of the milk to slake the lime, add the oil a few drops at a time, stirring the mixture with a flat stick till the whole of the oil is incorporated in the mass; then add the remainder of the milk, and afterwards the Spanish white or whiting, finely powdered, and sifted gently over the mixture by degrees. Curded milk will do for the purpose, but it must not be sour. One coat of this will do for ceilings and staircases in general; two coats or more for new wood. Where color is required, you may use powdered umber, ochres, chromes, greens, blues, pinks, &c., &c., ground in milk. For particular work, strain the color through a hair sieve.

Lime Whitewash.—Lime whitewash is made from lime

well slaked. Dissolve two pounds and a half of alum in boiling water, and add it to every pailful of whitewash. Lime whitewash should be used very thin, and when it is sufficiently bound on the wall by means of alum, two thin coats will cover the work better; this may be used for the first coat, thinned with water. Most whitewashers apply their wash too thick, and do not mix a proportionate quantity of alum to bind it, consequently the operation of the brush rubs off the first coat in various parts and leaves an uneven surface, and the original smooth surface of the wall is entirely destroyed.

Italian Marble.—This looks bold, and is well adapted for columns, &c., and is easy to imitate. The ground a light buff. For the graining colors, prepare a rich, warm buff, made in the following manner: Mix stiff in boiled oil white lead and good stone ochre, and tinge with vermilion, then grind some burnt terra sienna very fine in burnt oil, and put it into another pot; mix some pure white stiff in oil, and keep this separate. Thin these colors with turpentine, have ready a brush for the buff, and another for the terra sienna. Proceed to work as follows: Take the brush intended for the buff moderately full of color, and dab it on freely and carefully in different patches, some of them larger than others, and varying them as much as possible. When these are laid on, take the other brush and fill in with the terra sienna the spaces between; as soon as this is done, take a dry duster or softener and blend the

edges together, making it appear as soft as possible. Proceed in this manner till the whole is finished, then take a hair pencil and draw a few thin white veins over the work, varying them as much as is necessary; take another pencil for the terra sienna, and run a few thin lines intermixing with the whole; varnish when dry.

To Imitate Granite.—For the ground color, stain your white lead to a light lead color, with lamp-black and a little rose-pink. Throw on black spots with a graniting machine, a pale red, and fill up with white a little before the ground is dry.

A Cheap Oak Varnish.—Two quarts of boiled oil, one and a half pound of litharge, three quarters of a pound of gum shellac, one ounce of gum. All boiled together, and stirred up till dissolved, then take off the fire and add two quarts of turps. When settled, strain into a bottle and cork for use.

Common Oil Varnish.—Take one gallon of quick drying oil, two pounds of resin, and one quart of turpentine; put the resin with the drying oil into a varnish kettle, and let it dissolve in a gentle heat; take it from the fire and gradually pour in the spirits of turpentine. If too thick add more of the turpentine.

Transparent Varnish for Pictures.—Take the white of four eggs and two ounces of loaf sugar; beat them up in lime water to the proper consistency for varnishing.

For Varnishing on Wood, Unpainted.—Quarter of a pint of wood naphtha, quarter of a pint of spirits of wine, four ounces of benzoin, four ounces of orange shellac, added all together. If not thick enough with those ingredients for your purpose, add more of the gums benzoin and shellac.

Waterproof Varnish, for Linen or Calico.—One pint of turpentine, one and a half pint of linseed oil, seven ounces of litharge, one ounce of sugar of lead. Strain it, apply it with a brush, and dry it in the sun or in a warm place.

Instructions.—Oil of turpentine deadens the color of paints; varnishes, copal, &c., brighten the color.

SOLDERS.

For lead solder.—Melt 1 part block tin, and when fused, add 2 parts of lead. Use resin with it.

For tin solder.—Melt 4 parts of pewter, 1 part of tin, and 1 part bismuth together. Use resin with it.

CEMENTS.

Glue.—Melt 1 lb. glue in 2 quarts warm water. For a glue that will resist the action of water, boil 1 lb. of glue in 2 quarts of skimmed milk. Pulverized chalk added to glue strengthens it.

Soft cement.—For boilers, steam-pipes, &c.: 4 parts red or white lead, ground in oil, with 2 or 3 parts iron filings.

Hard cement.—Mix iron borings or filings with salt water, then add a small quantity of sal ammoniac with water.

Hydraulic cement—for cisterns, sewers, cellars, pipes, &c., is purchased by the barrel, which contains 300 lbs.

Dry cement—which will resist the weather equal to marble, is made of 2 parts sifted ashes, 3 parts clay, and 1 part sand, mixed with oil, and applied while soft.

Brown mortar, for masonry, brick-work, &c.

Mix 1 part lime, 2 parts sand, a small quantity of hair with water.

CONTENTS.

ALPHABETICALLY ARRANGED.

A.

	PAGE
ACCOUNTS, KEEPING OF.....	227
Accounts by single entry, with examples.....	227
" double " " " 	229
Form of bill of sundries.....	231
" receipt in full.....	237
" check.....	238
" due-bill.....	238
" promissory note.....	238
" " " with surety.....	238
" draft or bill of exchange.....	239
Explanation of all the above.....	240
ALCOHOL, PROPORTION OF, IN LIQUORS.....	190
ANGULAR MEASURE, Illustrated.....	23
ANIMALS, LIFE AND INCREASE OF.....	197
Table, showing the period of reproduction and gestation of domestic animals and fowls.....	198
Table, showing when forty weeks (the period of gestation in a cow) will expire, from any day throughout the year.....	199
Growth and life of animals.....	199

	PAGE
ANIMALS, AGE OF, Illustrated.....	201
To find the age of a horse.....	201
“ “ “ “ “ cattle.....	206
“ “ “ “ “ sheep.....	208
“ “ “ “ “ goats.....	208
ANIMALS, FOOD OF, Illustrated.....	212
Table, showing the comparative difference between good hay and other food for stock—being the results of experiments.....	212
Table, showing the comparative difference between good hay and other food for stock—being the mean between experiment and theory	213
Table, showing the quantity of food different animals require per day to each 100 lbs of their live weight.....	213
Table, showing the daily food required by the ox	214
Table, showing the effects produced by an equal quantity of dif- ferent kinds of substances as food for sheep.....	215
ANIMALS, COMPUTED WEIGHT OF.....	209
ANNUITIES.....	219
Table, showing the amount of \$1 for any number of years from 1 to 24, at 5 and 6 per cent. compound interest.....	219
Table, showing the present worth of \$1 annuity at 5 and 6 per cent. compound interest for any number of years from 1 to 34,	218
APOTHECARIES' WEIGHT, table.....	158
“ fluid measure, table.....	156
ARITHMETICAL CHARACTERS, EXPLANATION OF.....	14
ARTIFICIAL MANURES.....	357
AVOIRDUPOIS WEIGHT, table.....	152

B.

	PAGE
BALANCES, FALSE, Illustrated.....	84
To detect false balances.....	84
To find the true weight.....	84
BOARD FENCE (see fences).....	125
BOARD MEASURE.....	62
BOOKS, SIZES OF (see printing).....	260
BONDS—U. S. BONDS.....	242
BOXES, CAPACITY OF.....	81
BRICK-WORK (see masonry).....	276
BUTTER, PROPERTIES AND COMPOSITION OF.....	387
BUTTER AND CHEESE-MAKING.....	391
The butter dairy.....	391
The milk room.....	391
Cleanliness.....	392
Setting the milk.....	392
Cream-churning.....	393
Packing for market.....	394
Test of good butter.....	396
The cheese dairy.....	397
Quality of cheese.....	397
To construct an ice-house for the dairy.....	399
Analysis of butter.....	390

C.

CAPACITY OF BOXES.....	81
" WAGON-BEDS, Illustrated.....	82
CASK-GAUGING, Illustrated.....	78

	PAGE
To find the contents of a cask by three dimensions....	79
“ “ “ “ “ “ “ four “	79
CATTLE, SOILING.....	401
CATTLE, TO FIND THE AGE OF.....	206
CATTLE, COMPUTED WEIGHT OF, Illustrated.....	209
Table, showing the compute weight of cattle from their girth, &c.	211
CEMENTS.....	473
Glue.....	473
Soft cement.	473
Hard cement.....	473
Hydraulic cement.....	473
Dry cement.....	473
Brown mortar for masonry, brickwork, &c.....	473
CIRCLES.....	296
To find the circumference of a circle....	296
“ “ diameter “	296
“ “ area “	296
To find the side of an equal square containing the same area as a given circle.....	297
To find the solidity of a sphere... ..	297
Table, showing the areas of circles and the sides of their equal squares, from 1 to 100.....	298
To find, by means of the table, the square or circle that will con- tain the area of a board or surface of given length and width. .	302
CIRCULAR MEASURE, Illustrated.....	23
CISTERNS, Illustrated.....	86
To find the number of gallons in square or oblong cisterna....	86

CONTENTS.

479

	PAGE
To find the number of gallons in triangular cisterns.....	86
" " " " circular " 	87
Table, showing the contents of circular cisterns from 1 foot to 25 feet in diameter for each 10 inches in depth.....	87
To find the number of gallons in tub-shaped cisterns.....	88
To ascertain the size of cisterns adapted to roofs.....	89
Table, showing the contents of circular cisterns in barrels for each foot in depth.....	90
To construct filtering cisterns to furnish pure water for domestic use.....	91
CHARCOAL, TO PREPARE.....	115
CHEESE DAIRY (see butter and cheese).....	397
CLOTH MEASURE, table.....	169
COKE.....	118
COMMERCIAL ABBREVIATIONS.....	18
COMPOUND INTEREST, table of.....	218
CONTENTS.....	475
CORN ON THE COB IN CRIBS, TO MEASURE, Illustrated.....	57
When the crib is equilateral.....	57
When the crib is flared at the sides.....	58
CORN, RELATION OF PORK TO.....	194
Table, showing price of pork per lb. at different prices per bushel for corn.....	194
To find the price of pork, the price of corn being given.....	195
To find the price of corn, the price of pork being given.....	195
CROPS, ROTATION OF.....	378
Rotation of field crops.....	385

	PAGE
Rotation of garden crops.....	386
Chemical theory of rotation.....	381
CROPS, NUTRITIVE VALUE OF.....	190
CUBIC MEASURE, table.....	171
To find the cubic contents of any solid body.....	171
CUBES AND CUBE ROOTS, TABLE OF, FROM 1 TO 1000.....	303
CULTIVATION, STEAM, Illustrated.....	450
D.	
DECIMALS—FRACTIONS.....	257
To reduce fractions to decimals.....	257
To add decimals.....	257
To subtract decimals.....	258
To multiply decimals.....	258
To divide decimals.....	259
Table of useful decimals.....	259
DECREASE AND EXPECTATION OF HUMAN LIFE.....	216
Table, showing the decrement and expectation of human life...	216
Table of St. Maur.....	217
DEFINITIONS OF MATHEMATICAL FORMS.....	292
Parallel Lines.....	292
An Angle.....	292
A Right Angle.....	292
An Acute Angle.....	292
An Obtuse Angle.....	292
A Surface.....	292
A Triangle.....	292
The Altitude.....	292

	PAGE
A Right Angle Triangle.....	293
A Parallelogram.....	293
A Rectangle.....	293
A Square.....	293
A Trapezoid.....	293
The Altitude.....	293
A Circle.....	293
The Circumference.....	293
The Diameter.....	293
The Radius.....	293
A Solid.....	294
A Prism.....	294
Triangular Prism.....	294
Hexagonal Prism.....	294
Cylinder.....	294
Cube.....	294
A Pyramid.....	294
The Altitude.....	294
A Cone.....	294
A Frustum.....	295
An Ellipse.....	295
A Sphere.....	295
A Spheroid.....	295
DEPTH OF SOWING WHEAT.....	193
DIET, SOLID MATTER AND WATER IN ARTICLES OF.....	198
Table, showing the proportion of solid matter and water in 100 parts each of various articles of diet.....	198

	PAGE
DRAINING TILE.....	362
How to make a drain.....	364
Different kinds of tile used.....	364
Boynton's improvement in making tile.....	365
Rules to be observed in making tile drains.....	367
Size and quantity of tile required to the acre.....	368
Tools used in laying drain tile.....	370
How to make drain tile, Illustrated.....	373
Why should land be drained.....	373
The effects of drainage.....	374
DRY MEASURE, table.....	162

E.

EARTH, PRESSURE OF, AGAINST WALLS.....	255
ENGLISH MONEY, table.....	149
" Gold and silver coin.....	149
" Copper coin.....	150
Canadian currency.....	150
EXHAUSTION OF SOILS (see soils, exhaustion of).....	320
EXPECTATION AND DECREASE OF HUMAN LIFE.....	216
Table, showing the decrement and expectation of human life. .	216
St. Maur's Table.....	217

F.

FALSE BALANCES, Illustrated.....	84
To detect false balances.....	84
To find the true weight.....	84
FENCES AND FENCING, Illustrated.....	125

Rail fence.....	PAGE 126
Table, showing the number of rails, stakes, and riders required for each 10 rods of fence.....	127
Post and rail fence.....	128
Table, showing the number of posts and rails required for each 10 rods of post and rail fence.....	128
Post and board fence.....	128
To find the number of feet of boards required for each rod of post and board fence.....	129
To find the number of posts required for a given length of post and board fence.....	129
FENCES, HEDGE (see hedge plants).....	130
FENCES, WIRE.....	134
FOOD OF ANIMALS (see animals, food of), Illustrated.....	212
FOOD FOR STOCK, STEAMING.....	415
FRACTIONS (see decimals).....	257
Table of useful decimals	259
FREIGHTS, By-laws of N. Y. Chamber of Commerce.....	442
FUEL	113
Table, showing the comparative values of fire woods.....	113
Table, showing the weights per cubic foot of different kinds of coal.....	115
Properties of charcoal.	116
To prepare charcoal.....	115
Table, showing the number of parts of charcoal afforded by 100 parts of different kinds of wood.....	118
Coke.....	118

	PAGE
Table, showing the weight, evaporative powers for weight, bulk, and character of fuel.	119
Combustible matter of fuel.	120
Table, showing the heating power of different combustibles. . . .	121
Table, showing the effects of heat upon certain bodies.	121
Table, showing the relative value of different fuels by weight. . .	121
Table, showing the number of gallons of water which may be lifted to various heights by the consumption of 112 lbs. of coal	122
Table, showing the price of parts of a cord of wood at given rates per cord.	123

G.

GARDEN SEEDS, QUANTITY OF, TO PLANT, &c.	192
GARDENING FOR MARKET.	428
Size, arrangement, and equipment of the garden.	430
Construction and care of the hot-bed.	432
Profits of the same.	434
Management of field crops.	435
Vegetables best adapted for market.	437
Harvesting the crops.	440
Prices of early vegetables.	441
Profits of the business.	446
GAUGING, CASK, Illustrated.	78
GLUE, TO MELT AND APPLY.	473
GOATS, TO FIND THE AGE OF.	208
GOVERNMENT LAND MEASURE.	50
GRAIN, MEASUREMENT OF, IN GRANARIES, Illustrated.	60

	PAGE
GRAIN, WEIGHT OF, as established by the Legislatures of the different States.....	189
GRAIN, PER CENT. OF OIL IN.....	191
GRAIN, QUANTITY OF, TO SOW PER ACRE, &c.....	192
GRAVITY, SPECIFIC, Illustrated.....	182
To find the specific gravity of a body.....	183
“ “ “ “ “ “ “ lighter than water.....	183
To reduce the gravity of a body to its weight in lbs. per cubic foot	184
Table, showing the specific gravity of various bodies.....	185
To find the weight of a cubic foot of substance, the specific gravity being given.....	185
To find the number of cubic feet in any irregular body.....	186
Table, showing the weight of a cubic foot of different substances	187

H.

HAY, MEASUREMENT OF, Illustrated.....	51
To find the number of tons of hay raked into windrows.....	52
“ “ “ “ “ “ in a mow.....	52
“ “ “ “ “ “ in old stacks.....	53
“ “ “ “ “ “ in long, square stacks.....	53
“ “ “ “ “ “ when taken out of old mows or stacks.....	54
Table, showing the price per cwt. of hay at given prices per ton	54
An easy mode of ascertaining the value of a given number of lbs. of hay at a given price per ton of 2000 lbs.....	55
HEAT, EFFECTS OF, ON DIFFERENT BODIES.....	121
HEATING INCLOSED AIR.....	122

	PAGE
HEATING BY STEAM-PIPE	123
HEDGE PLANTS	130
Directions for setting and trimming.....	131
To preserve plants during the winter.....	131
Setting evergreens.....	131
Osage orange.....	132
Honey locust.....	132
Buck thorn.....	132
Privet.....	132
Hawthorn	133
Norway spruce.....	133
Arbor vitæ.....	133
Hemlock.....	133
HOP, ANALYSIS OF, Illustrated.....	323
HORSE POWER, Illustrated.....	136
HORSE POWER, origin and definition of.....	139
Table, showing the labor one horse is able to perform at different rates of speed on canals, railroads, and turnpikes.....	140
Table, showing how much one team and plough will perform in a day in acres and tenths	141
Draught of a horse.....	136
Power of the horse when aided by horse-mill.....	136
Travel per day of the horse.....	136
Burden of the horse.....	136
Endurance of the horse.....	139
To compute the power of a waterfall.....	139
" " " " of a steam-engine.....	140

CONTENTS.

487

	PAGE
To find the age of the horse.....	201
Food of the horse.....	213
HOUSE PAINTING.....	460
HUMAN STRENGTH.....	135
HYDRAULICS, Illustrated.....	93
The fundamental laws of hydraulics, &c.....	93
To find the velocity of a stream issuing from a head of water...	95
To find the head, the velocity being given.....	96
To find the quantity of water that will issue from an opening, the dimensions of the opening and the head being given....	96
To find the velocity of currents in ditches, sluices, brooks or rivers	97
To find the volume of water discharged by drains, sluices, brooks, &c., of given dimensions, in a given time.....	98
To find the velocity of water running through pipes.....	98
To find the quantity of water discharged through pipes... ..	99
To find the pressure of a fluid on the bottom of a vessel, cistern, or reservoir.....	100
To find the pressure on the side of a vessel.....	100
HYDRAULIC RAM, THE, Illustrated.....	103
To ascertain the quantity of water and the height to which it may be elevated by a given fall and volume of water.....	105
Working results of water rams now in use.....	106
HYDRAULIC PRESS, THE, Illustrated.....	110

I.

ICE HOUSE, to construct an.....	399
ICE, STRENGTH OF.....	271

	PAGE
ILLUSTRATIONS, LIST OF.....	17
INCLINED PLANE, Illustrated.....	282
INTEREST, SIMPLE.....	218
INTEREST, COMPOUND.....	218
Table of simple interest at 7 per cent. for each day to a month, from \$1 to \$100.....	222
Table of simple interest at 6 per cent. for each day to a month, from \$1 to \$100.....	224
Table, showing the interest of \$1 to \$5,000 from 1 day to 2000 days, at 6 or 7 per cent.....	220
IRON, WEIGHT OF SQUARE ROLLED, Illustrated.....	273
" " " ROUND "	275

K.

KEEPING ACCOUNTS (see accounts, keeping of).....	227
By single entry, with examples.....	227
By double entry, with examples.....	229

L.

LAND, MEASUREMENT OF, Illustrated.....	43
When the field is a square, a parallelogram, a rhombus, or a rhom- boid	44
When the field is triangular.....	44
When the field is a trapezium or trapezoid.....	45
When the field is an irregular polygon.....	45
When the field is long and the sides crooked and irregular.....	46
When the field is long and the sides and ends crooked and irregular	46
When the field is a circle.....	47

CONTENTS.

489

	PAGE
Plots containing an acre.....	47
Table, showing the square feet and the feet square of the fractions of an acre.....	48
Table, showing the number of hills or plants on an acre, for any distance apart, from 10 inches to 6 feet—the lateral and longi- tudinal distances being unequal.....	48
Table, showing the number of plants, hills, or trees contained in an acre at equal distances apart, from 3 inches up to 66 feet .	49
LAND MEASURE, GOVERNMENT.....	50
LATHS, size of, number in a bundle, &c.....	279
LATITUDE	24
LEAD PIPE, WEIGHT OF... ..	112
Table, showing the weight of lead pipe per yard from $\frac{1}{2}$ to $4\frac{1}{2}$ inches diameter.....	112
Table, showing the weight of very light lead pipe.... .	112
LEVER, THE, Illustrated.....	279
LIFE AND INCREASE OF ANIMALS, Illustrated.....	197
Table, showing the period of reproduction and gestation of ani- mals and fowls.....	198
Table, showing when forty weeks (the period of gestation in a cow) will expire from any day throughout the year.....	199
Growth and life of animals.....	199
LIFE, DECREASE AND EXPECTATION OF.....	216
Table, showing the decrement and expectation of life.....	216
Table of St. Maur.....	217
LIGHTNING RODS, Illustrated	251
To construct a lightning-rod.. ..	251

Conductors of electricity.....	PAGE 253
Non-conductors.....	253
Dr. Franklin's theory.....	253
LIQUORS, PROPORTION OF ALCOHOL IN.....	190
LIST OF ILLUSTRATIONS.....	17
LOGS REDUCED TO INCH-BOARD MEASURE.....	70
Table, showing the number of feet (board measure) of inch boards contained in round saw-logs of various dimensions...	71
LONGITUDE (see seasons, &c.), Illustrated.....	19
LONG MEASURE, table.....	167

III.

MANURES.....	327
The use of manures.....	328
Rules in deciding what manures should be used.....	331
Classification of manures.....	332
Mineral manures.....	332
Vegetable manures.....	332
Animal manures.....	332
Analysis of fish-guano.....	332
Analysis of Peruvian guano.....	333
Analysis of Bolivian guano.....	333
How to select a good article of guano.....	334
How to apply guano.....	334
Analysis of bone (crushed) manure.....	335
Table, showing the comparative value of animal manures, with farm-yard manure as the standard.....	335

	PAGE
Decomposed vegetables as manures.....	336
Distinction between animal and vegetable manures.....	336
Table, showing the relative values of decomposed vegetables as manures from the <i>inorganic</i> matter they contain.....	336
Table, showing the relative value of decomposed vegetables as manures from the <i>nitrogen</i> they contain.....	336
Analysis of a manure-heap in the condition usually applied to the field.....	337
Analysis of other specimens of fresh farm-yard manure.....	338
Analysis of green sand marl (of New Jersey).....	340
"Digestion and its products".....	341
Value of liquid manures.....	344
Poudrette and urate.....	345
Analysis of night-soil.....	346
The dry earth system.....	347
Invention of Rev. Mr. Moule.....	347
To construct earth closets and their use.....	348
Table, showing the effect produced on the quantity of the crop by equal quantities of different manures applied to the same soil.....	351
Table, showing the comparative increase of corn by different fertilizers.....	350
Moisture absorbed by different manures.....	351
Table, showing the number of loads of manure and the number of heaps to each load required to each acre, the heaps at given distances apart.....	352
Weight of manure per cubic foot.....	353

	PAGE
Load of manure, how much it is.....	353
To find the number of loads of manure required to the acre for a given number of lbs. per square foot.....	354
MANURES, ARTIFICIAL.....	354
Analysis of Mape's improved superphosphate of lime.....	358
Analysis of Coe's superphosphate.....	358
Analysis of Deburg's bone meal.....	358
Analysis of bone dust.....	359
Analysis of fish guano.....	359
Prices of standard fertilizers.....	360
Average composition per cent. and per ton of various agricultural products.....	361
MARKING GOODS.....	247
MARKET, GARDENING FOR.....	446
MASONRY, Illustrated.....	276
To find the number of perches in stone walls.....	276
Brick-work.....	277
Dimensions of bricks.....	277
To find the number of bricks in a wall.....	277
Laths.....	278
MATERIALS, STRENGTH OF.....	264
Tensile strength.....	264
Table, showing the weight in lbs. necessary to tear asunder one square inch of various materials.....	264
To find the tensile strength.....	265
Table, showing the strength of iron-wire rope and hempen cable	266
To find the strength of cables.....	266

	PAGE
To find the strength of ropes and hawsers.....	267
Table, showing what weight hemp rope will bear with safety...	267
Strength of metal and wooden rods.....	267
Hempen cords	268
Lateral or transverse strength.....	268
Table, showing the transverse strength of timber.....	268
Tables, showing the transverse strength of iron.....	269
To find the transverse strength when the bar or beam is fixed at one end, and the load applied at the other.....	269
When the bar or beam is fixed at both ends and the weight applied in the middle	270
When the bar or beam is supported at both ends, and the weight applied in the middle.....	270
Table, showing the resistance of materials to crushing.....	271
Strength of ice.....	271
MATHEMATICAL FORMS, DEFINITIONS of, Illustrated.....	292
MEASUREMENT OF LAND (see land, measurement of), Illustrated....	43
" HAY (see hay, measurement of), Illustrated.....	51
" GRAIN IN GRANARIES, Illustrated.....	60
" TIMBER (see timber measurement), Illustrated... ..	61
MEASURES AND WEIGHTS, TABLES OF U. S. (see weights and mea- sures).....	145
MEASURES OF CAPACITY COMPARED.....	163
Tables of English weights and measures	163
MECHANICAL POWERS, Illustrated.....	279
The lever, Illustrated.....	279
The inclined plane, Illustrated.....	282

	PAGE
The wheel and the axle.....	284
The wedge, Illustrated.....	286
The screw, Illustrated.....	288
The pulley, Illustrated.....	290
METALS, FUSING HEAT OF.....	121
METRIC SYSTEM OF WEIGHTS AND MEASURES.....	173
" " origin and history of.....	173
" " countries that have adopted the.....	173
" " act of Congress authorizing.....	174
" " formation of tables.....	176
" " table of linear measure.....	177
" " table of square measure.....	178
" " table of cubic or solid measure.....	179
" " table of dry and liquid measure.....	179
" " table of weights.....	180
" " table of angles.....	181
" " tables of equivalents.....	181
MILK, PROPERTIES AND COMPOSITION OF.....	387
Analysis of milk.....	387
Table, showing the effects of various degrees of heat in making new milk cream.....	387
Analysis of the milk of different animals.....	390
MISCELLANEOUS WEIGHTS.....	153
MIXING PAINTS.....	460
MONEY (see United States Money).....	145
MORTAR, BROWN, FOR MASONRY, BRICK-WORK, &c.....	463

N.

NUTRITIVE VALUE OF CERTAIN CROPS.....	PAGE 190
Table, showing the nutritive value of certain crops.....	190

O.

OIL, per cent. in different seeds, grain, &c.....	191
Table, showing the per cent. of oil in different seeds, grain, &c...	191
OSAGE ORANGE—hedge plants.....	130

P.

PAINTING.....	460
House painting.....	460
Mixing paints.....	463
A beautiful white paint.....	463
A pure white paint.....	464
Common white paint.....	464
For knotting.....	464
Common flesh color.....	464
Fine flesh color.....	464
A beautiful color for carriages.....	464
Cream color.....	464
Pearl gray.....	464
Fawn color.....	464
Blue.....	465
Buff.....	465
Straw.....	465
Drab.....	465
Steel.....	465

	PAGE
Purple.....	465
Violet.....	465
French gray.....	465
Silver.....	465
Gold.....	465
Dark chestnut.....	466
Salmon.....	466
Peach blossom.....	466
Drab.....	466
Lead.....	466
Chocolate.....	466
Dark red.....	466
Orange.....	466
Bright yellow.....	466
Dark yellow.....	467
Light yellow.....	467
Olive green.....	467
Light green.....	467
Grass green.....	468
Invisible green.....	468
Bronze.....	468
Imitation of gold.....	468
Tar paint.....	468
Paint driers.....	468
Milk paint.....	469
Lime whitewash.....	469
Italian marble.....	470

	PAGE
Imitation granite.....	471
Oak varnish.....	471
Oil varnish.....	471
Varnish for pictures.....	471
Varnish for unpainted wood.....	472
Waterproof varnish for cloth, &c.....	472
PENDULUMS (see time, seasons, &c.), Illustrated.....	31
PLANK MEASURE.....	62
Table, showing the contents (board measure) of planks of various dimensions.....	67
PLANTS (see hedge plants).....	130
PORK, RELATION OF CORN TO.....	194
Table, showing the price of pork per lb., at different prices per bushel for corn... ..	194
To find the price of pork per lb., the price of corn being given..	195
To find the price of corn, the price of pork being given.....	195
POST AND RAIL FENCE (see fences), Illustrated.....	128
POST AND BOARD " (see fences).....	128
POWERS, THE MECHANICAL, Illustrated.....	279
PRACTICAL READER, TO THE.....	11
PREFACE	7
PRESSURE OF EARTH AGAINST WALLS.....	255
PRINTING, FACTS ABOUT.....	260
The different types used in book printing.....	260
The number of <i>ems</i> made by different type.....	261
Press-work.....	262
Sizes of books.....	262

	PAGE
Table, showing the number of leaves and pages from the folding of a sheet.....	263
PROPERTIES AND COMPOSITION OF MILK, BUTTER, &c.....	387
PROPORTION OF ALCOHOL IN LIQUORS.....	190
“ WEIGHT TO BULK OF VARIOUS SUBSTANCES.....	193
PULLEY, THE, Illustrated.	290

R.

RAIL FENCE, Illustrated.....	126
RAIN, AVERAGE FALL OF (see temperature and average fall of rain)..	37
RELATIVE VALUE OF GOLD AND CURRENCY.....	243
Table, showing the greenback value of \$1 at the different quotations of gold.....	243
Highest quotation of gold in New York during the civil war..	244
“ “ “ “ Richmond “ “ “ “ ..	244
English bonds and consols, explanation of.....	244
“Selling Short,” explanation of.....	245
“Seller’s Option,” “ “ ..	245
“Buyer’s Option,” “ “ ..	245
Stock Quotations, “ “ ..	246
“Bull,” commercial definition of.....	245
“Bear,” “ “ “ ..	245
“Stag,” “ “ “ ..	245
RODS, LIGHTNING.....	251
To construct a lightning-rod	251
ROOTS, SQUARE AND CUBE, TABLE OF.....	303

CONTENTS.

499

	PAGE
ROTATION OF CROPS.....	378
" field crops.....	385
" garden crops....	386

S.

SCANTLING MEASURE.....	72
Table, showing the contents (board measure) of scantling of various dimensions.....	72
SCREW, THE, Illustrated.....	288
SEASONS, TIME, &c. (see time), Illustrated.....	19
SEEDS, WEIGHT OF, as established by the Legislatures of the different States	189
SEEDS, OIL PER CENT. IN.....	191
" quantity of, to sow or plant per acre, &c.....	192
SHEEP, TO FIND THE AGE OF... ..	208
SOILING CATTLE.....	401
Experiments by the author.....	402
Arrangement of crops for soiling.....	405
Arguments in favor of soiling.....	409
SOILS, EXHAUSTION OF.....	320
Table, showing the organic substances removed from the soil in 1000 lbs. each of the various crops.....	321
Table, showing the inorganic matter removed from the soil in 1000 lbs. each of the various crops... ..	321
Table, showing the <i>kinds</i> of inorganic matter removed from the soil in 1000 lbs. each of the various crops.....	322

	PAGE
Analysis of the hop, showing the elements it removes from the soil.....	323
Table, showing amount of inorganic matter removed from the soil by ten bushels of grain.....	324
SOILS.	311
Classification of soils.....	312
To analyze soils.....	313
General results of analytical examinations of soils.....	316
Table, showing the composition in 1000 parts of different soils..	317
Analytic table of three very fertile soils.....	317
Analytic table of arable lands of great fertility.....	318
Depth of soil—its importance.....	318
Table, showing the weight per cubic foot of the different kinds of earth.....	319
SOLDERS	473
Lead solder.....	473
Tin solder.....	473
SOLID MATTER AND WATER IN ARTICLES OF DIET.....	188
Table, showing the proportion of solid matter and water in 100 parts each of the various articles of diet.....	188
SPECIFIC GRAVITY (see gravity), Illustrated.....	182
SQUARE MEASURE, table.....	165
SQUARES AND SQUARE ROOTS, table of, from 1 to 1000.....	303
STEAMING FOOD FOR STOCK.....	415
Report of the Department of Agriculture.....	416
How to make a steaming apparatus.....	421
Prindle's Agricultural Steamer and Cauldron (Illustrated).....	423

CONTENTS.

501

	PAGE
Advantages of cooked food.	425
STEAM CULTIVATION, Illustrated.....	450
Advantages claimed.....	452
Report of the Royal Agricultural Society.....	455
STEAM POWER, rules for computing.....	140
STOCK, STEAMING FOOD FOR...	415
STOCK QUOTATIONS.....	246
STRENGTH, HUMAN.....	135
STRENGTH OF MATERIALS (see materials, strength of).....	204
SUCCESS IN BUSINESS.....	246
Short credits.....	246
Small profits.....	246
Economy in expense.....	247
Marking goods.....	247
SURVEYORS' MEASURE, table.....	168

T.

TEMPERATURE AND FALL OF RAIN, AVERAGE OF	37
Table, showing the average temperature of the four seasons at points on the Pacific and Atlantic coasts, and the interior of this continent.....	37
Periodical rains, region of.....	38
Frequent rains, ".....	39
Scanty rains, ".....	39
Table, showing the latitude and longitude, the elevation above the level of the sea, the mean annual temperature, and the average fall of rain in various places in the United States....	49
TILE DRAINING (see draining).....	362

	PAGE
TIMBER, MEASUREMENT OF, Illustrated.....	61
Board measure.....	62
To ascertain the contents (board measure) of boards, scantling, and plank.....	62
Table, showing the contents of inch boards from 6 inches to 30 broad, and from 4 to 24 feet long.....	63
Square timber.....	65
To measure square timber.....	65
Plank measure.....	62
Table, showing the contents (board measure) of planks of various dimensions.....	67
Round and square timber.....	64
To measure round timber, Illustrated.	65
Logs reduced to inch-board measure.....	70
Table, showing the number of feet (board measure) of inch boards contained in round saw-logs of various dimensions....	71
TIME, SEASONS, &c., Illustrated.....	19
To reduce longitude to time.....	19
Time, apparent and mean.....	21
To ascertain the length of the day and night.....	21
Pendulums, Illustrated....	31
To find the length of a pendulum for a given number of vibra- tions per minute.....	31
To find the vibrations per minute, the length of the pendulum being given.....	32
Measure of time, table, Illustrated.....	25
Division of the calendar year.....	26

	PAGE
Old style (O. S.) and new style (N. S.).....	27
Decade, what period it is.....	27
Century, “ “ “	27
Lunar Cycle, what it is.....	27
Golden Number, what it is.....	27
Solar Cycle, what it is.....	28
To find the lunar cycle or golden number.....	28
Table, showing the number of days from any day in the month to the same day in any other.....	28
Table, finding the number of days between two dates.....	29
Table, showing the planets, &c., in the solar system.....	27
Distance of the planets, and size compared with the earth.....	32

U.

UNITED STATES BONDS, explanation of.....	242
Five-Twenties.....	242
Ten-Forties.....	242
Seven-Thirties.....	242
Six per cents. of '81.....	243
UNITED STATES MONEY, table.....	145
“ “ Gold coin.....	146
“ “ Silver coin.....	147
“ “ Copper coin.....	148
Alloy of Gold and Silver.	147

V.

VELOCITY, table of.....	188
-------------------------	-----

W.

	PAGE
WAGES.....	224
Table of wages at \$3 to \$25 per month of 26 working days....	224
WAGON-BEDS, CAPACITY OF, Illustrated.....	82
To find the contents of wagon-beds.....	82
WALLS, PRESSURE OF EARTH AGAINST.....	255
WATER RAM, Illustrated.....	103
WEATHER.....	33
Table, for telling the weather through all the lunations of the year	33
WEDGE, THE, Illustrated.....	286
WEIGHTS AND MEASURES, TABLES OF U. S., Illustrated.....	145
Long measure.....	167
Hair's breadth.....	168
Gunter's chain.....	168
Ropes and cables.....	168
Geographical and nautical measure.....	169
Miscellaneous long measures.....	168
Measures of circles.....	23
Measures of surfaces.....	165
Land measure.....	43
Paper measure.....	263
Liquid measure.....	160
Standard gallon measure.....	161
Dry measure.....	162
Standard bushel measure.....	162
Imperial or British bushel.....	163
Miscellaneous dry measures.....	163

	PAGE
Measure of weights, avoirdupois.....	153
Troy weight.....	156
Troy weight reduced to avoirdupois.....	157
Diamond measure.....	157
Measure of time.....	25
Measure of value.....	145
Standard of gold and silver.....	148
Miscellaneous weights and measures.....	153
Heaping measure.....	163
Barrel measure.....	153
Ton weight and ton measure.....	172
A sack of wool.....	155
A pack of wool.....	155
A truss of hay.....	56
A load of hay.....	56
A balé of hay.....	56
A firkin of butter.....	155
A bale of cotton.....	155
WEIGHT, COMPUTE, OF CATTLE, Illustrated.....	209
WEIGHT OF LEAD PIPE.....	112
WEIGHTS OF GRAIN, SEEDS, &c.....	189
Table, showing the weight of grain, seeds, &c., as established by the Legislatures of the different States.....	189
WEIGHT OF SQUARE ROLLED IRON.....	273
Table, showing the weight of square rolled iron from $\frac{1}{16}$ inch to 12 inches, and 1 foot long.....	273
WEIGHT OF ROUND ROLLED IRON.....	275

352
70
—
282

	PAGE
Table, showing the weight of round rolled iron from $\frac{1}{4}$ inch to 12 inches diameter and 1 foot long	275
WEIGHT, PROPORTION OF, TO BULK OF VARIOUS SUBSTANCES.....	193
Table, showing the weight per cubic foot of various substances, and the number of cubic feet required to make a ton of each..	193
WHEAT, DEPTH OF SOWING.	193
WHEEL AND AXLE.....	284
WHITEWASH	469
WIND.....	35
Table, showing the force and velocity of wind.....	36
To find the force of wind acting against a surface.....	35
WIRE FENCES.....	134
WOOD MEASURE, Illustrated.....	62
To ascertain the number of cords in a given pile of wood.	62

MANN
Spec
Coll
S
SDI
C86
1868



